

RINGS, MODULES AND LINEAR ALGEBRA

N. P. STRICKLAND

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike license.



1. INTRODUCTION

At the end of the course, you should be able to prove the following:

- (1) Let G be an Abelian group of order p^3 , where p is prime. Then G is isomorphic to either Z_{p^3} , $Z_{p^2} \times Z_p$ or $Z_p \times Z_p \times Z_p$.
- (2) Let A be a 3×3 matrix of rational numbers satisfying $(A + I)^3 = 0$. Then A is conjugate to one of the following three matrices:

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & -1 \end{bmatrix} \quad \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \quad \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

(In other words, there is an invertible matrix P such that PAP^{-1} is one of the three matrices listed.)

- (3) Let $f: \mathbb{R} \rightarrow \mathbb{C}$ satisfy the differential equation $f'''' + f = 0$. Then $f(x) = ae^x + be^{-x} + ce^{ix} + de^{-ix}$ for some constants a, b, c, d .

The remarkable thing is that all these problems can be addressed using essentially the same ideas: the theory of modules over a Euclidean domain.

2. RINGS AND FIELDS

A *commutative ring* is a set R of things that can be added, negated and multiplied in a sensible way to get new elements of R . More precisely, we require that the following axioms be satisfied:

- (a) If $a, b \in R$ then $a + b \in R$. [closure under addition]
- (b) There is an element $0 \in R$ such that $a + 0 = a$ for all $a \in R$. [additive identity]
- (c) For each element $a \in R$ there is an element $-a \in R$ such that $a + (-a) = 0$. [additive inverses]
- (d) $a + (b + c) = (a + b) + c$ for all $a, b, c \in R$. [associativity of addition]
- (e) $a + b = b + a$ for all $a, b \in R$. [commutativity of addition]
- (f) If $a, b \in R$ then $ab \in R$. [closure under multiplication]
- (g) There is an element $1 \in R$ such that $1a = a$ for all $a \in R$. [multiplicative identity]
- (h) $a(bc) = (ab)c$ for all $a, b, c \in R$. [associativity of multiplication]
- (i) $ab = ba$ for all $a, b \in R$. [commutativity of multiplication]
- (j) $a(b + c) = ab + ac$ for all $a, b, c \in R$. [distributivity]

Strictly speaking, we should say that a ring is a set R together with a definition of addition, negation and multiplication such that the axioms hold. In all the examples that we will consider, there is a unique obvious way to define these operations.

We will not consider noncommutative rings in this course, so we will just use the word “ring” to mean “commutative ring”. We will use without comment various standard consequences of the axioms, such as the facts that $-(-a) = a$, $0.a = 0$ and $(-1).a = -a$.

Definition 2.1. [defn-domain]

A ring R is an *integral domain* if $1 \neq 0$, and whenever $a, b \neq 0$ we also have $ab \neq 0$.

Definition 2.2. An element a in a ring R is *invertible* if there is an element $b \in R$ such that $ab = 1$. If so, then this element b is unique and we call it a^{-1} . A *field* is a ring K such that $1 \neq 0$ and every nonzero element is invertible. Every field is an integral domain.

Example 2.3. [eg-Z-ring]

The set \mathbb{Z} of integers is a ring. If we add, subtract or multiply any two integers, we get another integer, so axioms (a), (c) and (f) hold. The numbers 0 and 1 are integers so axioms (b) and (g) hold. The remaining axioms are familiar properties of addition, subtraction and multiplication. It is also clear that \mathbb{Z} is an integral domain.

Example 2.4. [eg-N-not-ring]

The set \mathbb{N} of natural numbers is not a ring, because $0 \notin \mathbb{N}$, so there is no additive identity, in other words axiom (b) does not hold. Moreover, if $n \in \mathbb{N}$ then $-n \notin \mathbb{N}$, so \mathbb{N} does not have additive inverses.

Example 2.5. [eg-even-not-ring]

The set $2\mathbb{Z}$ of even integers is not a ring, because it does not contain the multiplicative identity element 1.

Example 2.6. [eg-QRC]

The set \mathbb{Q} of rational numbers, the set \mathbb{R} of real numbers, and the set \mathbb{C} of complex numbers, are all fields.

Example 2.7. [eg-irrational]

The set $X = \mathbb{R} \setminus \mathbb{Q}$ of irrational numbers is not a ring. Indeed, we have $\pi \in X$ and $-\pi \in X$ but $0 = \pi + (-\pi) \notin X$, so X is not closed under addition. Moreover, $\sqrt{2} \in X$ but $\sqrt{2} \cdot \sqrt{2} = 2 \notin X$, so X is not closed under multiplication. Even more obviously, X contains neither 0 nor 1, so it does not have an additive identity or a multiplicative identity.

Example 2.8. [eg-Zn]

For any natural number n , the set $\mathbb{Z}_n = \{\bar{0}, \bar{1}, \dots, \overline{n-1}\}$ of integers modulo n is a ring. For example, we have

$$\begin{aligned}\mathbb{Z}_4 &= \{\bar{0}, \bar{1}, \bar{2}, \bar{3}\} \\ \bar{2} + \bar{3} &= \bar{5} = \bar{1} \\ \bar{1} - \bar{2} &= \overline{-1} = \bar{3} \\ \bar{2} \cdot \bar{3} &= \bar{6} = \bar{2}.\end{aligned}$$

Note that \mathbb{Z}_4 is not an integral domain, because $\bar{2} \neq 0$ but $\bar{2} \cdot \bar{2} = 0$. In general, it can be shown that when n is prime, \mathbb{Z}_n is a field (and thus an integral domain), but when n is not prime, \mathbb{Z}_n is neither a field nor an integral domain.

Example 2.9. [eg-poly]

We write $\mathbb{Z}[x]$ for the set of all polynomials with integer coefficients. For example, $7x^3 - 22x + 3$ and x^{1001} are elements of $\mathbb{Z}[x]$ but $(x+1)/(x-1)$ and $x^2 - 1/2$ and $x - x^{-1}$ are not. The general form of an element of $\mathbb{Z}[x]$ is $f(x) = a_0 + a_1x + \dots + a_nx^n$ for some integer $n \geq 0$ and integers a_0, \dots, a_n . Integers are polynomials of degree zero, so $\mathbb{Z} \subseteq \mathbb{Z}[x]$. The usual operations of addition, multiplication and negation of polynomials make $\mathbb{Z}[x]$ into a ring.

More generally, given any ring R we can consider the ring $R[x]$ of polynomials with coefficients in R . For example, we can consider $f(x) = \bar{2}x^2 + \bar{3} \in \mathbb{Z}_6[x]$ and $g(x) = \bar{3}x + \bar{2} \in \mathbb{Z}_6[x]$ and we find that

$$\begin{aligned}f(x)g(x) &= \bar{6}x^3 + \bar{4}x^2 + \bar{9}x + \bar{6} \\ &= \bar{4}x^2 + \bar{3}x.\end{aligned}$$

This gives us rings $\mathbb{Z}[x] \subseteq \mathbb{Q}[x] \subseteq \mathbb{R}[x] \subseteq \mathbb{C}[x]$. We can also use more than one variable; for example, we have a ring $\mathbb{Q}[x, y, z]$ containing elements like $(x^2 + y^2 + z^2)/4$ or $1 + xyz$ (but not x/y or $\sqrt{2}x$ or e^{x+y}).

Example 2.10. [eg-diffop]

Let D denote the operation of differentiation with respect to t , so $D(t^3) = 3t^2$, $D(\sin(t)) = \cos(t)$, $D^3(f(t)) = f'''(t)$ and so on. Using this we can build more complicated operators like $(D^2 + 2D + 3)(f(t)) = f''(t) + 2f'(t) + 3f(t)$ and so on. By a *differential operator* we mean an operation of the form $a_0 + a_1D + \dots + a_nD^n$ for some $n \geq 0$ and some list of coefficients $a_i \in \mathbb{R}$. The set of differential operators is the polynomial ring $\mathbb{R}[D]$; it is essentially the same as $\mathbb{R}[x]$ except for the notation, and the fact that the elements are interpreted as operators rather than functions.

Remark 2.11. [rem-diffop]

Here we are only considering linear ordinary differential operators with constant coefficients. For more serious work on differential equations one needs to work with more general operators, which generally form noncommutative rings.

Example 2.12. [eg-Zplocal]

Let p be a prime number. Let $\mathbb{Z}_{(p)}$ be the set of rational numbers x that can be written in the form a/b , where a and b are integers and b is not divisible by p . For example, $123/101$ and $-4/12 = (-1)/3$ and $8 = 8/1$ are elements of $\mathbb{Z}_{(2)}$, but $5/6$ and $1/8$ are not. As any integer n can be written as $n/1$ and 1 is not divisible by p , we see that $\mathbb{Z} \subseteq \mathbb{Z}_{(p)}$. Now suppose that $x, y \in \mathbb{Z}_{(p)}$, so we can write $x = a/b$ and $y = c/d$ for some integers a, b, c and d , where b and d are not divisible by p . As p is prime this means that bd is also not divisible by p . We have

$$\begin{aligned}x + y &= (ad + bc)/(bd) \\xy &= (ac)/(bd) \\-x &= -a/b.\end{aligned}$$

As $ad + bc$, ac , $-a$, b and bd are integers, and bd and b are not divisible by p , this means that $x + y$, xy and $-x$ lie in $\mathbb{Z}_{(p)}$. Thus $\mathbb{Z}_{(p)}$ is a subring of \mathbb{Q} , called the ring of integers localized at p . (There is a long story coming from algebraic geometry that explains why the word “localized” is appropriate.)

Example 2.13. [eg-gaussian]

We write $\mathbb{Z}[i]$ for the set of complex numbers of the form $a + bi$, where a and b are integers (possibly zero). Thus 7 , $6 - 4i$ and $12i$ are elements of $\mathbb{Z}[i]$, but $2/3$ and $1 - i/5$ are not. Note that

$$\begin{aligned}(a + bi) + (c + di) &= (a + c) + (b + d)i \\(a + bi)(c + di) &= (ac - bd) + (ad + bc)i \\-(a + bi) &= (-a) + (-b)i.\end{aligned}$$

It follows easily that $\mathbb{Z}[i]$ is closed under addition, multiplication and negation, so it is a subring of \mathbb{C} . The elements of $\mathbb{Z}[i]$ are called *Gaussian integers*.

Exercises

Exercise 2.1. [ex-which-ring]

Which of the following are commutative rings?

- R_0 is the set of polynomials $f(x) \in \mathbb{R}[x]$ such that $f(-x) = f(x)$.
- R_1 is the set of polynomials $f(x) \in \mathbb{R}[x]$ such that $f(-x) = -f(x)$.
- R_2 is the set of 2×2 matrices over \mathbb{R} , with the usual definition of matrix multiplication.
- R_3 is the set of 2×2 matrices over \mathbb{R} , with multiplication given by the definition

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a' & b' \\ c' & d' \end{bmatrix} = \begin{bmatrix} aa' & bb' \\ cc' & dd' \end{bmatrix}.$$

- R_4 is the set of vectors in \mathbb{R}^3 , with multiplication given by the cross product:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \times \begin{bmatrix} x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} yz' - y'z \\ zx' - z'x \\ xy' - x'y \end{bmatrix}.$$

Exercise 2.2. [ex-typical-elts]

Choose two typical elements a and b of the ring $\mathbb{Z}_{(5)}$. Find $a + b$ and ab and check that they lie in $\mathbb{Z}_{(5)}$. Repeat this for the rings $\mathbb{Z}[i]$, $\mathbb{Q}[x, y]$ and \mathbb{Z}_{12} .

Exercise 2.3. [ex-six-local]

Let R be the set of rational numbers that can be written in the form a/b , where b is not divisible by 6. (We would call this $\mathbb{Z}_{(6)}$ if 6 were prime, which of course it isn't). Prove that R is *not* a subring of \mathbb{Q} .

Exercise 2.4. [ex-field-domain]

Let K be a field; prove that K is an integral domain.

Exercise 2.5. [ex-PX-ring]

Let X be a set. Let R be the set of all subsets of X , and define addition and multiplication of subsets as follows.

$$\begin{aligned} A + B &= (A \cup B) \setminus (A \cap B) \\ &= \{x \in X \mid x \in A \text{ or } x \in B \text{ but not both.}\} \\ AB &= A \cap B. \end{aligned}$$

For any $A \in R$, we define a function $\chi_A: X \rightarrow \mathbb{Z}_2$ by

$$\chi_A(x) = \begin{cases} \bar{1} & \text{if } x \in A \\ \bar{0} & \text{if } x \notin A \end{cases}$$

- Check that $A + \emptyset = A$ and $A + A = \emptyset$.
- Show that $\chi_{A+B}(x) = \chi_A(x) + \chi_B(x)$ and $\chi_{AB}(x) = \chi_A(x)\chi_B(x)$.
- Show that if $\chi_A = \chi_B$ then $A = B$.
- Prove that the definitions above make R into a commutative ring. (You may wish to use (b) and (c) to help check some of the axioms.)

3. MODULES

Definition 3.1. [defn-module]

Let R be a ring. A *module* over R is a set M of things with a definition of $m + n$ for all $m, n \in M$ and a definition of am for all $a \in R$ and $m \in M$ such that the following axioms are satisfied:

- If $m, n \in M$ then $m + n \in M$. [closure under addition]
- There is an element $0 \in M$ such that $m + 0 = m$ for all $m \in M$. [additive identity]
- For each $m \in M$ there is an element $-m \in M$ such that $m + (-m) = 0$. [additive inverses]
- $m + (n + p) = (m + n) + p$ for all $m, n, p \in M$. [associativity of addition]
- $m + n = n + m$ for all $m, n \in M$. [commutativity of addition]
- If $a \in R$ and $m \in M$ then $am \in M$. [closure of M under multiplication by R]
- $1.m = m$ for all $m \in M$.
- $(ab)m = a(bm)$ for all $a, b \in R$ and $m \in M$. [associativity of multiplication]
- $(a + b)m = am + bm$ for all $a, b \in R$ and $m \in M$. [left distributivity of multiplication]
- $a(m + n) = am + an$ for all $a \in R$ and $m, n \in M$. [right distributivity of multiplication]

Remark 3.2. [rem-additive-group]

Note that axioms (a) to (e) say that M is in particular an Abelian group under addition.

Example 3.3. [eg-free-module]

Let R be any ring, and let d be a natural number. We then write R^d for the set of d -tuples (x_1, \dots, x_d) with

$x_1, \dots, x_d \in R$. We make R^d into a module over R by defining

$$\begin{aligned}(x_1, \dots, x_d) + (y_1, \dots, y_d) &= (x_1 + y_1, \dots, x_d + y_d) \\ a(x_1, \dots, x_d) &= (ax_1, \dots, ax_d).\end{aligned}$$

It is straightforward to check that the axioms are satisfied. In particular, the case $d = 1$ says that we can regard R as a module over itself.

If R is a field, then an R -module is just a vector space over R . Modules are just the natural generalization of vector spaces defined over arbitrary rings rather than just fields. It is a basic fact of linear algebra that if K is a field and V is a vector space over K with a finite spanning set, then V is isomorphic to K^d for some integer d , called the *dimension* of V . The situation for modules over non-fields is more complicated; a module is usually not isomorphic to R^d for any d . The next simplest case after fields is when R is a Euclidean domain, and most of the course will be devoted to the study of modules over such rings.

Proposition 3.4. [prop-Z-module]

A \mathbb{Z} -module is just an Abelian group. More precisely, if M is an Abelian group (with the group operation written as addition) then there is a unique way to define am for all $a \in \mathbb{Z}$ and $m \in M$ such that axioms (f) to (j) hold, making M a \mathbb{Z} -module.

Sketch proof. Rather than giving a complete proof of this, we will give an outline of the argument with examples.

The basic idea is very simple. We just define

$$\begin{aligned}3m &= m + m + m \\ -5m &= -(m + m + m + m + m) = (-m) + (-m) + (-m) + (-m) + (-m)\end{aligned}$$

and so on. This defines multiplication (of integers by group elements) in terms of addition and negation of group elements. We actually have no choice about these definitions if we want the axioms to be satisfied: as $3 = 1 + 1 + 1$, axiom (i) says we must have $3m = (1 + 1 + 1)m = 1m + 1m + 1m$, and axiom (g) says that $1m = m$ so we must have $3m = m + m + m$, and so on.

We now need to check that axioms (f) to (j) are satisfied. Axioms (f) and (g) are immediate. The remaining axioms are easy to check when a and b are nonnegative: for example

$$\begin{aligned}2(3m) &= 2(m + m + m) \\ &= (m + m + m) + (m + m + m) \\ &= m + m + m + m + m + m \\ &= (2 \times 3)m \\ 2m + 3m &= (m + m) + (m + m + m) \\ &= m + m + m + m + m \\ &= (2 + 3)m \\ 3(m + n) &= (m + n) + (m + n) + (m + n) \\ &= (m + m + m) + (n + n + n) \\ &= 3m + 3n.\end{aligned}$$

If we allow a or b to be negative then there are quite a few more cases to check depending on the various possible combinations of signs, but they are all quite straightforward. For example

$$\begin{aligned}5m + (-3)m &= (m + m + m + m + m) + ((-m) + (-m) + (-m)) \\ &= m + m + (m + (-m)) + (m + (-m)) + (m + (-m)) \\ &= m + m \\ &= (5 + (-3))m.\end{aligned}$$

□

We will next give an example involving differential operators. To avoid annoying technicalities, it is best to restrict attention to functions that can be differentiated as many times as we like. We therefore introduce the following definition.

Definition 3.5. [defn-smooth]

A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is *smooth* if the n 'th derivatives $f^{(n)}(t)$ are defined and continuous everywhere on \mathbb{R} for all $n \geq 0$. In particular, the function $f = f^{(0)}$ itself must be defined and continuous everywhere.

For example, $\sin(t)$, $\cos(t)$, e^t , t^2 and so on are smooth. However, the functions $1/t$ and $\log(t)$ are not defined at $t = 0$, so they are not smooth. The function $f(t) = |t|$ is defined and continuous everywhere and $f'(t) = -1$ for $t < 0$ and $f'(t) = 1$ for $t > 0$ but $f'(0)$ is undefined so f is not smooth. Similarly, if $g(t) = t^{1/3}$ then $g'(t) = t^{-2/3}/3$, which is undefined at $t = 0$ so g is not smooth.

We write $C^\infty(\mathbb{R}, \mathbb{R})$ for the set of all smooth functions from \mathbb{R} to \mathbb{R} . If f and g are smooth and a is constant then one can show that $f + g$ and af are smooth. It follows that $C^\infty(\mathbb{R}, \mathbb{R})$ is a vector space over \mathbb{R} . Similarly, the set $C^\infty(\mathbb{R}, \mathbb{C})$ of smooth functions from \mathbb{R} to \mathbb{C} is a vector space over \mathbb{C} .

Example 3.6. [eg-smooth-module]

$C^\infty(\mathbb{R}, \mathbb{R})$ is a module over the ring $\mathbb{R}[D]$ of differential operators. For an operator $p(D) = a_0 + a_1D + \dots + a_nD^n$ and a smooth function $f(t)$, the product $p(D)f$ is defined by the usual rule

$$p(D)f = a_0f + a_1f' + a_2f'' + \dots + a_nf^{(n)}.$$

One can check the axioms directly, or use a more abstract approach discussed in the next section. Similarly, $C^\infty(\mathbb{R}, \mathbb{C})$ is a module over $\mathbb{C}[D]$.

Example 3.7. [eg-diffop-calc]

$$\begin{aligned} (1 + D + D^2).(1 + t + t^2) &= (1 + t + t^2) + (1 + t + t^2)' + (1 + t + t^2)'' \\ &= (1 + t + t^2) + (1 + 2t) + (2) \\ &= 4 + 3t + t^2. \end{aligned}$$

Example 3.8. [eg-t-sint]

Consider the function $f(t) = t \sin(t)$; I claim that $(D^2 + 1)^2 f = 0$. Indeed, we have

$$\begin{aligned} f'(t) &= \sin(t) + t \cos(t) \\ f''(t) &= 2 \cos(t) - t \sin(t) \\ ((D^2 + 1)f)(t) &= f(t) + f''(t) = 2 \cos(t) \end{aligned}$$

We also have $\cos'(t) = -\sin(t)$ and so $\cos''(t) = -\cos(t)$ so $(D^2 + 1) \cos = 0$. It follows that $(D^2 + 1)^2 f = (D^2 + 1)(2 \cos) = 0$.

Example 3.9. [eg-two-rates]

Consider a function of the form $f(t) = e^{\lambda t} + e^{\mu t}$. I claim that

$$(p(D)f)(t) = p(\lambda)e^{\lambda t} + p(\mu)e^{\mu t}.$$

Indeed, we have

$$\begin{aligned} f'(t) &= \lambda e^{\lambda t} + \mu e^{\mu t} \\ f''(t) &= \lambda^2 e^{\lambda t} + \mu^2 e^{\mu t} \end{aligned}$$

and more generally $f^{(k)}(t) = \lambda^k e^{\lambda t} + \mu^k e^{\mu t}$ (as one can easily check by induction). If $p(D) = a_0 + a_1D + \dots + a_mD^m$ then we have

$$\begin{aligned} (p(D)f)(t) &= a_0(e^{\lambda t} + e^{\mu t}) + a_1(\lambda e^{\lambda t} + \mu e^{\mu t}) + \dots + a_m(\lambda^m e^{\lambda t} + \mu^m e^{\mu t}) \\ &= (a_0 + a_1\lambda + \dots + a_m\lambda^m)e^{\lambda t} + (a_0 + a_1\mu + \dots + a_m\mu^m)e^{\mu t} \\ &= p(\lambda)e^{\lambda t} + p(\mu)e^{\mu t}. \end{aligned}$$

Exercises

Exercise 3.1. [ex-Z-ten]

List all the elements of the Abelian group $\mathbb{Z}_2 \oplus \mathbb{Z}_5$. Find an element that has order 10.

Exercise 3.2. [ex-diffop-calc]

- Calculate $(1 + D + D^2/2 + D^3/6).t^3$. What do you notice? Can you guess a generalisation?
- Put $f(t) = e^{-t} \sin(t)$ so $f \in C^\infty(\mathbb{R}, \mathbb{R})$. Calculate $(D + 1)^2 f$.
- Put $g_k(t) = t^k e^t$, so $g_k \in C^\infty(\mathbb{R}, \mathbb{R})$. Calculate $(D - 1)g_k$ and thus $(D - 1)^k g_k$. (You may wish to try $k = 3$ first.)
- Put $f(t) = te^t$. Show that $(D^k f)(t) = (k + t)e^t$ for all $k \geq 0$ and thus that $(p(D)f)(t) = (p'(1) + p(1)t)e^t$.

Exercise 3.3. [ex-gaussian-poly]

Define $v(t) = e^{t^2/2}$, so $u \in C^\infty(\mathbb{R}, \mathbb{R})$. Let V be the set of functions of the form $f(t)v(t)$, where f is a polynomial. For example, the function $(1 + t + t^2)e^{t^2/2}$ is an element of V .

- Prove that V is an $\mathbb{R}[D]$ -submodule of $C^\infty(\mathbb{R}, \mathbb{R})$.
- Calculate $D^k v$ for $0 \leq k \leq 3$.
- Show that for all $k \geq 0$ there is a polynomial $p_k(t)$ of the form $t^k +$ lower terms such that $D^k v = p_k.v$.
- Show that if $q(D)$ is a nonzero element of $\mathbb{R}[D]$ then $q(D)v \neq 0$ (look at leading terms).
- Suppose that $f(t)$ is a polynomial of degree k , say $f(t) = at^k +$ lower terms. Prove by induction on k that $f v = q(D)v$ for some element $q(D) \in \mathbb{R}[D]$.
- Deduce that $V \simeq \mathbb{R}[D]$ as an $\mathbb{R}[D]$ -module.

4. MODULES OVER POLYNOMIAL RINGS

We next consider modules over $K[x]$, where K is a field. The upshot here is that the study of modules over $K[x]$ is essentially the same as the study of square matrices over K , or of endomorphisms of vector spaces over K .

We start with some comments about the process of “substituting a matrix into a polynomial”. Let K be a field, and let A be an $n \times n$ matrix over K . Using the usual matrix multiplication we can define A^2, A^3 and so on; all of these are again $n \times n$ matrices over K . Thus, given a polynomial $f(x) = a_0 + a_1 x + \dots + a_d x^d \in K[x]$ we can define another $n \times n$ matrix $f(A)$ by $f(A) = a_0 I + a_1 A + \dots + a_d A^d$.

Example 4.1. [eg-fA]

If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $f(x) = 7 + 6x + 5x^2$ then $A^2 = \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix}$ and so

$$\begin{aligned} f(A) &= 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + 6 \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + 5 \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix} \\ &= \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} + \begin{bmatrix} 6 & 12 \\ 18 & 24 \end{bmatrix} + \begin{bmatrix} 35 & 50 \\ 75 & 110 \end{bmatrix} \\ &= \begin{bmatrix} 48 & 62 \\ 93 & 141 \end{bmatrix}. \end{aligned}$$

Example 4.2. [eg-diag-poly]

Consider a diagonal matrix $A = \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}$. Then

$$A^2 = \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix} \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix} = \begin{bmatrix} \lambda^2 & 0 \\ 0 & \mu^2 \end{bmatrix},$$

and more generally it is not hard to see that

$$A^k = \begin{bmatrix} \lambda^k & 0 \\ 0 & \mu^k \end{bmatrix}.$$

(Exercise: prove this by induction.) It follows that

$$\begin{aligned} f(A) &= a_0 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + a_1 \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix} + \dots + a_d \begin{bmatrix} \lambda^d & 0 \\ 0 & \mu^d \end{bmatrix} \\ &= \begin{bmatrix} a_0 + a_1\lambda + \dots + a_d\lambda^d & 0 \\ 0 & a_0 + a_1\mu + \dots + a_d\mu^d \end{bmatrix} \\ &= \begin{bmatrix} f(\lambda) & 0 \\ 0 & f(\mu) \end{bmatrix}. \end{aligned}$$

More generally, if A is an $n \times n$ matrix with entries $\lambda_1, \dots, \lambda_n$ on the diagonal and zeros elsewhere, then $f(A)$ has entries $f(\lambda_1), \dots, f(\lambda_n)$ on the diagonal and zeros elsewhere.

Example 4.3. [eg-jordan-poly]

Consider the matrix $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. It is easy to check that

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & k+1 \\ 0 & 1 \end{bmatrix},$$

and thus that $A^k = \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$ for all k . It follows that

$$\begin{aligned} f(A) &= a_0 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + a_1 \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + \dots + a_d \begin{bmatrix} 1 & d \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} a_0 + \dots + a_d & a_1 + 2a_2 + \dots + da_d \\ 0 & a_0 + \dots + a_d \end{bmatrix}. \end{aligned}$$

Note that $f(1) = a_0 + \dots + a_d$. Note also that the derivative $f'(x)$ is given by $f'(x) = a_1 + 2a_2x + \dots + da_dx^{d-1}$, so that $f'(1) = a_1 + 2a_2 + \dots + da_d$. We can thus rewrite the above result as

$$f(A) = \begin{bmatrix} f(1) & f'(1) \\ 0 & f(1) \end{bmatrix}.$$

We next need to check that some things work out as they “ought” to when we substitute matrices into polynomials. (Recall that matrix multiplication is noncommutative, there are nonzero matrices whose square is zero, and numerous other funny things can happen; so we need to be on our guard.)

Proposition 4.4. [prop-func-calc]

Let A be an $n \times n$ matrix over a field K . Then for any two polynomials $f, g \in K[x]$ we have

$$\begin{aligned} (f + g)(A) &= f(A) + g(A) \\ (fg)(A) &= f(A)g(A). \end{aligned}$$

Proof. Suppose that $f(x) = \sum_i a_i x^i$ and $g(x) = \sum_j b_j x^j$. Then $(f + g)(x) = \sum_i c_i x^i$ where $c_i = a_i + b_i$, and

$$(fg)(x) = \left(\sum_i a_i x^i \right) \left(\sum_j b_j x^j \right) = \sum_{i,j} a_i b_j x^{i+j} = \sum_k d_k x^k,$$

where $d_k = \sum_{i=0}^k a_i b_{k-i}$.

Thus

$$\begin{aligned} (f + g)(A) &= \sum_i c_i A^i \\ &= \sum_i (a_i A^i + b_i A^i) \\ &= \sum_i a_i A^i + \sum_i b_i A^i \\ &= f(A) + g(A). \end{aligned}$$

Similarly

$$\begin{aligned}
(fg)(A) &= \sum_k d_k A^k \\
&= \sum_k \sum_{i=0}^k a_i b_{k-i} A^k \\
&= \sum_k \sum_{i=0}^k (a_i A^i)(b_{k-i} A^{k-i}) \\
&= \sum_i \sum_j (a_i A^i)(b_j A^j) \\
&= \sum_i a_i A^i \sum_j b_j A^j \\
&= f(A)g(A).
\end{aligned}$$

□

We are now ready to construct some modules over $K[x]$.

Construction 4.5. [cons-MA]

Let A be an $n \times n$ matrix over a field K ; we will use this to define a module M_A over $K[x]$. The elements of M_A are just the vectors $v = (v_1, \dots, v_n)$ of length n over K , so $M_A = K^n$ as a set. Addition and subtraction of vectors is defined in the usual way. All that is left is to define the product fv for $f \in K[x]$ and $v \in K^n$, which we do by the formula $fv = f(A)v$. Here $f(A)$ is an $n \times n$ matrix, so the right hand side is defined by the ordinary multiplication of vectors by matrices.

We need to check the module axioms. Axioms (a) to (e) only involve addition and negation so they are clear. Axiom (f) is also clear because fv is certainly a vector in K^n . If $f(x)$ is constant polynomial 1, then $f(A)$ is the identity matrix, so $fv = Iv = v$ for all v ; this gives axiom (g). For axiom (h) we recall that $(fg)(A) = f(A)g(A)$ so

$$\begin{aligned}
(fg)v &= (fg)(A)v \\
&= f(A)g(A)v \\
&= f(A)(gv) \\
&= f(gv).
\end{aligned}$$

Similarly, axiom (i) follows from the fact that $(f+g)(A) = f(A) + g(A)$. Finally, axiom (j) is clear, because $B(v+w) = Bv + Bw$ for any matrix B and any vectors v and w .

Remark 4.6. [rem-MA-MB]

Let A and B be two different $n \times n$ matrices. Then M_A and M_B have the same elements but the multiplication rules in M_A and M_B are different, so M_A and M_B are different modules.

Example 4.7. [eg-fA-diagonal]

Let A be the matrix $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ over \mathbb{Q} , so that $f(A) = \begin{bmatrix} f(2) & 0 \\ 0 & f(3) \end{bmatrix}$ (by Example 4.2). Then $M_A = \mathbb{Q}^2$, with the multiplication rule $f.(s, t) = (f(2)s, f(3)t)$. For example, if $g(x) = x^2 - 6$ then $g(2) = -2$ and $g(3) = 3$ so we have

$$(x^2 - 6)(10, 11) = (-2 \times 10, 3 \times 11) = (-20, 33).$$

Example 4.8. [eg-fA-jordan]

Let A be the matrix $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ over \mathbb{Q} , so that $f(A) = \begin{bmatrix} f(1) & f'(1) \\ 0 & f(1) \end{bmatrix}$ (by Example 4.3). Then M_A is the set \mathbb{Q}^2 with the group operation $f.(s, t) = (f(1)s + f'(1)t, f(1)t)$. For example, if $g(x) = x^2 - 6$ then $g(1) = -5$ and $g'(1) = 2$ so we have

$$(x^2 - 6)(10, 11) = (-5 \times 10 + 2 \times 11, -5 \times 11) = (-28, -55).$$

Example 4.9. [eg-eigen-module]

The simplest examples are where A is just a 1×1 matrix, or in other words just an element $\lambda \in K$. The module M_λ is just a copy of K , with the multiplication rule $f.a = f(\lambda)a$. For example, the polynomial $f(x) = 1 + x + x^2$ satisfies $f(2) = 7$, so in the module M_2 over $\mathbb{Q}[x]$ we have $f.6 = 7 \times 6 = 42$.

There is a well-known correspondence between matrices and endomorphisms, and for many purposes it is more natural to use the latter. Let V be a vector space over a field K , and let ϕ be an endomorphism of V (in other words, a linear map from V to itself). Then we can define $\phi^2(v) = \phi(\phi(v))$ to get a new endomorphism of V , and similarly we can define ϕ^k for all $k \geq 0$. More generally, for any polynomial $f(x) = a_0 + a_1x + \dots + a_dx^d \in K[x]$ we can define an endomorphism $f(\phi)$ by

$$f(\phi)(v) = a_0v + a_1\phi(v) + \dots + a_d\phi^d(v).$$

We can then make V into a module over $K[x]$ by defining $fv = f(\phi)(v)$.

Example 4.10. [eg-diffop-endo]

We can define a map $\partial: C^\infty(\mathbb{R}, \mathbb{R}) \rightarrow C^\infty(\mathbb{R}, \mathbb{R})$ by $\partial(f) = f'$. As $(f + g)' = f' + g'$ and $(cf)' = cf'$ for constant c , we see that ∂ is an \mathbb{R} -linear endomorphism of $C^\infty(\mathbb{R}, \mathbb{R})$, making $C^\infty(\mathbb{R}, \mathbb{R})$ into a module over $\mathbb{R}[x]$. The multiplication rule is as follows: if $p(x) = \sum_i a_ix^i$ and $f(t) \in C^\infty(\mathbb{R}, \mathbb{R})$ then

$$\begin{aligned} p.f &= a_0\partial^0(f) + a_1\partial^1(f) + a_2\partial^2(f) + \dots \\ &= a_0f + a_1f' + a_2f'' + \dots \end{aligned}$$

Thus, this example is just the same as Example 2.10, with a slightly different viewpoint and less natural notation.

We explained above how a vector space V over K with an endomorphism ϕ can be regarded as a $K[x]$ -module. We conclude this section by showing that *every* $K[x]$ -module arises in this way.

Indeed, let M be a module over $K[x]$. As mentioned previously, axioms (a) to (e) say that M is an Abelian group under addition. Also, if $a \in K$ then we can regard a as a constant polynomial, so am is defined for all $m \in M$. As M is a module over $K[x]$, axioms (f) to (j) are valid for all polynomials a and b , so certainly they are valid for the special case of constant polynomials. Thus, we can regard M as a module over K . A module over a field is the same thing as a vector space, so M is a vector space over K .

Next, if $m \in M$ then xm is another element of M , so we can define a function $\phi: M \rightarrow M$ by $\phi(m) = xm$. I claim that this is a K -linear endomorphism. Indeed, for any $m, n \in M$ we have $x(m+n) = xm + xn$ by the right distributivity law, which means that $\phi(m+n) = \phi(m) + \phi(n)$. Moreover, for $a \in K$ we have $ax = xa$, so

$$a\phi(m) = a(xm) = (ax)m = (xa)m = x(am) = \phi(am)$$

(using axiom (h) twice). This shows that ϕ is linear, as claimed. Now consider a polynomial $f(x) = \sum_i a_ix^i \in K[x]$. I claim that $fm = \sum_i a_i\phi^i(m) = f(\phi)(m)$ for all $m \in M$. Indeed, we have

$$\begin{aligned} (x^2)m &= x(xm) = x\phi(m) = \phi(\phi(m)) = \phi^2(m) \\ (x^3)m &= x(x^2m) = x\phi^2(m) = \phi(\phi^2(m)) = \phi^3(m). \end{aligned}$$

Extending this by induction, we see that $x^k m = \phi^k(m)$ for all k . Thus

$$\begin{aligned} fm &= \left(\sum_i a_ix^i \right) m \\ &= \sum_i a_ix^i m \\ &= \sum_i a_i\phi^i(m) \\ &= f(\phi)(m). \end{aligned}$$

Thus, the module structure is obtained from the endomorphism ϕ in the way considered previously.

Exercises

Exercise 4.1. [ex-jordan-powers]

Let A be the matrix $\begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$. Find A^2 , A^3 and A^4 . Can you give a general rule for A^n ?

Exercise 4.2. [ex-MA-calc]

- (a) Put $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $m = (1, 1, 1) \in M_A$. Calculate $(x^3 - 1)m$.
- (b) Put $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$ and $m = (1, 2, 3) \in M_A$. Calculate $(x^3 - 1)m$.
- (c) Put $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $m = (1, -1) \in M_A$. Calculate $(14x^{12} + 5x^{11} - 36x^7 - 22x^4 + 13x - 5)m$. (You may wish to start by calculating fm for some very simple polynomials f first.)

Exercise 4.3. [ex-cayley-two]

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be a 2×2 matrix, and put $f(x) = x^2 - (a + d)x + (ad - bc)$. Show that $f(A) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. (This is the 2×2 case of the Cayley-Hamilton theorem.)

Exercise 4.4. [ex-fA-block]

Put $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $f(x) = x^4 - 3x$. Calculate $f(A)$.

Exercise 4.5. [ex-fA-swap]

Consider the matrix

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

and a polynomial $f(x) = \sum_i a_i x^i$.

- (a) Calculate A^i for some small numbers i , then give the general rule.
- (b) Write $b = a_0 + a_2 + a_4 + \dots = \sum_j a_{2j}$ and $c = a_1 + a_3 + \dots = \sum_j a_{2j+1}$. Express $f(1)$ and $f(-1)$ in terms of b and c .
- (c) Show that

$$f(A) = \frac{f(1)}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + \frac{f(-1)}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}.$$

5. GENERAL MODULE THEORY

Let R be a ring.

Definition 5.1. [defn-submodule]

Let M be an R -module. A *submodule* of M is a subset $N \subseteq M$ such that

- (a) $0 \in N$
- (b) If $n, m \in N$ then $n + m \in N$ (ie N is closed under addition)
- (c) If $n \in N$ and $a \in R$ then $an \in N$ (ie N is closed under multiplication by elements of R).

Note that if N is a submodule and $n \in N$ then $-n = (-1)n \in N$, so N is closed under negation and thus is a subgroup of M under addition. It is easy to see that N can itself be considered as an R -module.

Example 5.2. [eg-vector-subspace]

If R is a field, then modules are just the same as vector spaces, and submodules are just the same as vector subspaces.

Example 5.3. [eg-subgroup]

If $R = \mathbb{Z}$, then modules are just the same as Abelian groups, and submodules are just the same as subgroups.

Example 5.4. [eg-trivial-submodule]

If M is a module over any ring R , it is clear that $\{0\}$ and M itself are submodules of M .

Example 5.5. [eg-stable-subspace]

Let V be a vector space over a field K , equipped with a K -linear endomorphism $\phi: V \rightarrow V$. We regard V as a $K[x]$ -module in the usual way. We say that a subset $W \subseteq V$ is *stable under ϕ* if $\phi(w) \in W$ for all $w \in W$ (or more briefly, if $\phi(W) \subseteq W$).

I claim that a subset $W \subseteq V$ is a $K[x]$ -submodule if and only if it is a vector subspace and is stable under ϕ . Indeed, suppose that W is a submodule. Then it is certainly closed under addition and under multiplication by constant polynomials (ie elements of K) so it is a vector subspace. Also, it is closed under multiplication by x , so for $w \in W$ we have $\phi(w) = xw \in W$; this shows that W is stable under ϕ , as claimed.

Conversely, suppose that W is a vector subspace and is stable under ϕ . Clearly W is closed under addition. For any $w \in W$ we have $\phi(w) \in W$. Thus $\phi^2(w) = \phi(\phi(w)) = \phi(\text{an element of } W) = \text{another element of } W$, so $\phi^2(W) \subseteq W$. Thus $\phi^3(w) = \phi(\phi^2(w)) = \phi(\text{an element of } W) = \text{another element of } W$, so $\phi^3(W) \subseteq W$, and so on, so $\phi^k(w) \in W$ for all $k \geq 0$. Now consider a polynomial $f(x) = a_0 + \dots + a_d x^d \in K[x]$. We then have $fw = \sum_i a_i \phi^i(w)$. The vectors $\phi^i(w)$ lie in W , the coefficients a_i lie in K , and W is a vector subspace of V , so we see that $\sum_i a_i \phi^i(w) \in W$. Thus $fw \in W$ for all $w \in W$ and $f \in K[x]$, so W is a submodule of V .

Example 5.6. Let A be the matrix $\begin{bmatrix} 0 & -6 \\ 1 & 5 \end{bmatrix}$ over \mathbb{Q} , and use this to make \mathbb{Q}^2 into a module over $\mathbb{Q}[x]$. Put $W_0 = \{(u, v) \in \mathbb{Q}^2 \mid u = -3v\}$ and $W_1 = \{(u, v) \in \mathbb{Q}^2 \mid u = -4v\}$. A typical element of W_0 has the form $(-3v, v)$ and we have

$$\begin{bmatrix} 0 & -6 \\ 1 & 5 \end{bmatrix} \begin{bmatrix} -3v \\ v \end{bmatrix} = \begin{bmatrix} -6v \\ 2v \end{bmatrix},$$

which also lies in W_0 . Thus W_0 is stable under A and thus is a submodule of \mathbb{Q}^2 .

However, W_1 is not a submodule. Indeed, the vector $(-4, 1)$ lies in W_1 but

$$\begin{bmatrix} 0 & -6 \\ 1 & 5 \end{bmatrix} \begin{bmatrix} -4 \\ 1 \end{bmatrix} = \begin{bmatrix} -6 \\ 1 \end{bmatrix},$$

which does not lie in W_1 .

Example 5.7. [eg-eigenspaces]

Suppose that $\lambda, \mu \in K$ and $\lambda \neq \mu$. Define $\phi: K^2 \rightarrow K^2$ by $\phi(u, v) = (\lambda u, \mu v)$, and use this to make K^2 into a module over $K[x]$. Define

$$L = \{(u, 0) \mid u \in K\} \subset K^2$$

$$M = \{(0, v) \mid v \in K\} \subset K^2.$$

I claim that L and M are $K[x]$ -submodules of K^2 , and moreover that the only submodules are $\{0\}$, L , M and K^2 itself.

It is clear that L and M are vector subspaces of K^2 . Moreover we have $\phi(u, 0) = (\lambda u, 0) \in L$, so L is stable under ϕ and thus is a submodule. Similarly $\phi(0, v) = (0, \mu v) \in M$, so M is a submodule. It is trivial to check that $\{0\}$ and K^2 are subspaces of K^2 .

Now let W be any submodule of K^2 . Then W is also a vector subspace, with $0 \leq \dim(W) \leq \dim(K^2) = 2$. If $\dim(W) = 0$ then clearly $W = 0$, and if $\dim(W) = 2$ then clearly $W = K^2$. We can thus assume that $\dim(W) = 1$, so $W = K \cdot (u, v)$ for some vector $(u, v) \neq (0, 0)$. Because W is a $K[x]$ -submodule, we know that $(\lambda u, \mu v) \in W$, but $W = K \cdot (u, v)$ so $(\lambda u, \mu v) = \nu \cdot (u, v)$ for some $\nu \in K$, so $(\lambda - \nu)u = (\mu - \nu)v = 0$. If $u \neq 0$ then we deduce that $\nu = \lambda$ so $(\mu - \lambda)v = 0$, but $\mu \neq \lambda$ so $v = 0$. This means that $W = K \cdot (u, 0) = L$. Similarly, if $v \neq 0$ we deduce that $W = M$.

Example 5.8. [eg-poly-diff]

The set $\mathbb{R}[t]$ of polynomial functions is a vector subspace of the space $C^\infty(\mathbb{R}, \mathbb{R})$ of all smooth functions from \mathbb{R} to \mathbb{R} . Moreover if $f \in \mathbb{R}[t]$ then the derivative of f is again a polynomial, in other words $\partial(f) = f' \in \mathbb{R}[t]$. This means that the subspace $\mathbb{R}[t]$ is stable under the endomorphism ∂ , so it is an $\mathbb{R}[D]$ -submodule of $C^\infty(\mathbb{R}, \mathbb{R})$.

Example 5.9. [eg-sin-cos]

Let W be the space of functions of the form $f(t) = a \cos(t) + b \sin(t)$ (with $a, b \in \mathbb{R}$). Because $\partial(a \cos(t) + b \sin(t)) = b \cos(t) - a \sin(t)$, we see that W is stable under ∂ . It is thus an $\mathbb{R}[D]$ -submodule of $C^\infty(\mathbb{R}, \mathbb{R})$.

Remark 5.10. [rem-intersection]

Suppose that N_0 and N_1 are two submodules of an R -module M . I claim that $N_0 \cap N_1$ is again a submodule. Indeed, as $0 \in N_0$ and $0 \in N_1$ we have $0 \in N_0 \cap N_1$. Suppose that $n, m \in N_0 \cap N_1$. As $n, m \in N_0$ and N_0 is a submodule we have $n + m \in N_0$. As $n, m \in N_1$ and N_1 is a submodule we have $n + m \in N_1$. Thus $n + m \in N_0 \cap N_1$. Now suppose that $a \in R$. As N_0 is a submodule and $n \in N_0$ we have $an \in N_0$. As N_1 is a submodule and $n \in N_1$ we also have $an \in N_1$, so $an \in N_0 \cap N_1$. This shows that $N_0 \cap N_1$ is a submodule, as claimed.

Definition 5.11. [defn-sum]

Suppose that N_0 and N_1 are two submodules of an R -module M . We define $N_0 + N_1$ to be the set of elements $x \in M$ that can be written in the form $x = n_0 + n_1$ for some $n_0 \in N_0$ and $n_1 \in N_1$. I claim that this is a submodule of M . Indeed, suppose that $x, y \in N_0 + N_1$, so we can write $x = n_0 + n_1$ and $y = m_0 + m_1$ with $n_0, m_0 \in N_0$ and $n_1, m_1 \in N_1$. Then $x + y$ can be written as $(n_0 + m_0) + (n_1 + m_1)$, with $n_0 + m_0 \in N_0$ and $n_1 + m_1 \in N_1$, so $x + y \in N_0 + N_1$. Similarly, if $a \in R$ then $an_0 \in N_0$ and $an_1 \in N_1$ so $ax = an_0 + an_1 \in N_0 + N_1$. This shows that $N_0 + N_1$ is closed under addition and under multiplication by R , so it is a submodule as claimed.

Definition 5.12. [defn-direct-ext]

Let N_0 and N_1 be R -modules. We define $N_0 \oplus N_1$ to be the set of pairs (n_0, n_1) with $n_0 \in N_0$ and $n_1 \in N_1$. We make this set into an R -module by defining

$$\begin{aligned}(n_0, n_1) + (m_0, m_1) &= (n_0 + m_0, n_1 + m_1) \\ a(n_0, n_1) &= (an_0, an_1).\end{aligned}$$

(It is a longish but straightforward exercise to check that the axioms are satisfied.) This R -module is called the *external direct sum* of N_0 and N_1 .

Example 5.13. [eg-Z-six]

The group \mathbb{Z}_2 has elements $\bar{0}$ and $\bar{1}$, and the group \mathbb{Z}_3 has elements $\bar{0}$, $\bar{1}$ and $\bar{2}$. Thus, the group $\mathbb{Z}_2 \oplus \mathbb{Z}_3$ has elements $(\bar{0}, \bar{0})$, $(\bar{0}, \bar{1})$, $(\bar{0}, \bar{2})$, $(\bar{1}, \bar{0})$, $(\bar{1}, \bar{1})$ and $(\bar{1}, \bar{2})$. To illustrate the addition law, we have $(\bar{1}, \bar{2}) + (\bar{1}, \bar{2}) = (\bar{2}, \bar{4})$. The first component is to be interpreted as an element of \mathbb{Z}_2 , so $\bar{2} = \bar{0}$. The second component is to be interpreted as an element of \mathbb{Z}_3 , so $\bar{4} = \bar{1}$. Thus $(\bar{1}, \bar{2}) + (\bar{1}, \bar{2}) = (\bar{0}, \bar{1})$.

Example 5.14. [eg-Rn-Rm]

An element of $R^n \oplus R^m$ is a pair (u, v) with $u \in R^n$ and $v \in R^m$, or in other words a list $(u_1, \dots, u_n, v_1, \dots, v_m)$ where each u_i and v_j is an element of R . Thus, $R^n \oplus R^m = R^{n+m}$.

The next example relies on the following definition:

Definition 5.15. [defn-block-sum]

Let A and B be matrices over a field K , of sizes $p \times q$ and $n \times m$. The *block sum* of A and B is the matrix $\left(\begin{array}{c|c} A & 0_{n \times q} \\ \hline 0_{p \times m} & B \end{array} \right)$, of size $(p+n) \times (q+m)$. This is denoted by $A \oplus B$. For example, if $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix}$ then the block sum of A and B is

$$A \oplus B = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \oplus \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 0 & 0 & 5 & 6 \\ 0 & 0 & 7 & 8 \end{bmatrix}.$$

Note that an element $w \in R^{p+n}$ can be written as $w = (u, v)$ with $u \in R^p$ and $v \in R^n$, and we have

$$(A \oplus B)w = \left(\begin{array}{c|c} A & 0 \\ \hline 0 & B \end{array} \right) \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} Au \\ Bv \end{pmatrix}.$$

Example 5.16. [eg-block-sum]

Let A and B be square matrices over a field K , of sizes n and m say. We then have modules M_A and M_B over $K[x]$. The elements of $M_A \oplus M_B$ are pairs $w = (u, v)$ with $u \in K^n$ and $v \in K^m$, or equivalently they are elements of K^{n+m} . The module structure is given by the rule $x(u, v) = (xu, xv) = (Au, Bv)$, or in other words $xw = (A \oplus B)w$. Thus $M_A \oplus M_B = M_{A \oplus B}$.

Definition 5.17. [defn-direct-int]

Let M be an R -module, and let N_0 and N_1 be submodules. We say that M is the *internal direct sum* of N_0 and N_1 if $N_0 + N_1 = M$ and $N_0 \cap N_1 = \{0\}$.

Remark 5.18. [rem-int-ext]

We can define a function $\sigma: N_0 \oplus N_1 \rightarrow M$ by $\sigma(n_0, n_1) = n_0 + n_1$. When we have defined homomorphisms and isomorphisms of modules, we will see that σ is always a homomorphism, and that σ is an isomorphism if and only if M is the internal direct sum of N_0 and N_1 . This is the precise sense in which internal direct sums are “the same” as external ones.

Example 5.19. [eg-eigen-splitting]

In example 5.7 we see that K^2 is the internal direct sum of L and M .

Example 5.20. Consider the Abelian group $M = \mathbb{Z}_{12}$ as a module over \mathbb{Z} . Put $N_0 = \{\bar{0}, \bar{3}, \bar{6}, \bar{9}\}$ and $N_1 = \{\bar{0}, \bar{4}, \bar{8}\}$. It is easy to see that N_0 and N_1 are subgroups, and obviously $N_0 \cap N_1 = \{\bar{0}\}$. I claim that we also have $N_0 + N_1 = M$. Indeed, we have $\bar{1} = \bar{9} + \bar{4} \in N_0 + N_1$ and $N_0 + N_1$ is a submodule so for any $a \in \mathbb{Z}$ we have $\bar{a} = a \cdot \bar{1} \in N_0 + N_1$, as required. Thus M is the internal direct sum of N_0 and N_1 .

Example 5.21. [eg-trig-splitting]

Let V be the space of functions $f \in C^\infty(\mathbb{R}, \mathbb{R})$ that satisfy $f'' = f$. This is a vector space closed under differentiation, so it is an $\mathbb{R}[D]$ -submodule of $C^\infty(\mathbb{R}, \mathbb{R})$. Put $W_0 = \{f \mid f' = f\}$ and $W_1 = \{f \mid f' = -f\}$. These are also vector spaces closed under differentiation, so they are $\mathbb{R}[D]$ -submodules of $C^\infty(\mathbb{R}, \mathbb{R})$. If $f \in W_1$ then $f'' = (-f)' = -(-f) = f$, so $f \in V$. This shows that $W_1 \subseteq V$ and similarly $W_0 \subseteq V$, so W_0 and W_1 are submodules of V .

I claim that V is the direct sum of W_0 and W_1 . One way to see this is just to solve the differential equations. We find that V consists of all functions of the form $ae^t + be^{-t}$, that W_0 consists of all functions of the form ae^t , and that W_1 consists of all functions of the form be^{-t} , and the claim is clear from this.

We can also prove the claim without solving the differential equations explicitly. Indeed, if $f \in W_0 \cap W_1$ then $f = f'$ (because $f \in W_0$) and $f' = -f$ (because $f \in W_1$) so $f = -f$, so $f = 0$. This shows that $W_0 \cap W_1 = \{0\}$. Next, suppose that $g \in V$, so $g'' = g$. Put $g_0 = (g + g')/2$ and $g_1 = (g - g')/2$. We find that $g'_0 = (g' + g'')/2 = (g' + g)/2 = g_0$, so $g_0 \in W_0$. Similarly, we have $g'_1 = (g' - g'')/2 = (g' - g)/2 = -g_1$, so $g_1 \in W_1$. As $g = g_0 + g_1$ it follows that $g \in W_0 + W_1$, and we conclude that $V = W_0 + W_1$ as required.

Definition 5.22. [defn-submod-generated]

Let M be a module over a ring R , and let m_1, \dots, m_r be elements of M . Let N be the set of elements $x \in M$ that can be written in the form $x = a_1 m_1 + \dots + a_r m_r$ for some $a_1, \dots, a_r \in R$. I claim that this is a submodule of M . Indeed, if $x, y \in N$ then we have $x = \sum_i a_i m_i$ and $y = \sum_i b_i m_i$ for some $a_1, \dots, a_r, b_1, \dots, b_r \in R$. We then have $x + y = \sum_i (a_i + b_i) m_i$ so $x + y \in N$; this shows that N is closed under addition. Similarly, if $c \in R$ we have $cx = \sum_i (ca_i) m_i \in N$, so N is closed under multiplication by R , so it is a submodule as claimed.

We call N the *submodule generated by* $\{m_1, \dots, m_r\}$. In particular, we say that M is *generated by* $\{m_1, \dots, m_r\}$ if $N = M$, or equivalently if every element $x \in M$ can be written in the form $a_1 m_1 + \dots + a_r m_r$. We say that M is *finitely generated* if there is some finite list of elements that generates M .

Definition 5.23. [defn-cyclic]

We say that an R -module M is *cyclic* if there is a single element $m \in M$ that generates M , which means that every element $x \in M$ can be written in the form $x = am$ for some $a \in R$.

Example 5.24. [eg-standard-basis]

The module R^d is clearly generated by the standard basis elements $e_1 = (1, 0, \dots, 0)$, $e_2 = (0, 1, 0, \dots, 0)$ and so on. In particular it is finitely generated. It is not cyclic unless $d = 1$.

Example 5.25. [eg-finite-fg]

Let M be a finite Abelian group, considered as a \mathbb{Z} -module. Let the elements of M be m_1, \dots, m_d . Then any element $m \in M$ is equal to m_i for some i , so certainly it can be expressed in the form $\sum_i a_i m_i$ (for example, $m_2 = 0 \cdot m_1 + 1 \cdot m_2 + 0 \cdot m_3 + \dots + 0 \cdot m_d$). Thus, M is finitely generated as a \mathbb{Z} -module.

Example 5.26. [eg-poly-cyclic]

Let W_2 be the space of functions of the form $f(t) = a + bt + ct^2$, considered as a module over $\mathbb{R}[D]$ in the usual way. In particular, the function $g(t) = t^2$ gives an element of W_2 . I claim that W_2 is generated by g , and thus is cyclic. Indeed, we have $g'(t)/2 = t$ and $g''(t)/2 = 1$. It follows that for any function $f(t) = a + bt + ct^2$, we have $(c + (b/2)D + (a/2)D^2)g = cg + (b/2)g' + (a/2)g'' = f$, so $f \in \mathbb{R}[D]g$. This proves that $\mathbb{R}[D]g = W_2$ as required.

It is not hard to extend this method to show that the space W_d of polynomials of degree at most d is also a cyclic module over $\mathbb{R}[D]$ generated by the function $g(t) = t^d$.

Example 5.27. [eg-inf-gen]

Consider $\mathbb{R}[x]$ as a module over \mathbb{R} ; I claim it is not finitely generated. Indeed, suppose we have a finite list f_1, \dots, f_n of elements of $\mathbb{R}[x]$. Let d_i be the degree of the polynomial f_i , and put $d = \max(d_1, \dots, d_n)$. Then each of the polynomials f_i only involves the powers $1, x, x^2, \dots, x^d$, so any polynomial of the form $a_1f_1 + \dots + a_nf_n$ (with $a_1, \dots, a_n \in \mathbb{R}$) also involves only these powers. This means that x^{d+1} cannot be written in the form $a_1f_1 + \dots + a_nf_n$, so the elements f_1, \dots, f_n do not generate $\mathbb{R}[x]$ as a module over \mathbb{R} .

Exercises**Exercise 5.1.** [ex-image-annihilator]

Let $\alpha: M \rightarrow N$ be a homomorphism of modules over $\mathbb{C}[x]$. Suppose that $(x^3 - x)M = \{0\}$ and $x^5N = \{0\}$. Prove that for $n \in \text{image}(\alpha)$ we have $xn = 0$. Can you formulate a general theorem of which this is a special case?

Exercise 5.2. [ex-cyclic-retract]

Let R be a ring, and let M and N be R -modules. Show that if $M \oplus N$ is cyclic, then so are M and N .

Exercise 5.3. [ex-which-submodule]

- (a) Put $N_0 = \{(n, m) \in \mathbb{Z}^2 \mid n - m \text{ is even}\}$ and $N_1 = \{(n, m) \in \mathbb{Z}^2 \mid n - m \text{ is odd}\}$. Are these \mathbb{Z} -submodules of \mathbb{Z}^2 ?
- (b) Put $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $N_0 = \{(u, v) \in \mathbb{R}^2 \mid u - v = 0\}$ and $N_1 = \{(u, v) \mid u + v = 0\}$. Are these $\mathbb{R}[x]$ -submodules of M_A ?
- (c) Put $N_0 = \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid f(1) = 0\}$ and $N_1 = \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid \int_0^2 f = 0\}$. Are these $\mathbb{R}[D]$ -submodules of $C^\infty(\mathbb{R}, \mathbb{R})$?

Exercise 5.4. [ex-ten-elements]

Let R be a ring with exactly 10 elements, and let M be an R -module with exactly 20 elements. Prove that M is not a free module.

Exercise 5.5. [ex-twenty-four]

For any integer d , let N_d be the submodule of \mathbb{Z}_{24} generated by \bar{d} . The group N_6 has precisely 4 elements; list them. Find integers d and e such that $N_6 \cap N_d = N_d$ and $N_6 + N_d = N_e$.

Exercise 5.6. [ex-nine-hundred]

For any natural number d dividing 900, let N_d be the submodule of \mathbb{Z}_{900} generated by \bar{d} .

- (a) What is the order of N_{10} ?
- (b) Which standard group is isomorphic to \mathbb{Z}_{900}/N_{10} ?
- (c) Find d such that the submodule generated by $\bar{70}$ is N_d .
- (d) Find d such that $N_{12} + N_{30} + N_{100} = N_d$.
- (e) Find d such that $N_{30} \cap N_{50} = N_d$.

Exercise 5.7. [ex-MA-cyclic]

Consider the following matrix over \mathbb{Q} .

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

Show that the module M_A over $\mathbb{Q}[x]$ is cyclic, and give a polynomial $f(x)$ such that $M_A \simeq \mathbb{Q}[x]/f(x)$.

Exercise 5.8. [ex-Wd-cyclic]

Let W_d be the set of polynomials $f(t)$ of degree at most d . Prove that W_d is a cyclic module over $\mathbb{R}[D]$. What is the ideal $I \subseteq \mathbb{R}[D]$ such that $W_d \simeq \mathbb{R}[D]/I$?

Exercise 5.9. [ex-crossprod]

Fix a nonzero vector $u \in \mathbb{R}^3$, and write $r = \|u\|$. Define an endomorphism $\phi: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by $\phi(v) = u \times v$. Write M for \mathbb{R}^3 , considered as a module over $\mathbb{R}[x]$ using ϕ . Let L be the line through u and 0, and let K be the plane perpendicular to L .

- (a) Show that L is a submodule of M , and that $xL = 0$.
- (b) Show that K is a submodule of M , and that $(x^2 + r^2)K = 0$ and $xM \leq K$.
- (c) Show that $(x^3 + r^2x)M = 0$.

(You will need a number of standard facts about dot and cross products of vectors.)

6. HOMOMORPHISMS

Definition 6.1. [defn-homomorphism]

Let M and N be modules over a ring R . An R -module homomorphism (or just homomorphism) from M to N is a function $\alpha: M \rightarrow N$ such that

- (a) $\alpha(m_0 + m_1) = \alpha(m_0) + \alpha(m_1)$ for all $m_0, m_1 \in M$.
- (b) $\alpha(am) = a\alpha(m)$ for all $a \in R$ and $m \in M$.

Note that this implies that $\alpha(0) = \alpha(0 \cdot 0) = 0\alpha(0) = 0$ and $\alpha(-m) = \alpha((-1) \cdot m) = (-1) \cdot \alpha(m) = -\alpha(m)$.

An *isomorphism* is a homomorphism which is also a bijection.

Remark 6.2. [rem-inv-homo]

Let $\alpha: M \rightarrow N$ be an isomorphism. As α is a bijection, there is an inverse function $\alpha^{-1}: N \rightarrow M$ such that $\alpha(\alpha^{-1}(n)) = n$ for all $n \in N$ and $\alpha^{-1}(\alpha(m)) = m$ for all $m \in M$. I claim that α^{-1} is also a homomorphism. To see this, suppose that $n_0, n_1 \in N$. We then have elements $\alpha^{-1}(n_0)$ and $\alpha^{-1}(n_1)$ in M . As α is a homomorphism, we have $\alpha(\alpha^{-1}(n_0) + \alpha^{-1}(n_1)) = \alpha(\alpha^{-1}(n_0)) + \alpha(\alpha^{-1}(n_1)) = n_0 + n_1$. We can apply α^{-1} to this equation to get $\alpha^{-1}(\alpha(\alpha^{-1}(n_0) + \alpha^{-1}(n_1))) = \alpha^{-1}(n_0 + n_1)$. Because $\alpha^{-1}(\alpha(m)) = m$ for all m , the left hand side is just $\alpha^{-1}(n_0) + \alpha^{-1}(n_1)$. We thus have $\alpha^{-1}(n_0) + \alpha^{-1}(n_1) = \alpha^{-1}(n_0 + n_1)$, showing that α^{-1} respects addition.

Similarly, suppose that $n \in N$ and $a \in R$. As α respects multiplication by R , we have $\alpha(a\alpha^{-1}(n)) = a\alpha(\alpha^{-1}(n)) = an$. By applying α^{-1} to this equation we get $\alpha^{-1}(\alpha(a\alpha^{-1}(n))) = \alpha^{-1}(an)$. The left hand side is just $a\alpha^{-1}(n)$, so we have $a\alpha^{-1}(n) = \alpha^{-1}(an)$, completing the proof that α^{-1} is a homomorphism.

Example 6.3. [eg-tau-sg-d1]

Let R be any ring. Define $\tau: R^2 \rightarrow R^2$, $\sigma: R^3 \rightarrow R$ and $\delta: R^2 \rightarrow R^3$ by

$$\begin{aligned} \tau(u, v) &= (v, u) \\ \sigma(x, y, z) &= x + y + z \\ \delta(u, v) &= (u, v - u, -v). \end{aligned}$$

It is easy to check that these are all homomorphisms. For example, we have

$$\begin{aligned}\delta(u_0, v_0) + \delta(u_1, v_1) &= (u_0, v_0 - u_0, -v_0) + (u_1, v_1 - u_1, -v_1) \\ &= (u_0 + u_1, v_0 + v_1 - u_0 - u_1, -v_0 - v_1) \\ &= \delta(u_0 + u_1, v_0 + v_1) \\ &= \delta((u_0, v_0) + (u_1, v_1))\end{aligned}$$

and

$$\begin{aligned}a\delta(u, v) &= a.(u, v - u, -v) \\ &= (au, av - au, -av) \\ &= \delta(au, av) \\ &= \delta(a.(u, v)),\end{aligned}$$

so δ is a homomorphism.

Example 6.4. [eg-bad-defn]

I would like to define two homomorphisms $\alpha, \beta: \mathbb{Z}_3 \rightarrow \mathbb{Z}_{12}$ by $\alpha(\overline{m}) = \overline{4m}$ and $\beta(\overline{m}) = \overline{5m}$. There is a potential problem with this kind of definition, which means that the definition of β is actually invalid, although it turns out that α is OK. Consider the element $x = \overline{1} \in \mathbb{Z}_3$, which can also be described as $x = \overline{4}$. Using the description $x = \overline{1}$ we get $\beta(x) = \overline{5} \in \mathbb{Z}_{12}$. Using the description $x = \overline{4}$ we get $\beta(x) = \overline{20} \in \mathbb{Z}_{12}$. As $20 \not\equiv 5 \pmod{12}$, the elements $\overline{5}$ and $\overline{20}$ in \mathbb{Z}_{12} are not the same, so our definition of β is not self-consistent.

However, this problem does not occur with α . To see why, suppose we describe an element $y \in \mathbb{Z}_3$ in two different ways, say $y = \overline{n} = \overline{m}$. As $\overline{n} = \overline{m}$ in \mathbb{Z}_3 , we have $n = m \pmod{3}$, so $n = m + 3k$ for some integer k . This means that $4n = 4m + 12k$, so $\overline{4n} = \overline{4m}$ in \mathbb{Z}_{12} . This means that we get the same answer for $\alpha(y)$ no matter which description we use, so α is a well-defined function from \mathbb{Z}_3 to \mathbb{Z}_{12} .

We also have $\alpha(\overline{n}) + \alpha(\overline{m}) = \overline{4n} + \overline{4m} = \overline{4(n+m)} = \alpha(\overline{n+m})$, so α is a homomorphism of groups. It follows easily that it is also a homomorphism of \mathbb{Z} -modules.

The above example generalizes as follows:

Proposition 6.5. [prop-cyclic-hom]

Let p, q and r be integers such that $p, q > 0$ and pr is divisible by q . Then there is a unique homomorphism $\alpha: \mathbb{Z}_p \rightarrow \mathbb{Z}_q$ such that $\alpha(\overline{m}) = \overline{rm}$ for all $m \in \mathbb{Z}$.

Proof. By assumption we have $pr = qs$ for some integer s . If $\overline{n} = \overline{m}$ in \mathbb{Z}_p then $m = n + pk$ for some k , so $rm = rn + rpk = rn + qsk$, so $rm = rn \pmod{q}$, so $\overline{rm} = \overline{rn}$ in \mathbb{Z}_q . This shows that α is well-defined. We also have $\alpha(\overline{n}) + \alpha(\overline{m}) = \overline{rn} + \overline{rm} = \overline{r(n+m)} = \alpha(\overline{n+m})$, so α is a homomorphism. \square

Example 6.6. [eg-cyclic-hom]

Because 6×5 is divisible by 15, there is a unique homomorphism $\alpha: \mathbb{Z}_6 \rightarrow \mathbb{Z}_{15}$ such that $\alpha(\overline{m}) = \overline{5m}$ for all m .

Example 6.7. [eg-free-hom]

Let N be a module over a ring R , and let n_1, \dots, n_d be a list of elements of N . We define a function $\alpha: R^d \rightarrow N$ by

$$\alpha(u_1, \dots, u_d) = u_1 n_1 + \dots + u_d n_d.$$

I claim that this is a homomorphism. Indeed, we have

$$\begin{aligned}\alpha((u_1, \dots, u_d) + (v_1, \dots, v_d)) &= \alpha(u_1 + v_1, \dots, u_d + v_d) \\ &= (u_1 + v_1)n_1 + \dots + (u_d + v_d)n_d \\ &= (u_1 n_1 + \dots + u_d n_d) + (v_1 n_1 + \dots + v_d n_d) \\ &= \alpha(u_1, \dots, u_d) + \alpha(v_1, \dots, v_d),\end{aligned}$$

and

$$\begin{aligned}
\alpha(a(u_1, \dots, u_d)) &= \alpha(au_1, \dots, au_d) \\
&= au_1n_1 + \dots + au_dn_d \\
&= a(u_1n_1 + \dots + u_dn_d) \\
&= a\alpha(u_1, \dots, u_d)
\end{aligned}$$

as required.

Next, I claim that every homomorphism $\beta: R^d \rightarrow N$ is of the form just described. To see this, let e_k be the element of R^d given by $e_k = (0, \dots, 0, 1, 0, \dots, 0)$, with the 1 in the k 'th place. Put $n_k = \beta(e_k) \in N$. Any element $u = (u_1, \dots, u_d)$ in R^d can be written as $u = u_1e_1 + \dots + u_de_d$. For example, in the case $d = 3$ we have

$$(u_1, u_2, u_3) = u_1(1, 0, 0) + u_2(0, 1, 0) + u_3(0, 0, 1) = u_1e_1 + u_2e_2 + u_3e_3.$$

As β is a homomorphism, we have

$$\begin{aligned}
\beta(u_1, \dots, u_d) &= \beta(u_1e_1 + \dots + u_de_d) \\
&= u_1\beta(e_1) + \dots + u_d\beta(e_d) \\
&= u_1n_1 + \dots + u_dn_d,
\end{aligned}$$

so β has the form described previously.

We summarize the above discussion as follows:

Proposition 6.8. [prop-free-hom]

For any list (n_1, \dots, n_d) of elements of N we have a homomorphism $\alpha: R^d \rightarrow N$ given by $\alpha(u) = \sum_i u_i n_i$. Conversely, every homomorphism $\alpha: R^d \rightarrow N$ arises in this way from a unique list (n_1, \dots, n_d) . \square

Example 6.9. [eg-matrix]

We now consider homomorphisms $\alpha: R^d \rightarrow R^e$. By the previous example, α corresponds to a list u_1, \dots, u_d , where each u_i is an element of R^e , or in other words a vector of length e . We can construct a matrix A (with d columns and e rows) whose columns are the vectors u_1, \dots, u_d , and we find that $\alpha(x) = Ax$ for all $x \in R^d$.

Consider for example the homomorphism $\delta: R^2 \rightarrow R^3$ given by $\delta(s, t) = (s, t - s, -t)$. We have

$$\begin{aligned}
u_1 &= \delta(e_1) = \delta(1, 0) = (1, -1, 0) \\
u_2 &= \delta(e_2) = \delta(0, 1) = (0, 1, -1) \\
A &= \begin{pmatrix} 1 & 0 \\ -1 & 1 \\ 0 & -1 \end{pmatrix}
\end{aligned}$$

so

$$A \begin{bmatrix} s \\ t \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} s \\ t \end{bmatrix} = \begin{bmatrix} s \\ t - s \\ -t \end{bmatrix} = \alpha(s, t)$$

as claimed.

We summarize the above discussion as follows:

Proposition 6.10. [prop-matrix]

Homomorphisms from R^d to R^e are essentially the same as $d \times e$ matrices over R . \square

We next consider homomorphisms of modules over polynomial rings.

Proposition 6.11. [prop-Kx-hom]

Let V and W be vector spaces over a field K , and let $\phi: V \rightarrow V$ and $\psi: W \rightarrow W$ be K -linear maps. We use these to make V and W into modules over $K[x]$ in the usual way, so that $xv = \phi(v)$ for $v \in V$ and $xw = \psi(w)$ for $w \in W$. Then the $K[x]$ -module homomorphisms from V to W are precisely the K -linear maps $\gamma: V \rightarrow W$ such that $\psi\gamma = \gamma\phi$, or in other words $\psi(\gamma(v)) = \gamma(\phi(v))$ for all $v \in V$.

Proof. Let $\gamma: V \rightarrow W$ be a $K[x]$ -module homomorphism, so $\gamma(v_0 + v_1) = \gamma(v_0) + \gamma(v_1)$ for all $v_0, v_1 \in V$, and $\gamma(av) = a\gamma(v)$ for all $a \in K[x]$ and $v \in V$. By taking a to be a constant polynomial, we see that $\gamma(av) = a\gamma(v)$ for all $a \in K$, so γ is a K -linear map. Now take $a = x$ instead. As $v \in V$ we have $xv = \phi(v)$, and as $\gamma(v) \in W$ we have $x\gamma(v) = \psi(\gamma(v))$. Thus, the equation $\gamma(xv) = x\gamma(v)$ becomes $\gamma(\phi(v)) = \psi(\gamma(v))$, as required.

Conversely, suppose we have a K -linear map $\gamma: V \rightarrow W$ satisfying $\gamma\phi = \psi\gamma$. It follows that

$$\gamma\phi^2 = (\gamma\phi)\phi = (\psi\gamma)\phi = \psi(\gamma\phi) = \psi^2\gamma.$$

This can be extended by induction to show that $\gamma\phi^k = \psi^k\gamma$ for all $k \geq 0$. Thus, for any polynomial $p(x) = \sum_i a_i x^i$ we have

$$\gamma p(\phi) = \sum_i a_i \gamma\phi^i = \sum_i a_i \psi^i \gamma = p(\psi)\gamma,$$

so $\gamma(p(x)v) = p(x)\gamma(v)$, so γ is a $K[x]$ -module homomorphism. □

Remark 6.12. [rem-diagram]

We can indicate where the maps γ , ϕ and ψ go by a diagram as follows:

$$\begin{array}{ccc} V \ni t & \xrightarrow{\gamma} & W \\ \phi \downarrow & & \downarrow \psi \\ V & \xrightarrow{\gamma} & W. \end{array}$$

The condition $\psi\gamma = \gamma\phi$ means that the two way round the square are the same. This is usually expressed by saying that the diagram *commutes*.

Remark 6.13. [rem-Hom-MA-MB]

The proposition can be restated as follows in terms of matrices: If A and B are square matrices of size n and m over a field K , then the $K[x]$ -module homomorphisms from M_A to M_B are the $n \times m$ matrices C over K such that $CA = BC$. Here the matrices A , B and C correspond to the linear maps ϕ , ψ and γ respectively.

Remark 6.14. [rem-usually-zero]

If matrices A and B are chosen randomly, then usually the only matrix C satisfying $CA = BC$ will be the zero matrix. Of course, most of the examples in these notes are chosen specially so that there are some nonzero solutions.

Example 6.15. [eg-Hom-MA-MB]

Put $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}$. The homomorphisms from M_A to M_B are then the matrices $C = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for which $CA = BC$, or in other words

$$\begin{bmatrix} a+b & a+b \\ c+d & c+d \end{bmatrix} = \begin{bmatrix} 2c & 2d \\ 2a & 2b \end{bmatrix},$$

or equivalently

$$\begin{aligned} a+b &= 2c \\ a+b &= 2d \\ c+d &= 2a \\ c+d &= 2b. \end{aligned}$$

Solving these equations gives $a = b = c = d$. Thus, the homomorphisms from M_A to M_B are precisely the matrices of the form $\begin{bmatrix} a & a \\ a & a \end{bmatrix}$ for some $a \in K$.

Example 6.16. Define $\phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $\phi(u, v) = (v, u)$, and define $\psi: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by $\psi(u, v, w) = (v, w, u)$. We use these to make \mathbb{R}^2 and \mathbb{R}^3 into modules over $\mathbb{R}[x]$ as usual. Define $\gamma: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by $\gamma(u, v) = (u + v, u + v, u + v)/2$. I claim that this is an $\mathbb{R}[x]$ -module homomorphism. It is clearly \mathbb{R} -linear, so it is enough to check that $\gamma\phi = \psi\gamma$. We have $\gamma\phi(u, v) = \gamma(v, u) = (v + u, v + u, v + u)/2$. We also have $\gamma(u, v) = (u + v, u + v, u + v)/2$ and $\psi(w, w, w) = (w, w, w)$ for any w , so $\psi\gamma(u, v) = (u + v, u + v, u + v)/2$, which is the same as $\gamma\phi(u, v)$, as claimed.

Next, I claim that any other $\mathbb{R}[x]$ -module homomorphism $\beta: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ is actually a multiple of γ . To see this, note that $\beta(1, 1) \in \mathbb{R}^3$, so $\beta(1, 1) = (\lambda, \mu, \nu)$ for some $\lambda, \mu, \nu \in \mathbb{R}$. As β is a homomorphism we must have $\beta\phi(1, 1) = \psi\beta(1, 1) = \psi(\lambda, \mu, \nu) = (\mu, \nu, \lambda)$. On the other hand, we have $\phi(1, 1) = (1, 1)$ so $\beta\phi(1, 1) = \beta(1, 1) = (\lambda, \mu, \nu)$. Thus $(\lambda, \mu, \nu) = (\mu, \nu, \lambda)$, so $\nu = \mu = \lambda$. This means that $\beta(1, 1) = (\lambda, \lambda, \lambda) = \lambda \cdot (1, 1, 1)$.

We next claim that $\beta(1, -1) = (0, 0, 0)$. To see this, note that $\beta\phi^3 = \psi^3\beta$. Moreover $\psi^3(u, v, w) = (u, v, w)$ for all $(u, v, w) \in \mathbb{R}^3$, and $\phi^3(1, -1) = (-1, 1) = -(1, -1)$, so the equation $\psi^3\beta(1, -1) = \beta\phi^3(1, -1)$ becomes $\beta(1, -1) = \beta(-1, 1) = -\beta(1, -1)$. By rearranging we see that $\beta(1, -1) = 0$ as claimed.

Now consider an arbitrary element $(u, v) \in \mathbb{R}^2$. We can write this as

$$(u, v) = \frac{u+v}{2}(1, 1) + \frac{u-v}{2}(1, -1),$$

so

$$\begin{aligned} \beta(u, v) &= \frac{u+v}{2}\beta(1, 1) + \frac{u-v}{2}\beta(1, -1) \\ &= \frac{u+v}{2}(\lambda, \lambda, \lambda) + \frac{u-v}{2}(0, 0, 0) \\ &= \lambda\gamma(u, v). \end{aligned}$$

This proves that $\beta = \lambda\gamma$, as claimed.

Example 6.17. [eg-shift]

Define $\tau: C^\infty(\mathbb{R}, \mathbb{R}) \rightarrow C^\infty(\mathbb{R}, \mathbb{R})$ by $\tau(f)(t) = f(t+1)$. Thus if $g(t) = 3t$ and $h(t) = \sin(2\pi t)$ and $k(t) = 2^t$ then $\tau(g) = g+3$ and $\tau(h) = h$ and $\tau(k) = 2k$. It is clear that τ is an \mathbb{R} -linear map. By the chain rule we have

$$\frac{d}{dt}f(t+1) = f'(t+1)\frac{d}{dt}(t+1) = f'(t+1),$$

so $\partial(\tau(f)) = \tau(\partial(f))$, so τ is a homomorphism of $\mathbb{R}[D]$ -modules.

Proposition 6.18. [prop-conj-iso]

Let A and B be $n \times n$ matrices over a field K . Then M_A is isomorphic to M_B if and only if A is conjugate to B , in other words there is an invertible $n \times n$ matrix P such that $PAP^{-1} = B$.

Proof. A homomorphism from M_A to M_B is an $n \times n$ matrix P such that $PA = BP$. Such a homomorphism is an isomorphism if and only if P is invertible, and if so, the condition $PA = BP$ is equivalent to the condition $PAP^{-1} = B$. \square

Corollary 6.19. [cor-semisimple]

Suppose that A is a diagonalizable $n \times n$ matrix over K . Then there exists an invertible matrix P such that $D := PAP^{-1}$ is a diagonal matrix, with diagonal entries $\lambda_1, \dots, \lambda_n$ say. Then M_A is isomorphic to M_D and thus to $M_{\lambda_1} \oplus \dots \oplus M_{\lambda_n}$. \square

Definition 6.20. [defn-ker-img]

Let $\alpha: M \rightarrow N$ be a homomorphism of modules over a ring R . We define the kernel and image of α by

$$\begin{aligned} \ker(\alpha) &= \{m \in M \mid \alpha(m) = 0\} \\ \text{image}(\alpha) &= \{n \in N \mid n = \alpha(m) \text{ for some } m \in M\}. \end{aligned}$$

Proposition 6.21. [prop-img-submodule]

$\ker(\alpha)$ is a submodule of M and $\text{image}(\alpha)$ is a submodule of N .

Proof. Suppose that $m_0, m_1 \in \ker(\alpha)$ and that $a \in R$. We then have $\alpha(m_0) = \alpha(m_1) = 0$ and so

$$\begin{aligned} \alpha(m_0 + m_1) &= \alpha(m_0) + \alpha(m_1) = 0 + 0 = 0 \\ \alpha(am_0) &= a\alpha(m_0) = a \cdot 0 = 0, \end{aligned}$$

so $m_0 + m_1 \in \ker(\alpha)$ and $am_0 \in \ker(\alpha)$. This shows that $\ker(\alpha)$ is a submodule of M .

Now suppose that $n_0, n_1 \in \text{image}(\alpha)$ and $a \in R$. We then have $n_0 = \alpha(m_0)$ for some $m_0 \in M$ and $n_1 = \alpha(m_1)$ for some $m_1 \in M$. It follows that $\alpha(m_0 + m_1) = n_0 + n_1$, so $n_0 + n_1$ is $\alpha(\text{something})$, so $n_0 + n_1 \in \text{image}(\alpha)$. Similarly, we have $an_0 = \alpha(am_0)$, so $an_0 \in \text{image}(\alpha)$. This shows that $\text{image}(\alpha)$ is a submodule of N . \square

Example 6.22. [eg-times-four]

Define $\alpha: \mathbb{Z} \rightarrow \mathbb{Z}_{12}$ by $\alpha(n) = \overline{4n}$. We then have

$$\begin{array}{lll} \alpha(0) & = \overline{0} & \alpha(1) & = \overline{4} & \alpha(2) & = \overline{8} \\ \alpha(3) & = \overline{12} = \overline{0} & \alpha(4) & = \overline{16} = \overline{4} & \alpha(5) & = \overline{20} = \overline{8} \\ \alpha(6) & = \overline{24} = \overline{0} & \alpha(7) & = \overline{28} = \overline{4} & \alpha(8) & = \overline{32} = \overline{8} \end{array}$$

and everything repeats with period three. It follows that the only elements of \mathbb{Z}_{12} that can be written in the form $\alpha(n)$ are $\overline{0}$, $\overline{4}$ and $\overline{8}$, so $\text{image}(\alpha) = \{\overline{0}, \overline{4}, \overline{8}\}$. It also follows that $\alpha(n) = \overline{0}$ if and only if n is divisible by 3, so $\ker(\alpha) = \{n \in \mathbb{Z} \mid n = 0 \pmod{3}\} = 3\mathbb{Z}$.

Example 6.23. [eg-sum-hom]

Define $\sigma: R^3 \rightarrow R$ by $\sigma(x, y, z) = x + y + z$ and $\delta: R^2 \rightarrow R^3$ by $\delta(u, v) = (u, v - u, -v)$. I claim that $\text{image}(\delta) = \ker(\sigma)$.

To see this, first suppose that $(x, y, z) \in \text{image}(\delta)$. This means that $(x, y, z) = \delta(u, v) = (u, v - u, -v)$ for some u, v . It follows that $\sigma(x, y, z) = x + y + z = u + (v - u) + (-v) = 0$, so $(x, y, z) \in \ker(\sigma)$. This proves that $\text{image}(\delta) \subseteq \ker(\sigma)$.

Conversely, suppose that $(x, y, z) \in \ker(\sigma)$. This means that $x + y + z = 0$, so $-(x + y) = z$. From this it follows that $(x, y, z) = (x, (x + y) - x, -(x + y)) = \delta(x, x + y)$, so (x, y, z) is $\delta(\text{something})$, so $(x, y, z) \in \text{image}(\delta)$. This proves that $\ker(\sigma) \subseteq \text{image}(\delta)$ and thus that $\ker(\sigma) = \text{image}(\delta)$, as claimed.

Remark 6.24. [rem-exact-seq]

Suppose we have modules L, M, N and homomorphisms $\alpha: L \rightarrow M$ and $\beta: M \rightarrow N$ such that $\text{image}(\alpha) = \ker(\beta)$. We then say that the sequence $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is *exact*; for example, the sequence $R^2 \xrightarrow{\delta} R^3 \xrightarrow{\sigma} R$ in the above example is exact. This is a very important concept elsewhere in the theory of modules, although we will make little use of it in this course.

Proposition 6.25. [prop-jective]

Let $\alpha: M \rightarrow N$ be a homomorphism of R -modules. Then

- α is injective if and only if $\ker(\alpha) = \{0\}$.
- α is surjective if and only if $\text{image}(\alpha) = N$.
- α is an isomorphism if and only if $\ker(\alpha) = \{0\}$ and $\text{image}(\alpha) = N$.

Proof. First, suppose that $\ker(\alpha) = \{0\}$. If $\alpha(m_0) = \alpha(m_1)$ then $\alpha(m_0 - m_1) = \alpha(m_0) - \alpha(m_1) = 0$, so $m_0 - m_1 \in \ker(\alpha) = \{0\}$, so $m_0 - m_1 = 0$, so $m_0 = m_1$. This proves that α is injective.

Conversely, suppose that α is injective. If $m \in \ker(\alpha)$ then $\alpha(m) = 0$, so $\alpha(m) = \alpha(0)$, and as α is injective this means that $m = 0$. Thus $\ker(\alpha) = \{0\}$. This completes the proof of (a).

For (b), note that $\text{image}(\alpha)$ is the set of things in N that can be written in the form $\alpha(m)$ for some m . Thus $\text{image}(\alpha) = N$ if and only if every element in N can be written as $\alpha(m)$ for some m , and this is precisely the definition of surjectivity.

Finally, an isomorphism is just a bijective homomorphism. It is standard that a function is a bijection if and only if it is both injective and surjective, so (c) follows immediately from (a) and (b). \square

Exercises

Exercise 6.1. [ex-Hom-MA-MB-i]

In each of the following situations, find all the $\mathbb{Q}[x]$ -module homomorphisms from M_A to M_B .

- $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.
- $A = \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}$ and $B = \begin{bmatrix} \mu & 0 \\ 0 & \lambda \end{bmatrix}$, where $\lambda \neq \mu$.
- $A = I_2$ (the 2×2 identity matrix) and $B = I_3$.

Exercise 6.2. [ex-Hom-MA-MB-ii]

- Put $A = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$. Find all the $\mathbb{Q}[x]$ -module homomorphisms from M_A to M_B .
- Put $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Find all the $\mathbb{Q}[x]$ -module homomorphisms from M_A to M_B .

- (c) Suppose that $\lambda_1, \lambda_2, \lambda_3, \mu_1, \mu_2 \in \mathbb{C}$, and put $A = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}$ and $B = \begin{bmatrix} \mu_1 & 0 \\ 0 & \mu_2 \end{bmatrix}$. Show that for most values of the λ 's and μ 's, the only $\mathbb{C}[x]$ -module homomorphism from M_A to M_B is zero. What can you say about the exceptional cases?

Exercise 6.3. [ex-annihilator]

Let a be an element of a ring R . For any R -module M , put

$$\text{ann}(a, M) = \{m \in M \mid am = 0\}.$$

We will also write R/a for the factor module R/Ra .

- Find $\text{ann}(4, \mathbb{Z}_{12})$.
- Find $\text{ann}(x - 1, M_A)$, where A is the matrix $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$.
- Let $\alpha: R/a \rightarrow M$ be a homomorphism. Show that $\alpha(\bar{1}) \in \text{ann}(a, M)$.
- Conversely, given $m \in \text{ann}(a, M)$, show that there is a unique homomorphism $\alpha: R/a \rightarrow M$ such that $\alpha(\bar{1}) = m$.
- Describe all the $\mathbb{R}[D]$ -module homomorphisms from $\mathbb{R}[D]/(D^2 - 1)$ to $C^\infty(\mathbb{R}, \mathbb{R})$.

Exercise 6.4. [ex-Z-nine-SES]

Show that there are homomorphisms $\alpha: \mathbb{Z}_3 \rightarrow \mathbb{Z}_9$ and $\beta: \mathbb{Z}_9 \rightarrow \mathbb{Z}_3$ given by $\alpha(\bar{n}) = \overline{3n}$ and $\beta(\bar{m}) = \bar{m}$. Show that the sequence $\mathbb{Z}_3 \xrightarrow{\alpha} \mathbb{Z}_9 \xrightarrow{\beta} \mathbb{Z}_3$ is exact.

Exercise 6.5. [ex-no-homs]

- Let α be a $\mathbb{C}[x]$ -module homomorphism from $\mathbb{C}[x]/(x^2 - 1)$ to M_A , where $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$. Prove that $\alpha = 0$.
- Let V be the space of functions of the form $a \sin(t) + b \cos(t)$, and let W be the space of functions of the form $a \sinh(t) + b \cosh(t)$. Let $\beta: V \rightarrow W$ be an $\mathbb{R}[D]$ -module homomorphism. Show that $\beta = 0$.
- Let $\gamma: \mathbb{Z}_4 \rightarrow \mathbb{Z}^4$ be a homomorphism of \mathbb{Z} -modules. Prove that $\gamma = 0$.

Exercise 6.6. [ex-count-Hom]

Let R be a ring, and let M be an R -module with only finitely many elements, say $|M| = m$. How many homomorphisms are there from R^d to M ?

7. FACTOR MODULES

Let M be an R -module for some ring R , and let N be a submodule. We next define the factor module M/N .

For any element $m \in M$ we define the set $m + N = \{m + n \mid n \in N\}$, which is a subset of M . A *coset* of N in M is a subset $C \subseteq M$ that can be written in the form $C = m + N$ for some m . We write M/N for the set of all such cosets.

Example 7.1. [eg-three-cosets]

Take $R = \mathbb{Z}$ and $M = \mathbb{Z}$ and

$$\begin{aligned} N &= 3\mathbb{Z} = \{n \in \mathbb{Z} \mid n = 0 \pmod{3}\} \\ &= \{\dots, -9, -6, -3, 0, 3, 6, 9, \dots\}. \end{aligned}$$

Consider the following three sets:

$$\begin{aligned} A &= \{\dots, -10, -7, -4, -1, 2, 5, 8, \dots\} \\ B &= \{\dots, -9, -6, -3, 0, 3, 6, 9, \dots\} \\ C &= \{\dots, -8, -5, -2, 1, 4, 7, 10, \dots\}. \end{aligned}$$

The set A can be described as $-7 + N$ or as $-1 + N$ or as $26 + N$, so it is a coset. Similarly, B can be described as $0 + N$ or $999 + N$ and C can be described as $-8 + N$ or $1 + N$, so A , B and C are all cosets. In fact, they are the only cosets, so $\mathbb{Z}/N = \{A, B, C\}$.

Proposition 7.2. [prop-cosets]

Let R , M and N be as above, and suppose that $m_0, m_1 \in M$. Then the following are equivalent

- (a) $m_0 + N = m_1 + N$
- (b) $m_0 - m_1 \in N$
- (c) $m_0 \in m_1 + N$
- (d) $m_1 \in m_0 + N$
- (e) $(m_0 + N) \cap (m_1 + N)$ is nonempty.

Proof. If $m_0 \in m_1 + N$ then $m_0 = m_1 + n$ for some $n \in N$ so $m_0 - m_1 = n \in N$. Conversely, if $m_0 - m_1 \in N$ then the equation $m_0 = m_1 + (m_0 - m_1)$ shows that $m_0 \in m_1 + N$, so statements (b) and (c) are equivalent. Similarly, (b) and (d) are equivalent.

If (c) holds then m_0 lies in both $m_0 + N$ and $m_1 + N$, so $(m_0 + N) \cap (m_1 + N) \neq \emptyset$, so (e) holds.

Conversely suppose that (e) holds, so there is an element x lying in both $m_0 + N$ and $m_1 + N$. This means that $x = m_0 + n_0$ for some $n_0 \in N$ and $x = m_1 + n_1$ for some $n_1 \in N$, so $m_0 = x - n_0 = m_1 + (n_1 - n_0)$. This shows that $m_0 \in m_1 + N$, so (c) holds. We now see that (b), (c), (d) and (e) are all equivalent to each other.

If (a) holds then it is clear that (c) holds and thus that (b),(d) and (e) also hold. Conversely, suppose that (b) holds. If $x \in m_0 + N$ then $x = m_0 + n$ for some $n \in N$, so $x = m_1 + ((m_0 - m_1) + n)$ and $(m_0 - m_1) + n \in N$ so $x \in m_1 + N$. This shows that $m_0 + N \subseteq m_1 + N$, and a similar argument shows that $m_1 + N \subseteq m_0 + N$, so (a) holds. This now shows that all five statements are equivalent to each other. \square

We next want to define addition of cosets. Given two cosets C_0 and C_1 we choose $m_0, m_1 \in M$ such that $C_0 = m_0 + N$ and $C_1 = m_1 + N$, and then we would like to define $C_0 + C_1 = (m_0 + m_1) + N$. Similarly, if $a \in R$ we would like to define $aC_0 = (am_0) + N$. There is a potential problem here: suppose we chose a different description of the same coset C_0 (say as $m'_0 + N$) and a different description of C_1 (say as $m'_1 + N$). This gives an apparently different answer for $C_0 + C_1$: before we had $(m_0 + m_1) + N$, now we have $(m'_0 + m'_1) + N$. If these were genuinely different cosets then our definition of addition would be ambiguous and thus invalid. However, we will show that these are simply different descriptions of the same coset, so our definition is unambiguous after all.

Lemma 7.3. [lem-cosets]

If $m_0 + N = m'_0 + N$ and $m_1 + N = m'_1 + N$ then $(m_0 + m_1) + N = (m'_0 + m'_1) + N$ and $(am_0) + N = (am'_0) + N$.

Proof. By Proposition 7.2 we have $m_0 - m'_0 \in N$ and $m_1 - m'_1 \in N$. As N is closed under addition this means that $m_0 - m'_0 + m_1 - m'_1 \in N$, or in other words $(m_0 + m_1) - (m'_0 + m'_1) \in N$, so $(m_0 + m_1) + N = (m'_0 + m'_1) + N$ as claimed. Similarly, as $m_0 - m'_0 \in N$ and N is a submodule, we have $am_0 - am'_0 = a(m_0 - m'_0) \in N$, so $(am_0) + N = (am'_0) + N$ as claimed. \square

Corollary 7.4. [cor-cosets]

We can unambiguously define addition of cosets and multiplication of cosets by elements of R , using the formulae

$$(m_0 + N) + (m_1 + N) = (m_0 + m_1) + N$$

$$a(m + N) = (am) + N.$$

Notation 7.5. [ntn-cosets]

If there is no ambiguity about which submodule N is intended, we will write \bar{m} for $m + N$.

Proposition 7.6. [prop-factor-module]

The definitions in Corollary 7.4 make the set M/N into an R -module. Moreover, the function $\pi: M \rightarrow M/N$ defined by $\pi(m) = \bar{m} = m + N$ is an R -module homomorphism.

Proof. We need to check all the axioms in Definition 3.1. All the proofs follow the same pattern, so we will do only two of them. We first consider axiom (d), which says that addition is associative. Let A, B, C be any

three cosets; we must show that $A + (B + C) = (A + B) + C$. We can choose $a, b, c \in M$ such that $A = a + N$, $B = b + N$ and $C = c + N$. From the definition of addition in M/N we have $B + C = (b + c) + N$ and thus $A + (B + C) = (a + (b + c)) + N$. Similarly, we have $(A + B) + C = ((a + b) + c) + N$. As addition is associative in the module M that we started with, we have $(a + b) + c = a + (b + c)$ so $((a + b) + c) + N = (a + (b + c)) + N$, so $(A + B) + C = A + (B + C)$ as required.

We next check axiom (j) (which says that multiplication is right-distributive). Let A and B be elements of M/N , and let r be an element of R ; we must show that $r(A + B) = rA + rB$. Choose a and b such that $A = a + N$ and $B = b + N$. We then have $A + B = (a + b) + N$ so $r(A + B) = r((a + b) + N) = (r(a + b)) + N = (ra + rb) + N = (ra + N) + (rb + N) = r(a + N) + r(b + N) = rA + rB$, as required.

The very definition of addition and multiplication in M/N says that $\pi(m_0) + \pi(m_1) = \pi(m_0 + m_1)$ and $a\pi(m) = \pi(am)$, so π is an R -module homomorphism. \square

Theorem 7.7 (The first isomorphism theorem). *Let $\alpha: M \rightarrow N$ be a homomorphism of R -modules. Put $K = \ker(\alpha) \subseteq M$ and $L = \text{image}(\alpha) \subseteq N$. Then there is an isomorphism $\bar{\alpha}: M/K \rightarrow L$ such that $\bar{\alpha}(m + K) = \alpha(m)$ for all $m \in M$.*

Proof. Let C be a coset in M/K . We can choose an element $m \in M$ such that $C = m + K$, and clearly $\alpha(m) \in \text{image}(\alpha) = L$. We would like to define $\bar{\alpha}(C) = \alpha(m)$, but we need to check that this is well-defined. Suppose we describe the same coset C in a different way, say as $C = m' + K$. This gives an apparently different answer for $\bar{\alpha}(C)$: before we had $\alpha(m)$, now we have $\alpha(m')$. As $m' + K = m + K$ we have $m' - m \in K = \ker(\alpha)$, which means that $\alpha(m') - \alpha(m) = \alpha(m' - m) = 0$, so $\alpha(m') = \alpha(m)$, so our two answers are actually the same. Thus, we have a well-defined function $\bar{\alpha}: M/K \rightarrow L$ satisfying $\bar{\alpha}(m + K) = \alpha(m)$ for all m .

Next, we have

$$\begin{aligned} \bar{\alpha}((m_0 + K) + (m_1 + K)) &= \bar{\alpha}((m_0 + m_1) + K) \\ &= \alpha(m_0 + m_1) \\ &= \alpha(m_0) + \alpha(m_1) \\ &= \bar{\alpha}(m_0 + K) + \bar{\alpha}(m_1 + K) \end{aligned}$$

and

$$\begin{aligned} \bar{\alpha}(a(m + K)) &= \bar{\alpha}((am) + K) \\ &= \alpha(am) \\ &= a\alpha(m) \\ &= a\bar{\alpha}(m + K), \end{aligned}$$

so $\bar{\alpha}$ is a homomorphism.

We next show that $\bar{\alpha}: M/K \rightarrow L$ is surjective. As L was defined as the image of α , any element $n \in L$ has the form $n = \alpha(m)$ for some $m \in M$. This means that $n = \bar{\alpha}(m + K)$, so n is in the image of $\bar{\alpha}$. As this is true for every element of L , the homomorphism $\bar{\alpha}$ is surjective.

Finally, we show that $\bar{\alpha}$ is injective. Suppose we have cosets C_0, C_1 with $\bar{\alpha}(C_0) = \bar{\alpha}(C_1)$. Choose $m_0, m_1 \in M$ such that $C_0 = m_0 + K$ and $C_1 = m_1 + K$. Then the equation $\bar{\alpha}(C_0) = \bar{\alpha}(C_1)$ means that $\alpha(m_0) = \alpha(m_1)$, so $\alpha(m_0 - m_1) = 0$, so $m_0 - m_1 \in \ker(\alpha) = K$. As $m_0 - m_1 \in K$ we have $m_0 + K = m_1 + K$ or in other words $C_0 = C_1$. This proves that $\bar{\alpha}$ is injective as well as surjective, so it is an isomorphism. \square

Example 7.8. [eg-FIT]

Define $\alpha: \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ by $\alpha(u, v) = (u + v, u + v)$, and put $K = \ker(\alpha)$ and $L = \text{image}(\alpha)$. Clearly $\alpha(u, v) = (0, 0)$ if and only if $v = -u$, so

$$K = \{(u, v) \in \mathbb{Z}^2 \mid v = -u\} = \{(t, -t) \mid t \in \mathbb{Z}\}.$$

Next, note that that $\alpha(u, v)$ is always of the form (r, r) for some r . Conversely, any vector of the form (r, r) can be written as $\alpha(r, 0)$, so it lies in the image of α . Thus

$$L = \{(x, y) \in \mathbb{Z}^2 \mid x = y\} = \{(r, r) \mid r \in \mathbb{Z}\}.$$

Consider the set

$$C = \{\dots, (-3, 5), (-2, 4), (-1, 3), (0, 2), (1, 1), (2, 0), (3, -1), \dots\}.$$

This can be described as $C = (1, 1) + K$ or $C = (-3, 5) + K$ or $C = (-999, 1001) + K$, so it is a coset of K , or in other words an element of the group \mathbb{Z}^2/K . We have

$$\bar{\alpha}(C) = \alpha(1, 1) = \alpha(-3, 5) = \alpha(-999, 1001) = (2, 2).$$

Proposition 7.9. [prop-cyclic]

Let M be a cyclic R -module. Then $M \simeq R/I$ for some submodule I of R .

Proof. Choose an element m that generates M . Define a homomorphism $\alpha: R \rightarrow M$ by $\alpha(a) = am$. As m generates M , this homomorphism is surjective. Put $I = \ker(\alpha)$, which is a submodule of R . The First Isomorphism Theorem now tells us that $R/I \simeq M$. \square

Exercises

Exercise 7.1. [ex-SIT]

Let M be a module over a ring R , and let L and N be submodules of M . Prove that $L/(L \cap N)$ is isomorphic to $(L + N)/N$. [You may wish to consider the homomorphism $\pi: L \rightarrow (L + N)/N$ given by $\pi(x) = x + N$.]

Exercise 7.2. [ex-int-ext]

Let M be a module over a ring R , and let N_0 and N_1 be submodules of M . Define a homomorphism $\sigma: N_0 \oplus N_1 \rightarrow M$ by $\sigma(n_0, n_1) = n_0 + n_1$. Prove that σ is an isomorphism if and only if M is the internal direct sum of N_0 and N_1 .

8. IDEALS AND FACTOR RINGS

Definition 8.1. [defn-ideal]

An *ideal* in a ring R is a subset $I \subseteq R$ such that

- (a) $0 \in I$
- (b) If $b, c \in I$ then $b + c \in I$
- (c) If $a \in R$ and $b \in I$ then $ab \in I$.

Remark 8.2. [rem-ideal-submod]

We remarked earlier that R can be regarded as a module over itself. By comparing the above definition with Definition 5.1 we see that ideals are just the same as submodules of R .

Example 8.3. [eg-even]

Let I be the set of even integers; then I is an ideal in \mathbb{Z} . Indeed, 0 is even so (a) holds; the sum of two even integers is even so (b) holds; and the product of an even integer with any other integer is still even so (c) holds.

Example 8.4. [eg-ker-eval]

Put $R = \mathbb{Z}[x]$ and $I = \{f \in \mathbb{Z}[x] \mid f(1) = 0\}$. For example $x^{10} - x \in I$ and $(x - 3)(x - 2)(x - 1) \in I$ but $x + 7 \notin I$ (because $1 + 7 \neq 0$) and $\frac{1}{2}x^2 - x + \frac{1}{2} \notin I$ (because the coefficients are not integers, so $\frac{1}{2}x^2 - x + \frac{1}{2} \notin \mathbb{Z}[x]$). Clearly the zero polynomial is an element of I , so (a) holds. If $f, g \in I$ then $f(1) = g(1) = 0$ so $(f + g)(1) = f(1) + g(1) = 0 + 0 = 0$, so $f + g \in I$; thus (b) holds. If $f \in I$ and g is any polynomial then $(gf)(1) = g(1)f(1) = g(1) \cdot 0 = 0$ so $gf \in I$; thus (c) holds. This shows that I is an ideal in R .

Example 8.5. [eg-const]

Put $R = \mathbb{Z}[x]$ and $I = \{\text{constant polynomials}\} \subseteq R$. Then I is *not* an ideal. Axioms (a) and (b) certainly hold. Moreover, if a and b are elements of I then $ab \in I$ also. However, axiom (c) says more than this: it says that if $b \in I$ and a is *any* element of R , not necessarily in I , then ab must be in I . However, $x \in R$ and $1 \in I$ but $x \cdot 1 \notin I$ so axiom (c) is violated.

Example 8.6. [eg-field-ideal]

Let K be a field. It is easy to see that $\{0\}$ and K itself are ideals in K ; I claim that these are the only ideals. Indeed, let I be an ideal in K . If $I \neq \{0\}$ then we have some nonzero element $b \in I$. For any element $c \in K$ we have $cb^{-1} \in K$ and $b \in I$ so axiom (c) tells us that $(cb^{-1})b \in I$, or in other words $c \in I$. This means that $I = K$, as required.

Example 8.7. [eg-principal]

Let R be any ring, and let x be any element of R . Define $Rx = \{ux \mid u \in R\}$. I claim that this is an ideal (called the *principal ideal* generated by x). First, we have $0 = 0.x \in Rx$, so axiom (a) holds. Second, if $a, b \in Rx$ then there exist u, v such that $a = ux$ and $b = vx$ so $a + b = (u + v)x$, so $a + b \in Rx$. This shows that (b) holds. Finally, if $a \in R$ and $b \in Rx$ then $b = vx$ for some $v \in R$ so $ab = (av)x \in Rx$, so (c) holds.

In example 8.3, the ideal I is just $\mathbb{Z}.2$. In example 8.4, the ideal I is just $\mathbb{Z}[x].(x - 1)$.

Now let I be an ideal in a ring R . Recall that I is an R -submodule of R , so we can define the R -module $R/I = \{x + I \mid x \in R\}$ as before. In the case where I is the principal ideal Ra for some $a \in R$, we will generally write R/a rather than R/Ra .

We next show that R/I can itself be regarded as a ring.

Lemma 8.8. [lem-coset-product]

If $a + I = a' + I$ and $b + I = b' + I$ then $ab + I = a'b' + I$.

Proof. As $a + I = a' + I$, the element $u := a - a'$ lies in I . Similarly, the element $v := b - b'$ lies in I . We have $a = a' + u$ and $b = b' + v$ so

$$ab - a'b' = (a' + u)(b' + v) - a'b' = a'b' + b'u + uv = (a' + u)v + b'u.$$

As $a' + u \in R$ and $v \in I$, axiom (c) says that $(a' + u)v \in I$. As $b' \in R$ and $u \in I$, axiom (c) also says that $b'u \in I$. As I is closed under addition, this means that $(a' + u)v + b'u \in I$, so $ab - a'b' \in I$, so $ab + I = a'b' + I$ as required. \square

It follows that we can define multiplication of cosets unambiguously by $(a + I)(b + I) = ab + I$. By the method of Proposition 7.6 we see that this makes R/I into a ring.

Definition 8.9. [defn-ring-homomorphism]

Let R_0 and R_1 be rings. A *ring homomorphism* from R_0 to R_1 is a function $\alpha: R_0 \rightarrow R_1$ such that

- (a) $\alpha(a + b) = \alpha(a) + \alpha(b)$ for all $a, b \in R_0$
- (b) $\alpha(1) = 1$
- (c) $\alpha(ab) = \alpha(a)\alpha(b)$ for all $a, b \in R_0$.

One can check that a ring homomorphism automatically satisfies $\alpha(0) = 0$ and $\alpha(-a) = -\alpha(a)$. Moreover, if a is invertible in R_0 then $\alpha(a)$ is invertible in R_1 with $\alpha(a)^{-1} = \alpha(a^{-1})$. We say that α is an *isomorphism* if it is a bijection as well as a homomorphism. If so, one can check that the inverse function $\alpha^{-1}: R_1 \rightarrow R_0$ is also a ring homomorphism.

If we define $\pi: R \rightarrow R/I$ by $\pi(a) = a + I$, we find that $\pi(a + b) = \pi(a) + \pi(b)$ and also $\pi(1) = 1$ and $\pi(ab) = \pi(a)\pi(b)$, in other words π is a homomorphism of rings.

The following result is the *First Isomorphism Theorem for Rings*.

Theorem 8.10. [thm-FIT-rings]

Let $\alpha: R_0 \rightarrow R_1$ be a homomorphism of rings. Then $\ker(\alpha)$ is an ideal in R_0 and $\text{image}(\alpha)$ is a subring of R_1 . Moreover, there is a ring isomorphism $\bar{\alpha}: R_0/\ker(\alpha) \simeq \text{image}(\alpha)$ given by $\bar{\alpha}(a + \ker(\alpha)) = \alpha(a)$. In particular, if α is surjective then $R_0/\ker(\alpha) \simeq R_1$.

Proof. Put $K = \ker(\alpha)$ and $R_2 = \text{image}(\alpha)$. If $a, b \in K$ then $\alpha(a) = \alpha(b) = 0$ so $\alpha(a + b) = \alpha(a) + \alpha(b) = 0$ so $a + b \in K$. Also, if c is any element of R_0 then $\alpha(ca) = \alpha(c)\alpha(a) = \alpha(c).0 = 0$, so $ca \in K$. This shows that K is an ideal in R_0 .

As $\alpha(0) = 0$ and $\alpha(1) = 1$ we see that $0, 1 \in \text{image}(\alpha) = R_2$. If $u, v \in R_2$ then we have $u = \alpha(a)$ and $v = \alpha(b)$ for some $a, b \in R_0$. Thus $u + v = \alpha(a + b)$ and $-u = \alpha(-a)$ and $uv = \alpha(ab)$, so $u + v, -u, uv \in R_2$. This shows that R_2 is a subring of R_1 .

Just as in the proof of the first isomorphism theorem for modules, we have a well-defined bijection $\bar{\alpha}: R_0/K \rightarrow R_2$ given by $\bar{\alpha}(a+K) = \alpha(a)$. We then have

$$\bar{\alpha}((a+K)(b+K)) = \bar{\alpha}(ab+K) = \alpha(ab) = \alpha(a)\alpha(b) = \bar{\alpha}(a+K)\bar{\alpha}(b+K),$$

and similarly $\bar{\alpha}(1+K) = 1$ and $\bar{\alpha}((a+K)+(b+K)) = \bar{\alpha}(a+K) + \bar{\alpha}(b+K)$, so $\bar{\alpha}$ is a ring homomorphism. \square

Example 8.11. [eg-FIT-rings]

For any $n > 0$ we put $n\mathbb{Z} = \{nk \mid k \in \mathbb{Z}\} = \{m \in \mathbb{Z} \mid m = 0 \pmod{n}\}$. This is an ideal in \mathbb{Z} , so we have a factor ring $\mathbb{Z}/n\mathbb{Z}$. Note that $a+n\mathbb{Z} = b+n\mathbb{Z}$ iff $a-b \in n\mathbb{Z}$ iff $a = b \pmod{n}$. Using this, we see that $\mathbb{Z}/n\mathbb{Z}$ is just the usual ring \mathbb{Z}_n of residue classes modulo n .

Example 8.12. Let K be a field, and λ an element of K . Put $I = \{f \in K[x] \mid f(\lambda) = 0\}$, which is an ideal in the ring $R := K[x]$. To see this, define a function $\alpha: K[x] \rightarrow K$ by $\alpha(f) = f(\lambda)$. Clearly

$$\begin{aligned} \alpha(f+g) &= (f+g)(\lambda) = f(\lambda) + g(\lambda) = \alpha(f) + \alpha(g) \\ \alpha(fg) &= (fg)(\lambda) = f(\lambda)g(\lambda) = \alpha(f)\alpha(g) \\ \alpha(1) &= 1 \end{aligned}$$

so α is a ring homomorphism. If $c \in K$ then we can regard c as a constant polynomial and we find that $\alpha(c) = c$; this shows that α is surjective. It is clear that $\ker(\alpha) = I$, so $K[x]/I \simeq K$ by the First Isomorphism Theorem.

It is also a standard fact that $f(\lambda) = 0$ iff f is divisible by $x-\lambda$, so I is just the principal ideal $K[x] \cdot (x-\lambda)$, so we have shown that $K[x]/(x-\lambda) \simeq K$.

Example 8.13. [eg-C-as-quotient]

Let I be the principal ideal $\mathbb{R}[x](x^2+1)$ in $\mathbb{R}[x]$. I claim that $\mathbb{R}[x]/(x^2+1) = \mathbb{R}[x]/I$ is isomorphic to \mathbb{C} . The basic point is just that $\bar{x}^2 + 1 = \overline{x^2 + 1} = \bar{0}$, so $\bar{x}^2 = -\bar{1}$, so \bar{x} is a square root of -1 .

To give a formal proof, we consider the function $\alpha: \mathbb{R}[x] \rightarrow \mathbb{C}$ defined by $\alpha(f) = f(i)$. Just as in the last example we find that α is a ring homomorphism. Any complex number z can be written in the form $a+bi$ for some $a, b \in \mathbb{R}$ and we find that $\alpha(a+bx) = a+bi = z$, so α is surjective. Thus, if we put $I = \ker(\alpha) = \{f \in \mathbb{R}[x] \mid f(i) = 0\}$ we have $\mathbb{R}[x]/I \simeq \mathbb{C}$.

As $i^2 + 1 = 0$ we have $x^2 + 1 \in I$ and so $\mathbb{R}[x](x^2 + 1) \subseteq I$. Conversely, suppose that $f \in I$, so $f(i) = 0$. We can divide $f(x)$ by $x^2 + 1$ to get $f(x) = (x^2 + 1)q(x) + a + bx$ for some polynomial $q(x) \in \mathbb{R}[x]$ and $a, b \in \mathbb{R}$. We then have $0 = f(i) = q(i)(i^2 + 1) + a + bi = a + bi$, and by comparing real and imaginary parts we see that $a = b = 0$. This means that $f(x) = q(x)(x^2 + 1)$, so $f(x) \in \mathbb{R}[x] \cdot (x^2 + 1)$. This shows that $I = \mathbb{R}[x](x^2 + 1)$, and so $\mathbb{R}[x]/(x^2 + 1) = \mathbb{R}[x]/I \simeq \mathbb{C}$.

The next example relies on the following result.

Lemma 8.14. [lem-Zpk-units]

Let p be a prime number and $k \geq 0$. If a is not divisible by p then \bar{a} is invertible in \mathbb{Z}_{p^k} .

Proof. Let d be the greatest common divisor of a and p^k . This is in particular a divisor of p^k , so it must be of the form p^j for some $j \leq k$. It must also be a divisor of a , which is impossible if $j > 0$, because a is not divisible by p . We must therefore have $j = 0$ or in other words $d = 1$. As $(a, p^k) = 1$ we have $ab + p^k c = 1$ for some integers b, c . This means that $ab = 1 \pmod{p^k}$ or in other words $\bar{a}\bar{b} = \bar{1}$. This shows that \bar{a} is invertible (with inverse \bar{b}) as required. \square

Example 8.15. [eg-p-local-units]

Let p be a prime, and consider the ring $\mathbb{Z}_{(p)}$ as in example 2.12. Let I be the principal ideal $\mathbb{Z}_{(p)} \cdot p^k$ for some $k \geq 0$. I claim that $\mathbb{Z}_{(p)}/p^k$ is isomorphic to $\mathbb{Z}_{p^k} = \mathbb{Z}/p^k$. To see this, we must define a homomorphism $\rho: \mathbb{Z}_{(p)} \rightarrow \mathbb{Z}/p^k$. Any element of $x \in \mathbb{Z}_{(p)}$ can be written as $x = a/b$ where $a, b \in \mathbb{Z}$ and $b \neq 0 \pmod{p}$. This means that \bar{b} is invertible in \mathbb{Z}/p^k , so we have an element $\bar{a}\bar{b}^{-1} \in \mathbb{Z}/p^k$. Now suppose we describe x in a different way, say as $x = c/d$ with $c, d \in \mathbb{Z}$ and $d \neq 0 \pmod{p}$. Then $a/b = c/d$ so $ad = bc$ so $\bar{a}\bar{d} = \bar{b}\bar{c}$. As \bar{b} and \bar{d} are invertible we can divide through by them to get $\bar{a}\bar{b}^{-1} = \bar{c}\bar{d}^{-1}$. Thus, we can unambiguously define a function $\rho: \mathbb{Z}_{(p)} \rightarrow \mathbb{Z}/p^k$ by $\rho(a/b) = \bar{a}\bar{b}^{-1}$. It is easy to check that this is a ring homomorphism.

For any element $y \in \mathbb{Z}/p^k$ we can write $y = \bar{a}$ for some $a \in \{0, 1, \dots, p^k - 1\}$. Any of these numbers a can be regarded as an element of $\mathbb{Z}_{(p)}$ and then we have $\rho(a) = \bar{a} = y$. This shows that $\text{image}(\rho) = \mathbb{Z}/p^k$.

Next, suppose that $x \in \ker(\rho)$. Then $x = a/b$ for some $a, b \in \mathbb{Z}$ with $b \not\equiv 0 \pmod{p}$ and $\bar{a}\bar{b}^{-1} = \rho(a/b) = 0$ in \mathbb{Z}/p^k . We can multiply this equation by \bar{b} to see that $\bar{a} = 0$ in \mathbb{Z}/p^k , so $a = 0 \pmod{p^k}$, so $a = p^k c$ for some integer c . If we define $y = c/b$ we find that $y \in \mathbb{Z}_{(p)}$ and $x = p^k y$ so $x \in p^k \mathbb{Z}_{(p)}$. Conversely, if $x \in p^k \mathbb{Z}_{(p)}$ then $x = p^k y$ for some $y \in \mathbb{Z}_{(p)}$ so $\rho(y) \in \mathbb{Z}/p^k$ and $\rho(x) = p^k \rho(y)$. However, it is easy to see that $p^k z = 0$ for all $z \in \mathbb{Z}/p^k$, so $\rho(x) = 0$ so $x \in \ker(\rho)$.

The first isomorphism theorem now gives us an isomorphism $\bar{\rho}: \mathbb{Z}_{(p)}/\ker(\rho) \rightarrow \text{image}(\rho)$, or equivalently $\bar{\rho}: \mathbb{Z}_{(p)}/p^k \rightarrow \mathbb{Z}/p^k$.

Proposition 8.16. [prop-RI-mod]

Modules over R/I are the same thing as modules over R with the property that $am = 0$ for all $a \in I$ and $m \in M$.

Proof. Let M be a module over R/I . To make M into an R -module, we need to define am for each $a \in R$ and $m \in M$. Note that $\pi(a) = (a + I) \in R/I$, and M is a module over R/I , so $\pi(a)m$ is already defined. We can thus define am to be $\pi(a)m$. It is easy to check that the axioms are satisfied; for example, we have

$$(a + b)m = \pi(a + b)m = (\pi(a) + \pi(b))m = \pi(a)m + \pi(b)m = am + bm,$$

so multiplication is left distributive. Thus M is a module over R/I . If $a \in I$ then $\pi(a) = 0$ so $am = \pi(a)m = 0$ as required.

Conversely, suppose that M is a module over R with the property that $am = 0$ for all $a \in I$ and $m \in M$. To make M into an R/I -module, we must define Am for each coset $A \in R/I$ and each $m \in M$. We would like to do this by writing A in the form $a + I$ for some $a \in R$ and defining Am to be the same as am . This raises the usual problem of ambiguity, but if $a + I = a' + I$ then $a - a' \in I$, so $(a - a')m = 0$ (by our assumption on M) so $am = a'm$. Thus, we have an unambiguous definition of Am for $A \in R/I$ and $m \in M$. It is again straightforward to check that the module axioms are satisfied. For example, if $A, B \in R/I$ we can choose $a, b \in R$ such that $A = a + I$ and $B = b + I$. We then have $A + B = (a + b) + I$ so

$$(A + B)m = ((a + b) + I)m = (a + b)m = am + bm = (a + I)m + (b + I)m = Am + Bm,$$

so multiplication is left distributive. Thus M is a module over R/I , as required. \square

Example 8.17. [eg-vier]

Consider the Abelian group $V = \{0, a, b, c\}$ with addition table as follows:

$$\begin{aligned} a + a &= b + b = c + c = 0 \\ a + b &= c \\ b + c &= a \\ c + a &= b. \end{aligned}$$

Like any Abelian group, this can be regarded as a \mathbb{Z} -module. As $a + a = b + b = c + c = 0$, we see that $2v = 0$ for all $v \in V$, and thus that $nv = 0$ for all $n \in 2\mathbb{Z}$. It follows that V can be regarded as a module over the ring \mathbb{Z}_2 .

Example 8.18. [eg-lagrange]

Let M be an Abelian group of order d , considered as a module over \mathbb{Z} as usual. By Lagrange's theorem, if $m \in M$ then the order of m divides d . As we are using additive notation, this just means that $dm = 0$. It follows that $am = 0$ for all $a \in d\mathbb{Z}$, so M can be regarded as a module over \mathbb{Z}_d .

Example 8.19. [eg-trig-module]

Let W be the space of functions of the form $f(t) = a \cos(t) + b \sin(t)$ (with $a, b \in \mathbb{R}$). As in Example 5.9, we can regard this as a module over $\mathbb{R}[D]$. If f is as above then $f'(t) = -a \sin(t) + b \cos(t)$ and so $f''(t) = -a \cos(t) - b \sin(t) = -f(t)$, so $(D^2 + 1)f = f'' + f = 0$. If we let I be the principal ideal $(D^2 + 1)\mathbb{R}[D]$ we find that $p(D)f = 0$ for all $p(D) \in I$ and $f \in W$, so W can be regarded as a module over the ring $\mathbb{R}[D]/I = \mathbb{R}[D]/(D^2 + 1)$.

Exercises

Exercise 8.1. [ex-no-ring-homs]

- (a) Show that there are no ring homomorphisms from \mathbb{Z}_3 to \mathbb{Z} . [Consider the equation $\bar{1} + \bar{1} + \bar{1} = \bar{0}$.]
- (b) Show that there are no ring homomorphisms from \mathbb{Q} to \mathbb{Z} . [Consider the equation $\frac{1}{2} \cdot (1 + 1) = 1$.]
- (c) Show that there are no ring homomorphisms from \mathbb{C} to \mathbb{R} .
- (d) Find a ring homomorphism from \mathbb{C} to \mathbb{C} that is not the identity (there is only one reasonable example).

Exercise 8.2. [ex-FIT-rings]

- (a) Prove that $\mathbb{Q}[x]/(x^2 - 2)$ is isomorphic to a subring of \mathbb{R} .
- (b) Let I be the ideal $\mathbb{Z}[i] \cdot (2 + 3i)$ in $\mathbb{Z}[i]$, and define a homomorphism $\alpha: \mathbb{Z} \rightarrow \mathbb{Z}[i]/I$ by $\alpha(n) = n + I$.
 - (i) Show that $\alpha(-5) = i + I$, and deduce that α is surjective.
 - (ii) Suppose that $n \in \mathbb{Z}$ and that n is divisible by $2 + 3i$ in $\mathbb{Z}[i]$. Show that n^2 is divisible by 13 in \mathbb{Z} .
 - (iii) Show that $\mathbb{Z}[i]/I \simeq \mathbb{Z}_{13}$.

Exercise 8.3. [ex-C-as-quotient]

- (a) Prove that the ring $\mathbb{R}[x]/(x^2 + 4)$ is isomorphic to \mathbb{C} .
- (b) Prove that $\mathbb{R}[x]/(x^2 - 4)$ is not a field (and thus cannot be isomorphic to \mathbb{C}).

Exercise 8.4. [ex-F-nine]

Let R be the ring $\mathbb{Z}[i]/3$, and put $u = 1 + i + 3\mathbb{Z}[i] \in R$.

- (a) List the elements of R .
- (b) Calculate u^k for $0 \leq k \leq 8$.
- (c) Compare your list in (a) with your list in (b), and show that R is a field.
- (d) Do you know another proof that R is a field?

9. EUCLIDEAN DOMAINS

We next consider Euclidean domains, which are a particular kind of commutative ring. The idea is to generalize the following two facts:

- (1) If n and m are integers with $m \neq 0$ then we can divide n by m to get a quotient q with remainder r . We then have $n = mq + r$ and $|r| < |m|$.
- (2) If f and g are polynomials over \mathbb{C} with $g \neq 0$ then we can divide f by g to get a quotient q with remainder r . We then have $f = gq + r$ and the degree of r is less than the degree of g .

Given a ring R and elements $a, b \in R$ with $b \neq 0$, we would like to do a similar kind of division to get an equation $a = bq + r$ where the “size” of r is less than the “size” of b . If $R = \mathbb{Z}$ then “size” means absolute value, and if $R = \mathbb{C}[x]$ then “size” means degree. For a general ring there may not be a suitable notion of size, so there may not be a useful division algorithm. A suitable notion of size is called a *Euclidean valuation*; the formal definition is as follows.

Definition 9.1. [defn-valn]

A *Euclidean valuation* on a ring R is a function $\nu(a)$ defined for all nonzero elements a of R such that

- (a) $\nu(a)$ is a nonnegative integer whenever $a \neq 0$.
- (b) If $a, b \in R$ and $b \neq 0$ then there are elements $q, r \in R$ such that $a = bq + r$ and either $r = 0$ or $\nu(r) < \nu(b)$.
- (c) If $a, b \in R$ and $a \neq 0$ and $b \neq 0$ then $ab \neq 0$ and $\nu(a) \leq \nu(ab)$.

A *Euclidean domain* is a ring R for which there exists a Euclidean valuation.

Remark 9.2. The first part of condition (c) says that the product of two nonzero elements is nonzero, or in other words that R is an integral domain.

Example 9.3. [eg-Z-euclidean]

The function $\nu(a) = |a|$ is a Euclidean valuation on \mathbb{Z} . Indeed, conditions (a) and (c) are clear, and (b) is just the ordinary division algorithm for integers.

Example 9.4. [eg-poly-euclidean]

If K is a field then we can define a Euclidean valuation ν on $K[x]$ by $\nu(f) = \deg(f) =$ the degree of f . Indeed, conditions (a) and (c) are clear, and (b) is just the ordinary division algorithm for polynomials.

Example 9.5. [eg-Zi-euclidean]

The function $\nu(x + iy) = |x + iy|^2 = x^2 + y^2$ defines a Euclidean valuation on $\mathbb{Z}[i]$. Indeed, condition (a) is clear. For condition (b), suppose that $a = x + iy$ and $b = u + iv$ for some integers u, v, x, y . As $b \neq 0$ we can consider the complex number $a/b = s + it$ say, where $s, t \in \mathbb{R}$. Let s_0 be the closest integer to s , so $|s - s_0| \leq 1/2$. (If s has the form $m + 1/2$ for some integer m then we could take $s_0 = m$ or $s_0 = m + 1$; it doesn't matter which.) Similarly, we let t_0 be the closest integer to t , so $|t - t_0| \leq 1/2$. We put $q = s_0 + it_0$ and $r = a - qb$ so that $a = qb + r$. We find that

$$|a/b - q|^2 = |(s - s_0) + (t - t_0)i|^2 = (s - s_0)^2 + (t - t_0)^2 \leq 1/4 + 1/4 < 1,$$

so $|r|^2 = |a - qb|^2 = |b|^2|a/b - q|^2 < |b|^2$, or in other words $\nu(r) < \nu(b)$ as required.

Finally, for condition (c), we have $\nu(ab) = |ab|^2 = |a|^2|b|^2$. It is clear that $|b|^2$ is a nonnegative integer, and as $b \neq 0$ we have $|b|^2 \neq 0$ so $|b|^2 \geq 1$ so $\nu(ab) \geq \nu(a)$.

Example 9.6. [eg-p-local-euclidean]

We next define a Euclidean valuation on $\mathbb{Z}_{(p)}$. Any nonzero element $a \in \mathbb{Z}_{(p)}$ has the form u/w , where u and w are integers and w is not divisible by p . If we divide u by p as many times as possible we end up with an equation $u = p^t v$ where v is not divisible by p . We then define $\nu(a) = \nu(p^t v/w) = t$. For example, if $p = 3$ we have $\nu(567/13) = \nu(3^4 \times 7/13) = 4$.

Now suppose we have some other element $b \in \mathbb{Z}_{(p)}$ with $b \neq 0$. We can write this in the form $b = p^s x/y$ where p does not divide x or y , so $\nu(b) = s$. We then have $ab = p^{t+s}(vx)/(wy)$, from which it is not hard to see that $\nu(ab) = t + s = \nu(a) + \nu(b)$. Given this, condition (c) is clear.

Condition (b) is also satisfied, in a rather trivial way. If $t < s$ then we just put $q = 0$ and $r = a$ and we have $a = qb + r$ with $\nu(r) < \nu(b)$. On the other hand, if $t \geq s$ then the number $q := a/b = p^{t-s}(vy)/(wx)$ lies in $\mathbb{Z}_{(p)}$ and $a = bq$ so we can take $r = 0$.

For the rest of this section, we let R denote a Euclidean domain, with Euclidean valuation ν say.

Theorem 9.7. [thm-pid]

Every ideal I in R is principal.

Proof. Let I be an ideal; we must find an element $b \in R$ such that $I = Rb$. First, if $I = \{0\}$ then $I = R0$ as required; so we may assume that $I \neq \{0\}$. Each nonzero element $b \in I$ has a valuation $\nu(b) \in \mathbb{Z}$ with $\nu(b) \geq 0$. Choose such an element for which $\nu(b)$ is as small as possible. By assumption $b \in I$ and I is an ideal so $Rb \subseteq I$.

Conversely, suppose that $a \in I$. By axiom (b), then there are elements $q, r \in R$ such that $a = bq + r$ and either $r = 0$ or $\nu(r) < \nu(b)$. Note that $r = a - bq$ and $a, b \in I$ so $r \in I$. If r were a nonzero element of I with $\nu(r) < \nu(b)$, this would contradict our choice of b . We must therefore have $r = 0$ instead, so $a = bq$, so $a \in Rb$. This proves that $I \subseteq Rb$ and thus that $I = Rb$. \square

Now suppose we have two elements $a, b \in R$. Then $Ra + Rb$ is an ideal in R , so by the theorem there must be an element d such that $Ra + Rb = Rd$. This raises the question of how to find d explicitly.

The first thing to note is that $Ra + Rb$ is *not* the same as $R(a + b)$. For example, take $R = \mathbb{Z}$ and $a = 3$ and $b = 2$. We can write any number n as $n = n \times 3 + (-n) \times 2$ so $n \in \mathbb{Z}.3 + \mathbb{Z}.2$, which shows that $\mathbb{Z}.3 + \mathbb{Z}.2 = \mathbb{Z}$. However, $\mathbb{Z}.(2 + 3)$ consists only of the multiples of 5, so it is not the same.

Definition 9.8. [defn-divisible]

Let a and b be elements of R . We say that a is *divisible* by b if and only if there is an element $c \in R$ such that $a = bc$, or equivalently if $a \in Rb$, or equivalently if $Ra \subseteq Rb$. We also write $b|a$ if a is divisible by b .

Definition 9.9. [defn-gcd]

Let a and b be elements of R . A *common divisor* of a and b is an element $d \in R$ such that a and b are both divisible by d . A *greatest common divisor* (or *gcd*) of a and b is an element d such that

- (i) d is a common divisor of a and b ; and
- (ii) if d' is any other common divisor then d is divisible by d' .

Proposition 9.10. [prop-gcd]

We have $Ra + Rb = Rd$ if and only if d is a gcd of a and b . If so, then d can be written as $xa + yb$ for some $x, y \in R$.

Proof. First suppose that $Ra + Rb = Rd$. We can write a as $1a + 0b$, so $a \in Ra + Rb = Rd$, so a is divisible by d . Similarly b is divisible by d , so d is a common divisor of a and b .

Next, it is clear that $d \in Rd$, so $d \in Ra + Rb$, so $d = xa + yb$ for some $x, y \in R$.

Finally, suppose that d' is another common divisor of a and b . Then $a = u'd'$ and $b = v'd'$ for some $u', v' \in R$. This means that $d = xa + yb = xu'd' + yv'd' = (xu' + yv')d'$, so d is divisible by d' . This proves that d is a gcd of a and b .

Conversely, suppose that d is a gcd of a and b . We know from Theorem 9.7 that $Ra + Rb = Rc$ for some $c \in R$, and we know from the first half of this proof that c must also be a gcd of a and b . As d is a greatest common divisor and c is a common divisor, we see that d is divisible by c , so $Rd \subseteq Rc$. As c is a greatest common divisor and d is a common divisor, we see that c is divisible by d , so $Rc \subseteq Rd$, so $Rd = Rc = Ra + Rb$ as required. \square

We now explain how to actually find the gcd of two elements $a, b \in R$. The answer is to use the Euclidean algorithm, just as in \mathbb{Z} or $\mathbb{R}[x]$. We may assume that $a, b \neq 0$ (otherwise the problem is trivial). We then define $a_0 = a$ and $b_0 = b$. By the definition of a Euclidean valuation, we can find $p_0, a_1 \in R$ such that $a_0 = p_0b_0 + a_1$ and either $a_1 = 0$ or $\nu(a_1) < \nu(b_0)$. (Informally, a_1 is the remainder when we divide a_0 by b_0 .) Assuming that $a_1 \neq 0$, we can then find q_1, b_1 such that $b_0 = q_1a_1 + b_1$ and either $b_1 = 0$ or $\nu(b_1) < \nu(a_1)$. Assuming that $b_1 \neq 0$ we can find p_1, a_2 such that $a_1 = p_1b_1 + a_2$ and either $a_2 = 0$ or $\nu(a_2) < \nu(b_1)$. Assuming that $a_2 \neq 0$, we can then find q_2, b_2 such that $b_1 = q_2a_2 + b_2$ and either $b_2 = 0$ or $\nu(b_2) < \nu(a_2)$. Continuing in this way, we get a sequence

$$\nu(b_0) > \nu(a_1) > \nu(b_1) > \nu(a_2) > \nu(b_2) > \dots$$

Now, all these valuations are nonnegative integers so they cannot keep decreasing forever. Thus, after a finite number of steps we must end up with either $a_k = 0$ or $b_k = 0$, forcing the process to stop.

Suppose that the first term to be zero is a_k , so the elements a_0, \dots, a_{k-1} and b_0, \dots, b_{k-1} are all nonzero. I claim that $Ra + Rb = Rb_{k-1}$, so that b_{k-1} is a gcd of a and b .

To see this, put $I = Ra + Rb$ and $J = Rb_{k-1}$; we must show that $I = J$. Certainly $a_0, b_0 \in I$. Using the equation $a_1 = a_0 - p_0b_0$ we deduce that $a_1 \in I$. Using the equation $b_1 = b_0 - q_1a_1$ we deduce that $b_1 \in I$. Using the equation $a_2 = a_1 - p_1b_1$ we deduce that $a_2 \in I$. Continuing in this way, we see that $a_i, b_i \in I$ for all i and in particular that $b_{k-1} \in I$, so $J \subseteq I$.

We now use a similar argument in the opposite direction. Clearly $b_{k-1} \in J$. We have $a_{k-1} = p_{k-1}b_{k-1} + a_k$ but $a_k = 0$ so $a_{k-1} = p_{k-1}b_{k-1} \in J$. We can now use the equation $b_{k-2} = q_{k-1}a_{k-1} + b_{k-1}$ to show that $b_{k-2} \in J$, and then use the equation $a_{k-2} = p_{k-2}b_{k-2} + a_{k-1}$ to show that $a_{k-2} \in J$. Working backwards in this way, we eventually find that $a_0, b_0 \in J$ or in other words $a, b \in J$. This implies that $ua + vb \in J$ for all $u, v \in R$, or in other words that $I = Ra + Rb \subseteq J$. We have already seen that $J \subseteq I$, so $I = J$ as required.

All this assumed that the first term to be zero was a_k . It could instead happen that the first term to be zero was b_k , in which case a very similar argument would show that $Ra + Rb = Ra_k$, so a_k is a gcd of a and b .

Exercises

Exercise 9.1. [ex-roots-of-unity]

Let n and m be coprime positive integers, and put $f(x) = x^n - 1$ and $g(x) = x^m - 1$. By considering the roots of f and g , show that the gcd of f and g in $\mathbb{C}[x]$ is $x - 1$.

Exercise 9.2. [eg-p-local-gcd]

Let p be a prime, and let a and b be nonzero elements of $\mathbb{Z}_{(p)}$. Show that either a is a gcd of a and b in $\mathbb{Z}_{(p)}$, or b is a gcd of a and b in $\mathbb{Z}_{(p)}$.

10. FACTORISATION IN EUCLIDEAN DOMAINS

Let R be a Euclidean domain. We say that an element $a \in R$ is a *nonunit* if it is not invertible in R . We say that an element is *reducible* if it can be written as a product of two nonunits. We say that it is *irreducible* if it is a nonunit but is not reducible.

Note that 0 is a nonunit and $0 = 0 \cdot 0$ so 0 is a product of two nonunits and thus is reducible. For the rest of this section, we exclude the element 0 from consideration.

Example 10.1. [eg-prime-numbers]

Let p be a prime number. Then the only ways to factor p in \mathbb{Z} are $p = 1 \cdot p = (-1) \cdot (-p) = p \cdot 1 = (-p) \cdot (-1)$. In each case, one of the factors is either 1 or -1 and thus is invertible in \mathbb{Z} . Thus, p is irreducible. Similarly, $-p$ is irreducible, but if $n > 0$ and n is composite then n and $-n$ are reducible.

Example 10.2. [eg-irreducible-poly]

Consider the ring $\mathbb{C}[x]$. I claim that the irreducible elements are precisely the polynomials of the form $ax + b$ with $a \neq 0$. Firstly, polynomials of degree 0 are invertible (because we have excluded the zero polynomial from consideration). Thus, any nonunit has degree at least 1 , so any reducible polynomial has degree at least 2 . Thus when $a \neq 0$ the polynomial $ax + b$ is a nonunit but not reducible, so it is irreducible.

Next, let $f(x)$ be any polynomial of degree $d > 1$. By the Fundamental Theorem of Algebra, f has a root, in other words there is a complex number a such that $f(a) = 0$. This means that f is divisible by $x - a$, say $f(x) = (x - a)g(x)$ for some polynomial g . Note that g has degree $d - 1 > 0$, so both $g(x)$ and $x - a$ are nonunits, so f is reducible. This shows that the irreducibles are precisely the polynomials of degree exactly one, as claimed.

Example 10.3. [eg-p-local-irreducible]

Any element in $x \in \mathbb{Z}_{(p)}$ can be written as $x = p^v a/b$ where a and b are not divisible by p and so a/b is a unit in $\mathbb{Z}_{(p)}$. In other words, every element is a unit multiple of p^v for some $v \geq 0$. Using this we find that x is a unit iff $v = 0$, that x is irreducible iff $v = 1$, and that x is reducible iff $v > 1$.

Definition 10.4. [defn-associate]

We write $a \sim b$ if and only if there is an invertible element $u \in R$ such that $au = b$. If so, we say that a is a *unit multiple* or *associate* of b . It is not hard to see that $a \sim b$ if and only if $Ra = Rb$, and that the relation \sim is an equivalence relation.

Note that in \mathbb{Z} , both 7 and -7 are irreducible, but they are unit multiples of each other so for many purposes it makes no sense to use both of them. It is thus traditional to ignore -7 . Similarly, in $\mathbb{C}[x]$ both $x - 2$ and $2x - 4$ are irreducible but they are unit multiples of each other, so we usually ignore $2x - 4$. This leads us to make the following definitions.

Definition 10.5. [defn-comp-irr]

A *complete set of irreducibles* in a Euclidean domain R is a set \mathcal{P} of irreducibles such that for every irreducible p , there is a unique irreducible $p' \in \mathcal{P}$ such that $p' \sim p$.

Note that for any Euclidean domain, we can always choose a complete set of irreducibles. We simply divide the set of all irreducibles up into equivalence classes under the relation \sim , and we pick one irreducible from each equivalence class.

Example 10.6. [eg-complete-set]

The set of positive prime numbers is a complete set of irreducibles in \mathbb{Z} . The set $\{x - a \mid a \in \mathbb{C}\}$ is a complete set of irreducibles in $\mathbb{C}[x]$. The set $\{p\}$ (with just one element) is a complete set of irreducibles in $\mathbb{Z}_{(p)}$.

Proposition 10.7. [prop-Rp-field]

If p is irreducible in R then R/p is a field.

Proof. As p is not a unit we see that 1 is not divisible by p , so $\bar{1} \neq \bar{0}$ in R/p .

Every nonzero element of R/p has the form $\bar{a} = a + Rp$, where $a \in R$ but $a \notin Rp$. We then have $Ra + Rp = Rd$, where d is a gcd of a and p . Thus d divides a and p , say $a = ud$ and $p = vd$. As p is irreducible, one of v and d must be a unit. If v is a unit we have $a = ud = uv^{-1}p$ so $a \in Rp$, contrary to assumption. Thus d must be a unit instead, so $1 \in Rd = Ra + Rp$, so $1 = xa + yp$ for some $x, y \in R$. This means that $\bar{x}\bar{a} = \bar{1}$ in R/p , so \bar{a} is invertible as required. \square

Corollary 10.8. [cor-irr-prime]

If a and b are not divisible by p , then ab is not divisible by p .

Proof. $a + Rp$ and $b + Rp$ are nonzero elements of the field R/p , so $ab + Rp = (a + Rp)(b + Rp)$ is nonzero, so p does not divide ab . \square

Lemma 10.9. [lem-nu-greater]

If a and b are nonzero and b is a nonunit then $\nu(ab) > \nu(a)$.

Proof. Axiom (c) for Euclidean valuations says that $\nu(ab) \geq \nu(a)$. It is thus enough to suppose that $\nu(ab) = \nu(a)$ and deduce a contradiction. We can divide a by ab with remainder to get $a = abq + r$ for some q, r with either $r = 0$ or $\nu(r) < \nu(ab) = \nu(a)$. Note that $a(1 - bq) = r$. As b is not a unit we cannot have $bq = 1$, so $1 - bq \neq 0$. We also have $a \neq 0$ so $r \neq 0$, so $\nu(r) < \nu(a)$. However, Axiom (c) also says that $\nu(a(1 - bq)) \geq \nu(a)$, giving the required contradiction. \square

Lemma 10.10. [lem-nu-zero]

If $\nu(a) = 0$ then a is invertible.

Proof. Divide 1 by a to get $1 = qa + r$ with $r = 0$ or $\nu(r) < \nu(a)$. As valuations are always nonnegative we cannot have $\nu(r) < \nu(a)$ so we must have $r = 0$, so $1 = qa$, so q is an inverse for a . \square

Theorem 10.11. [thm-ufd]

Let R be a Euclidean domain, and let \mathcal{P} be a complete set of irreducibles in R . Then any nonzero element $a \in R$ can be written in the form $a = up_1^{n_1} \dots p_r^{n_r}$, where p_1, \dots, p_r are distinct irreducibles in \mathcal{P} and $n_1, \dots, n_r \in \mathbb{N}$ and u is invertible. Moreover, this factorization is unique (except that it could be written in a different order, for example $3^2 \times 5^3 = 5^3 \times 3^2$).

Proof. We first show that any nonzero element $a \in R$ can be written as a unit times a product of standard irreducibles, by induction on $\nu(a)$. If $\nu(a) = 0$ then a is a unit, which we think of as a unit times the product of the empty list of standard irreducibles. If $\nu(a) > 0$ then a is a nonunit. If a is irreducible then it has the form up , where u is a unit and p is a standard irreducible. Otherwise, a can be written as a product of two nonunits, say $a = bc$. By Lemma 10.9 we see that $\nu(b)$ and $\nu(c)$ are strictly less than $\nu(a)$. By induction we may assume that b and c can be written as units times products of standard irreducibles, and it follows that the same is true of a . By collecting factors together, we can write $a = up_1^{n_1} \dots p_r^{n_r}$, where p_1, \dots, p_r are distinct irreducibles in \mathcal{P} and $n_1, \dots, n_r \in \mathbb{N}$ and u is invertible.

Suppose we have another such factorization $a = vq_1^{m_1} \dots q_s^{m_s}$, where q_1, \dots, q_s are distinct irreducibles in \mathcal{P} and $m_1, \dots, m_s \in \mathbb{N}$ and v is invertible. I claim that $p_1 = q_j$ for some j . If not, then all the elements q_j would be indivisible by p_1 , as would v , and Corollary 10.8 would tell us that $v \prod_j q_j^{m_j}$ is indivisible by p_1 , so a is indivisible by p_1 , which is clearly false. Thus $p_1 = q_j$ for some j , and the same argument shows that each p_i is a q_j , and similarly that each q_j is a p_i . Thus, after renumbering the q 's if necessary, we may assume that $r = s$ and $p_i = q_i$ for all i .

Now suppose that $n_1 \leq m_1$. Put $b = up_1^{n_1} \dots p_r^{n_r}$ and $c = vp_1^{m_1 - n_1} p_2^{m_2} \dots p_r^{m_r}$. We then have $p_1^{n_1} b = a$ and $p_1^{n_1} c = a$ and $p_1^{n_1} \neq 0$ so $b = c$. Using Corollary 10.8 again we see that b is not divisible by p_1 so c is not divisible by p_1 so $m_1 - n_1 = 0$ so $m_1 = n_1$. A similar argument works if $n_1 \geq m_1$, so $n_1 = m_1$ in all cases. The same method shows that $n_i = m_i$ for all i , and given this, it is clear that $u = v$ as well. \square

11. FINITE FREE MODULES OVER A EUCLIDEAN DOMAIN

Throughout this section, we let R denote a Euclidean domain.

Definition 11.1. [defn-finite-free]

A finite free module over R is an R -module M that is isomorphic to R^d for some nonnegative integer d .

To understand this definition more explicitly, we introduce the notion of a basis.

Definition 11.2. [defn-basis]

Let M be an R -module. We say that a list $\{m_1, \dots, m_d\}$ of elements of M is a *basis* if for every element $m \in M$ there is precisely one list $(u_1, \dots, u_d) \in R^d$ such that $m = u_1m_1 + \dots + u_dm_d$.

Proposition 11.3. [prop-basis]

Let M be an R -module. Then M is a finite free module if and only if it has a basis.

Proof. Suppose that $\{m_1, \dots, m_d\}$ is a basis. As in example 6.7, we can define a homomorphism $\phi: R^d \rightarrow M$ by $\phi(u_1, \dots, u_d) = u_1m_1 + \dots + u_dm_d$. By the definition of a basis, for each $m \in M$ there is precisely one element $u \in R^d$ such that $\phi(u) = m$. This means that ϕ is a bijection, and thus an isomorphism. Thus $M \simeq R^d$, so M is free.

Conversely, suppose that M is free, so we have an isomorphism $\phi: R^d \rightarrow M$ for some d . By the second half of Example 6.7, there is a list m_1, \dots, m_d of elements of M such that $\phi(u_1, \dots, u_d) = u_1m_1 + \dots + u_dm_d$ for all $(u_1, \dots, u_d) \in R^d$. As ϕ is an isomorphism, for each element $m \in M$ there is a unique element $u \in R^d$ with $\phi(u) = m$, and this means precisely that $\{m_1, \dots, m_d\}$ is a basis. \square

Example 11.4. [eg-sub-free-i]

Put $M = \{(x, y, z) \in \mathbb{Z}^3 \mid x = y\}$. Then the vectors $m_1 := (1, 1, 0)$ and $m_2 := (0, 0, 1)$ clearly lie in M . Moreover, any vector in M can be written uniquely in the form $(x, x, z) = xm_1 + zm_2$. Thus $\{m_1, m_2\}$ is a basis for M as a \mathbb{Z} -module, so M is free.

Example 11.5. [eg-sub-free-ii]

Put $M = \{(x, y, z) \in \mathbb{Z}^3 \mid x + y + z = 0 \pmod{3}\}$, considered as a module over \mathbb{Z} . I claim that this is free as a \mathbb{Z} -module. To prove this, we need to find some elements of M which can form a basis. Firstly, if $x + y + z = 0$ then (x, y, z) will certainly lie in M . Some obvious vectors satisfying $x + y + z = 0$ are $m_1 := (1, -1, 0)$ and $m_2 := (0, 1, -1)$. Also, the vector $m_3 := (0, 0, 3)$ satisfies $x + y + z = 3$ so $x + y + z = 0 \pmod{3}$ so $m_3 \in M$. We can thus define a map $\phi: \mathbb{Z}^3 \rightarrow M$ by $\phi(u_1, u_2, u_3) = u_1m_1 + u_2m_2 + u_3m_3$.

Now suppose we have an element $m = (x, y, z) \in M$. Note that $x + y + z = 0 \pmod{3}$, so $x + y + z = 3t$ for some $t \in \mathbb{Z}$. We have

$$m - xm_1 = (x, y, z) - x(1, -1, 0) = (0, x + y, z),$$

and so

$$m - xm_1 - (x + y)m_2 = (0, x + y, z) - (x + y)(0, 1, -1) = (0, 0, x + y + z) = (0, 0, 3t) = tm_3,$$

so $m = xm_1 + (x + y)m_2 + tm_3 = \phi(x, x + y, t)$. This shows that ϕ is surjective.

Now suppose that $(u, v, w) \in \ker(\phi)$, so $um_1 + vm_2 + wm_3 = 0$. By looking at the x coordinates we see that $u \cdot 1 + v \cdot 0 + w \cdot 0 = 0$ so $u = 0$. Thus, the original equation becomes $vm_2 + wm_3 = 0$. By looking at the y coordinates we see that $v \cdot 1 + w \cdot 0 = 0$, so $v = 0$, so the equation becomes $wm_3 = 0$. By looking at the z coordinates we see that $3w = 0$ but w is just an integer so the only way $3w$ can be 0 is if $w = 0$. This proves that $\ker(\phi) = \{(0, 0, 0)\}$, so ϕ is injective and thus is an isomorphism.

We next give a convenient test for showing that certain modules are *not* free.

Definition 11.6. [defn-torsion]

For any R -module M , an element $m \in M$ is a *torsion element* if there is some nonzero element $a \in R$ such that $am = 0$. We write $\text{tors}(M)$ for the set of all torsion elements. We say that M is *torsion-free* if $\text{tors}(M) = \{0\}$. Equivalently, a module M is torsion-free if whenever $a \in R$ and $x \in M$ are both nonzero, their product ax is also nonzero.

Lemma 11.7. [lem-free-torsion-free]

Every finite free module is torsion-free.

Proof. Every finite free module is isomorphic to R^d for some d , so it is enough to show that R^d is torsion-free. Let a be a nonzero element of R , and let $u = (u_1, \dots, u_d)$ be a nonzero element of R^d . This means that $u_i \neq 0$ for some i . As R is an integral domain, it follows that $au_i \neq 0$, and thus that the vector $au = (au_1, \dots, au_d)$ is not the zero vector. \square

Example 11.8. [eg-Z-torsion]

Suppose that $n > 1$, and consider \mathbb{Z}_n as a \mathbb{Z} -module. Then $\bar{1}$ is a nonzero element of \mathbb{Z}_n and n is a nonzero element of \mathbb{Z} but $n\bar{1} = \bar{n} = \bar{0}$. Thus \mathbb{Z}_n is not torsion-free as a \mathbb{Z} -module, so it cannot be a finite free module.

Example 11.9. Consider $C^\infty(\mathbb{R}, \mathbb{R})$ as a module over $\mathbb{R}[D]$ in the usual way. Then the function $f(t) = e^t$ is a nonzero element of $C^\infty(\mathbb{R}, \mathbb{R})$, and $D - 1$ is a nonzero element of $\mathbb{R}[D]$, but $(D - 1)f = f' - f = 0$. Thus $C^\infty(\mathbb{R}, \mathbb{R})$ is not torsion-free, so it is not a finite free module over $C^\infty(\mathbb{R}, \mathbb{R})$. For an even simpler proof, just consider the constant function $g(t) = 1$, so $Dg = g' = 0$, again showing that $C^\infty(\mathbb{R}, \mathbb{R})$ is not torsion-free.

We now prove a key theorem, which makes the theory of modules over Euclidean domains much simpler than for other rings.

Theorem 11.10. [thm-hered]

Any submodule of a finite free module over a Euclidean domain is free.

Note that Examples 11.4 and 11.5 illustrate this.

Proof. As every finite free module is isomorphic to R^d for some d , it will be enough to show that every submodule of R^d is free. We do this by induction on d .

The case $d = 0$ is easy. The module R^0 is just $\{0\}$, the only submodule of this is $\{0\}$ itself, and this is free because it is R^0 .

The case $d = 1$ is essentially given by Theorem 9.7. Let M be a submodule of $R^1 = R$, or in other words an ideal in R . By Theorem 9.7 we have $M = Ra$ for some $a \in R$. If $a = 0$ then $M = \{0\} = R^0$ so M is free. If $a \neq 0$ then I claim that $\{a\}$ is a basis for M . Indeed, as $M = Ra$, every element $m \in M$ can certainly be written as $m = ua$ for some $u \in R$. If we have $ua = m = va$ then $(u - v)a = 0$ but $a \neq 0$ and R is an integral domain so $u - v = 0$ so $u = v$. Thus m can be written *uniquely* in the form $m = ua$, so $\{a\}$ is a basis and so M is free.

Now suppose that $d > 1$ and that we have proved that every submodule of R^{d-1} is free. We need to show that every submodule $M \leq R^d$ is free. Let F be the set of vectors of the form $(x_1, x_2, \dots, x_{d-1}, 0)$ in R^d . Note that F is a copy of R^{d-1} , so every submodule of F is free. In particular, $M \cap F$ is a submodule of F so it is free, with basis $\{m_1, \dots, m_r\}$ say. Define a homomorphism $\pi: M \rightarrow R$ by $\pi(x_1, \dots, x_d) = x_d$. The image of π is a submodule of R so it has the form Ra for some $a \in R$.

Suppose that $a = 0$. Then $\text{image}(\pi) = Ra = \{0\}$, so $\pi(m) = 0$ for all $m \in M$, so every element of M has the form $(x_1, \dots, x_{d-1}, 0)$. This means that every element of M is an element of F , so M is a submodule of F , so M is free.

Suppose instead that $a \neq 0$. As $a \in Ra = \text{image}(\pi)$, we can choose $m_{r+1} \in M$ such that $\pi(m_{r+1}) = a$. I claim that $\{m_1, \dots, m_r, m_{r+1}\}$ is a basis for M . To see this, let m be an element of M . Then $\pi(m) \in \text{image}(\pi) = Ra$, so $\pi(m) = u_{r+1}a$ for some $u_{r+1} \in R$. Put $m' = m - u_{r+1}m_{r+1}$, so $m' \in M$ and

$$\pi(m') = \pi(m) - u_{r+1}\pi(m_{r+1}) = u_{r+1}a - u_{r+1}a = 0.$$

This means that the last coordinate of m' is 0, so $m' \in F$. As $m' \in M \cap F$ and $\{m_1, \dots, m_r\}$ is a basis for $M \cap F$ we see that $m' = u_1m_1 + \dots + u_rm_r$ for some $u_1, \dots, u_r \in R$, so

$$m = m' + u_{r+1}m_{r+1} = u_1m_1 + \dots + u_{r+1}m_{r+1}.$$

This shows that the elements m_1, \dots, m_{r+1} generate M .

Now suppose that we have elements $v_1, \dots, v_{r+1} \in R$ satisfying $v_1m_1 + \dots + v_{r+1}m_{r+1} = 0$. I claim that $v_1 = \dots = v_{r+1} = 0$. Indeed, as $m_1, \dots, m_r \in F$ we have $\pi(m_1) = \dots = \pi(m_r) = 0$, so when we apply π to the previous equation we get $v_{r+1}\pi(m_{r+1}) = 0$, or equivalently $v_{r+1}a = 0$. As $a \neq 0$ and R is an integral domain this means that $v_{r+1} = 0$. Thus, our original equation becomes $v_1m_1 + \dots + v_rm_r = 0$. As $\{m_1, \dots, m_r\}$ is a basis for $M \cap F$, the only way we can have $v_1m_1 + \dots + v_rm_r = 0$ is if $v_1 = \dots = v_r = 0$, as claimed.

It now follows that $\{m_1, \dots, m_{r+1}\}$ is a basis for M , so M is free. This completes the inductive step, and thus the proof of the theorem. \square

Corollary 11.11. [cor-Noether]

Let M be a finitely generated module over a Euclidean domain R , and let N be a submodule of M . Then N is also finitely generated.

Proof. As M is finitely generated, there is a list m_1, \dots, m_d of elements of M such that an arbitrary element $m \in M$ can be written in the form $u_1m_1 + \dots + u_dm_d$. Define $\phi: R^d \rightarrow M$ by $\phi(u_1, \dots, u_d) = u_1m_1 + \dots + u_dm_d$; this is clearly a surjective homomorphism. Put $L = \{u \in R^d \mid \phi(u) \in N\}$. I claim that this is a submodule of R^d . Indeed, if $u, v \in L$ then $\phi(u), \phi(v) \in N$ so $\phi(u+v) = \phi(u) + \phi(v) \in N$ so $u+v \in L$. Similarly, if $u \in L$ and $a \in R$ then $\phi(u) \in N$ so $\phi(au) = a\phi(u) \in N$ so $au \in L$, so L is a submodule as claimed. Submodules of R^d are finite free modules by the theorem, so we can choose a basis $\{p_1, \dots, p_r\}$ for L . Put $n_i = \phi(p_i)$; as $p_i \in L$ we have $n_i \in N$. I claim that the elements n_1, \dots, n_r generate N . Indeed, suppose $n \in N$. Then $n \in M$ and the homomorphism $\phi: R^d \rightarrow M$ is surjective so we have $n = \phi(u)$ for some $u \in R^d$. As $\phi(u) = n \in N$ we see that $u \in L$, so u can be written in the form $u = v_1p_1 + \dots + v_rp_r$ for some $v_1, \dots, v_r \in R$. It follows that

$$n = \phi(u) = v_1\phi(p_1) + \dots + v_r\phi(p_r) = v_1n_1 + \dots + v_rn_r.$$

This shows that the elements n_1, \dots, n_r generate N as claimed, so N is finitely generated. \square

Exercises

Exercise 11.1. [ex-find-bases]

Find bases over \mathbb{Z} for the following submodules of \mathbb{Z}^3 . Justify your answers.

- (a) $M_0 = \{(x, y, z) \mid x - y + z = 0 \pmod{5}\}$
- (b) $M_1 = \{(x, y, z) \mid x = y \pmod{2} \text{ and } y = z \pmod{3}\}$
- (c) $M_2 = \{(x, y, z) \mid 6x + 15y + 10z = 0\}$.

Exercise 11.2. [ex-split-quotient]

Let d_1, \dots, d_n be elements of a ring R , and let N be the submodule of R^n generated by the elements d_1e_1, \dots, d_ne_n . Prove that $R^n/N \simeq R/d_1 \oplus \dots \oplus R/d_n$. [You may wish to start by defining a homomorphism $\alpha: R^n \rightarrow R/d_1 \oplus \dots \oplus R/d_n$.]

Exercise 11.3. [ex-compatible-bases]

Put

$$F = \{(w, x, y, z) \in \mathbb{Z}^4 \mid w + x + y + z \text{ is even}\}$$

$$G = \{(w, x, y, z) \in \mathbb{Z}^4 \mid w - x, x - y, \text{ and } y - z \text{ are divisible by } 4\}.$$

Find integers $d_1, d_2, d_3, d_4 > 0$ and vectors $u_1, u_2, u_3, u_4 \in \mathbb{Z}^4$ such that $\{u_1, u_2, u_3, u_4\}$ is a basis for F and $\{d_1u_1, d_2u_2, d_3u_3, d_4u_4\}$ is a basis for G . [It is possible to do this using matrix methods, but intelligent trial and error is likely to be easier.] It follows that $G \subseteq F$, so we can form the factor group F/G . Deduce a description of F/G as a direct sum of cyclic groups.

12. ROW AND COLUMN OPERATIONS

We saw in the last section that a submodule of a finite free module over a Euclidean domain is free. We next give a systematic method for finding a basis for such a submodule.

Suppose we have vectors u_1, \dots, u_n in R^m , and we let N be the submodule generated by u_1, \dots, u_n . It will be convenient to form the $n \times m$ matrix A whose columns are the vectors u_i .

Definition 12.1. [defn-col-ops]

An *elementary column operation* on a matrix A over a ring R is any of the following operations:

- (a) Add a multiple of one column to another column
- (b) Multiply a column by an invertible element of R
- (c) Exchange two columns.

We say that a matrix A is in (unreduced) column echelon form if

- (a) All nonzero columns occur to the left of all the zero columns.

- (b) If the i 'th and $(i + 1)$ 'st columns are nonzero, then the top nonzero entry in the $(i + 1)$'s column is below the top nonzero entry in the i 'th column.

Example 12.2. The matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 \\ 3 & 6 & 0 & 0 \\ 4 & 7 & 9 & 0 \\ 5 & 8 & 10 & 0 \end{bmatrix}$$

is in column echelon form, but the following matrices are not

$$\begin{bmatrix} 0 & 3 & 0 \\ 1 & 4 & 0 \\ 2 & 5 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 3 & 5 \end{bmatrix} \quad \begin{bmatrix} 0 & 0 & 8 \\ 0 & 4 & 0 \\ 2 & 0 & 0 \end{bmatrix}.$$

Proposition 12.3. [prop-col-red]

Let A be an $n \times m$ matrix over a Euclidean domain R , and let N be the submodule of R^m generated by the columns of A . Then A can be reduced by elementary column operations to a matrix B in column echelon form, and the nonzero columns of B form a basis for N .

Remark 12.4. [rem-unreduced-echelon]

The resulting matrix B is not unique; in general, a matrix A can be converted to many different matrices B in column echelon form.

The algorithm is a mixture of the Euclidean algorithm and the usual Gaussian algorithm for column-reducing a matrix over a field. Suppose that the top row of A is nonzero (if not, we simply ignore any rows of zeros at the top and work with the first nonzero row). Of all the nonzero elements in the top row, we find one whose valuation is as small as possible. After swapping the columns around, we can assume that this element occurs in the top left corner of the matrix, so it is the entry a_{11} . If we were working over a field, we would subtract a_{k1}/a_{11} times the first column from the k 'th column (for $k = 2, \dots, n$), and then the top row would have the form $(a_{11}, 0, \dots, 0)$. As we are not working over a field, we do not necessarily have $a_{k1}/a_{11} \in R$, so we cannot perform these operations. As the next best thing, we divide a_{k1} by a_{11} by the division algorithm to obtain $a_{k1} = a_{11}q_k + r_k$ with either $r_k = 0$ or $\nu(r_k) < \nu(a_{11})$. We then subtract q_k times the first column from the k 'th column for $k = 2, \dots, n$. The top row is now $(a_{11}, r_2, \dots, r_n)$. If $r_2 = \dots = r_n = 0$ then we have a matrix of the form $\left(\begin{array}{c|c} a_{11} & 0 \\ * & A' \end{array} \right)$. We can then column-reduce the smaller matrix A' by the same process, and eventually we get a column-reduced form for A . Generally, however, the elements r_2, \dots, r_n will not all be zero. Among those that are nonzero, we choose one whose valuation is as small as possible. Note that by construction this valuation is strictly less than that of a_{11} and valuations are always nonnegative integers, so this kind of step can only occur finitely many times. We swap the columns around to put the element of minimum valuation in the top left corner and repeat the whole process.

We end up with a matrix B in column echelon form. Let N' be the submodule of R^m generated by the columns of B . It is clear from the form of B that its columns are linearly independent, so they form a basis for N' . To prove Proposition 12.3, it will be enough to check that $N' = N$, or equivalently that when we perform an elementary column operation on a matrix, the module spanned by the columns does not change. We will explain this by example rather than giving a formal proof. Suppose that the matrix has three columns, say u_1, u_2 and u_3 .

- (a) A typical operation of the first type replaces the list (u_1, u_2, u_3) by $(u_1, u_2 + cu_1, u_3)$ for some $c \in R$. If a vector v can be written as $a_1u_1 + a_2u_2 + a_3u_3$, it can also be written as $(a_1 - ca_2)u_1 + a_2(u_2 + cu_1) + a_3u_3$, so $\text{span}\{u_1, u_2, u_3\} \subseteq \text{span}\{u_1, u_2 + cu_1, u_3\}$. Conversely, if v can be written as $b_1u_1 + b_2(u_2 + cu_1) + b_3u_3$, it can also be written as $(b_1 + cb_2)u_1 + b_2u_2 + b_3u_3$, so $\text{span}\{u_1, u_2 + cu_1, u_3\} \subseteq \text{span}\{u_1, u_2, u_3\}$.
- (b) A typical operation of the second type replaces the list (u_1, u_2, u_3) by (u_1, cu_2, u_3) for some invertible element $c \in R$. If a vector v can be written as $a_1u_1 + a_2u_2 + a_3u_3$, it can also be written as $a_1u_1 + (a_2c^{-1})(cu_2) + a_3u_3$, so $\text{span}\{u_1, u_2, u_3\} \subseteq \text{span}\{u_1, cu_2, u_3\}$. Conversely, if v can be written

as $b_1u_1 + b_2(cu_2) + b_3u_3$, it can also be written as $b_1u_1 + (b_2c)u_2 + b_3u_3$, so $\text{span}\{u_1, cu_2, u_3\} \subseteq \text{span}\{u_1, u_2, u_3\}$.

- (c) A typical operation of the third type replaces the list (u_1, u_2, u_3) by (u_2, u_1, u_3) , and this clearly does not change the span.

A formal proof would be essentially the same but would need more elaborate notation.

Example 12.5. [eg-col-red-i]

Here we perform a column reduction over \mathbb{Z} by strictly following the algorithm.

$$\begin{aligned} \begin{bmatrix} 5 & 8 & 11 & 3 \\ 16 & 25 & 34 & 9 \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} 3 & 5 & 8 & 11 \\ 9 & 16 & 25 & 34 \end{bmatrix} \\ &\xrightarrow{2} \begin{bmatrix} 3 & 2 & 2 & 2 \\ 9 & 7 & 7 & 7 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 2 & 3 & 2 & 2 \\ 7 & 9 & 7 & 7 \end{bmatrix} \\ &\xrightarrow{4} \begin{bmatrix} 2 & 1 & 0 & 0 \\ 7 & 2 & 0 & 0 \end{bmatrix} \\ &\xrightarrow{5} \begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 7 & 0 & 0 \end{bmatrix} \\ &\xrightarrow{6} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 3 & 0 & 0 \end{bmatrix} \end{aligned}$$

In step 1 we note that the element of smallest valuation in the top row is the 3 in the 4'th column. We therefore move the 4'th column round to be the first column. In step 2 we divide the remaining entries in the top row by 3. We have $5 = 1 \times 3 + 2$ and $8 = 2 \times 3 + 2$ and $11 = 3 \times 3 + 2$ so we subtract 1 times the first column from the second column, 2 times the first column from the third column, and 3 times the first column from the fourth column. In step 3 we note that the element of smallest valuation on the top row is 2, so we swap the first two columns to put a 2 in the top left corner. Then in step 4, we subtract the first column from each of the other columns. The element of smallest valuation in the top row is now 1, so we swap the first two columns to put the 1 in the top left corner; this is step 5. Finally, we subtract twice the first column from the second column. Our matrix is now in the form $\left(\begin{array}{c|c} a_{11} & 0 \\ * & A' \end{array} \right)$, and the matrix $A' = (3 \ 0 \ 0)$ is already in echelon form so we are finished.

If we use a little imagination rather than strictly following the algorithm, we can reduce the matrix more quickly:

$$\begin{aligned} \begin{bmatrix} 5 & 8 & 11 & 3 \\ 16 & 25 & 34 & 9 \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} -1 & -1 & -1 & 3 \\ -2 & -2 & -2 & 9 \end{bmatrix} \\ &\xrightarrow{2} \begin{bmatrix} -1 & 0 & 0 & 0 \\ -2 & 0 & 0 & 3 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 3 & 0 & 0 \end{bmatrix}. \end{aligned}$$

The conclusion is that if N is the submodule of \mathbb{Z}^2 generated by the vectors $(5, 16)$, $(8, 25)$, $(11, 34)$ and $(3, 9)$ then $\{(1, 2), (0, 3)\}$ is a basis for N .

Example 12.6. [eg-col-red-ii]

Here is a column reduction over $\mathbb{Q}[x]$.

$$\begin{aligned} \begin{bmatrix} x^2 & x-1 & x^3 \\ 0 & -1 & x \\ 0 & -1 & 0 \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} 1 & x-1 & 1 \\ x+1 & -1 & x^2+2x+1 \\ x+1 & -1 & x^2+x+1 \end{bmatrix} \\ &\xrightarrow{2} \begin{bmatrix} 1 & 0 & 0 \\ x+1 & -x^2 & x^2+x \\ x+1 & -x^2 & x^2 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 1 & 0 & 0 \\ x+1 & x^2 & x \\ x+1 & x^2 & 0 \end{bmatrix} \\ &\xrightarrow{4} \begin{bmatrix} 1 & 0 & 0 \\ x+1 & x & 0 \\ x+1 & 0 & x^2 \end{bmatrix} \end{aligned}$$

In step 1 we subtracted $x+1$ times the middle column from the first column, and x^2+x+1 times the middle column from the third column. In step 2 we subtracted $x-1$ times the first column from the middle column, and subtracted the first column from the last column. In step 3 we multiplied the middle column by -1 and then subtracted it from the last column. In step 4 we subtracted x times the last column from the middle column, and then swapped the middle and last columns.

The conclusion is that the vectors $m_1 := (1, x+1, x+1)$, $m_2 := (0, x, 0)$ and $m_3 := (0, 0, x^2)$ form a basis for the submodule of $\mathbb{Q}[x]^3$ generated by the columns of the original matrix.

In fact, the columns of the original matrix also form a basis for this submodule (just because they are linearly independent over $\mathbb{Q}[x]$). However, the basis $\{m_1, m_2, m_3\}$ is easier to work with because it is in echelon form.

Example 12.7. [eg-power-vecs]

Put $v_k = (k, k^2, k^3, k^4) \in \mathbb{Z}^4$, and let N be the subgroup of \mathbb{Z}^4 generated by v_1, \dots, v_5 . These vectors are the columns of the first matrix written below:

$$\begin{aligned} \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 9 & 16 & 25 \\ 1 & 8 & 27 & 64 & 125 \\ 1 & 16 & 81 & 256 & 625 \end{bmatrix} &\rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 6 & 12 & 20 \\ 1 & 6 & 24 & 60 & 120 \\ 1 & 14 & 78 & 252 & 620 \end{bmatrix} \\ &\rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 \\ 1 & 6 & 6 & 24 & 60 \\ 1 & 14 & 36 & 168 & 480 \end{bmatrix} \\ &\rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 \\ 1 & 6 & 6 & 0 & 0 \\ 1 & 14 & 36 & 24 & 0 \end{bmatrix} \end{aligned}$$

We deduce that the vectors $(1, 1, 1, 1)$, $(0, 2, 6, 14)$, $(0, 0, 6, 36)$ and $(0, 0, 0, 24)$ form a basis for N . Note that the “leading terms” of these vectors are $1!$, $2!$, $3!$ and $4!$; this is part of a general pattern.

The above algorithm helps us to find a basis for a submodule N of R^n . However, we often want instead to investigate the structure of R^n/N , and so far we do not have a method for this. The simplest case is where N is generated by the elements d_1e_1, \dots, d_re_r for some $r \leq n$ and some nonzero elements d_1, \dots, d_r in R , where e_i is the i 'th standard basis vector. It is then easy to see that

$$R^n/N \simeq R/d_1 \oplus \dots \oplus R/d_r \oplus R^{n-r}.$$

More generally, suppose we can find a (non-standard) basis u_1, \dots, u_n for R^n and nonzero elements d_1, \dots, d_r such that N is generated by d_1u_1, \dots, d_ru_r . We again find that $R^n/N \simeq R/d_1 \oplus \dots \oplus R/d_r \oplus R^{n-r}$. Our next algorithm will enable us to find bases of this type.

Definition 12.8. [defn-normal-form]

A matrix over a Euclidean domain is in *normal form* if it has the form $\left(\begin{array}{c|c} D & 0 \\ \hline 0 & 0 \end{array}\right)$, where

- (1) D is an $r \times r$ matrix for some r (called the *rank*)
- (2) The diagonal entries in D are nonzero, and the other entries are zero
- (3) If the diagonal entries are d_1, \dots, d_r , then $d_1 | d_2 | \dots | d_r$.

For example, the following matrix over \mathbb{Z} is in normal form:

$$\left(\begin{array}{ccc|cc} 2 & 0 & 0 & 0 & 0 \\ 0 & 6 & 0 & 0 & 0 \\ 0 & 0 & 12 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 \end{array}\right)$$

We have already defined elementary column operations over a Euclidean domain, and elementary row operations are defined in the obvious analogous way.

Theorem 12.9. [thm-gauss]

Let A be an $n \times m$ matrix over a Euclidean domain R , and let N be the quotient of R^m by the span of the columns of A . Then A can be transformed by row and column operations to a matrix B in normal form. Moreover, if the nonzero diagonal entries in B are d_1, \dots, d_r then $N \simeq R/d_1 \oplus \dots \oplus R/d_r \oplus R^{m-r}$.

The proof will follow after some auxiliary definitions and preliminary results.

Remark 12.10. [rem-mod-unit]

It could happen that some of the elements d_i are units. In this case we have $Rd_i = R$ and so $R/d_i = R/R = 0$ so we can drop the term R/d_i from the direct sum.

Definition 12.11. [defn-nu-A]

Let R be a Euclidean domain with valuation ν . For any nonzero matrix A over R , we let $\nu(A)$ be the smallest of the valuations of all the nonzero entries in A . For example, if $R = \mathbb{Z}$ we have $\nu \begin{bmatrix} 5 & -4 \\ 0 & -3 \end{bmatrix} = \nu(-3) = 3$.

Definition 12.12. We say that a nonzero $n \times m$ matrix A over R is *prenormal* if it has the form $\left(\begin{array}{c|c} d & 0 \\ \hline 0 & B \end{array}\right)$, where

- (1) d is a nonzero element of R
- (2) B is an $(n-1) \times (m-1)$ matrix over R
- (3) every entry in B is divisible by d .

Lemma 12.13. [lem-prenormal]

Let A be a nonzero matrix over a Euclidean domain R . Then A can be transformed into prenormal form by elementary row and column operations.

Proof. Let a_{ij} be the entry in the i 'th column and j 'th row of A . Put $n = \nu(A)$, which is a nonnegative integer. By definition we have $\nu(a_{ij}) = n$ for some i, j , and after reordering the rows and columns we may assume that $\nu(a_{11}) = n$. Put $d = a_{11}$.

Suppose that every entry in A is divisible by d . Then for each $i > 1$ we have $a_{1i} = q_i d$ for some $q_i \in R$. We can subtract q_i times the top row from the i 'th row for each i to get a new matrix in which every row except the first starts with 0. It is easy to see that in this new matrix, every entry is still divisible by d . We can then subtract multiples of the first column from the other columns to get a matrix of the form $\left(\begin{array}{c|c} d & 0 \\ \hline 0 & B \end{array}\right)$

where every entry in B is divisible by d . Thus A can be made prenormal, as claimed.

Now suppose instead that some entry in A is not divisible by d . I claim that A can be transformed to a new matrix A' with $\nu(A') < n = \nu(A)$. Indeed, if some entry a_{1i} in the first column is not divisible by d then we have $a_{1i} = dq + r$ for some nonzero remainder r with $\nu(r) < \nu(d) = n$. If we subtract q times the first row from the i 'th row we get a new matrix A' in which the i 'th row starts with r , so $\nu(A') \leq \nu(r) < n$. A similar argument works if some entry a_{j1} in the first row is not divisible by d . This leaves the case where all entries in the first row or the first column are divisible by d , but some entry a_{ij} (with $i, j > 1$) is not

divisible by d . After reordering some rows and columns we may assume that a_{22} is not divisible by d . We thus have $a_{22} = dq + r$ where $r \neq 0$ and $\nu(r) < \nu(d) = n$. We also have $a_{12} = ud$ and $a_{21} = vd$ for some u, v . The top left corner of our matrix looks like this:

$$\begin{bmatrix} a & ud \\ vd & qd + r \end{bmatrix}.$$

We now subtract v times the first row from the second row, then add the second row to the top row, then subtract $(q - uv + u)$ times the first column from the second column. The effect on the top left corner is as follows:

$$\begin{bmatrix} d & ud \\ vd & qd + r \end{bmatrix} \rightarrow \begin{bmatrix} d & ud \\ 0 & (q - uv)d + r \end{bmatrix} \rightarrow \begin{bmatrix} d & (q - uv + u)d + r \\ 0 & (q - uv)d + r \end{bmatrix} \rightarrow \begin{bmatrix} d & r \\ 0 & (q - uv)d + r \end{bmatrix}.$$

Thus, the matrix A' that we end up with has r as an entry, so $\nu(A') \leq \nu(r) < n$ as claimed.

We now repeat the whole process. We find that either A' can be made prenormal, or it can be transformed to another matrix A'' with $\nu(A'') < \nu(A')$, and so on. As valuations of matrices are nonnegative integers, we cannot keep on reducing them indefinitely, so eventually we must get to a matrix that can be made prenormal. \square

Proposition 12.14. [prop-gauss]

Any matrix A over a Euclidean domain R can be transformed to normal form by row and column operations.

Proof. Let A_0 be a matrix over a Euclidean domain, of shape $n \times m$ say. If $A_0 = 0$ then A_0 is already in normal form. Otherwise, by the Proposition, we can convert A_0 by row and column operations to the form $\left(\begin{array}{c|c} d_1 & 0 \\ \hline 0 & A_1 \end{array} \right)$, where every entry in A_1 is divisible by d_1 . Clearly A_1 has shape $(n - 1) \times (m - 1)$. If A_1 is nonzero we can convert it to the form $\left(\begin{array}{c|c} d_2 & 0 \\ \hline 0 & A_2 \end{array} \right)$, where the entries in A_2 are divisible by d_2 . As this is obtained from A_1 by row and column operations, we see that all the entries are still divisible by d_1 , and in particular $d_1|d_2$. This converts A_0 to the form

$$\left(\begin{array}{cc|c} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ \hline 0 & 0 & A_2 \end{array} \right).$$

The theorem follows by iterating this process in the evident way. \square

Example 12.15. [eg-gauss]

$$\begin{bmatrix} 13 & 3 \\ 7 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 6 & 1 \\ 7 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 6 & 1 \\ -5 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 5 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix}.$$

This row reduction follows the method implicit in the above proof, except that we have not bothered to reorder the columns until the end. We start by finding the entry of smallest valuation, which is 2. We subtract the bottom row from the top row to make the entry above the 2 have smaller valuation than 2 does. Now the 1 in the top right hand corner has smaller valuation than everything else, and everything is divisible by 1. We can thus use row and column operations to clear away the rest of the entries in the same row or column as the 1. We then swap the columns to put the matrix in normal form.

Example 12.16.

$$\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 3 \\ 0 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 \\ -6 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 6 \end{bmatrix}.$$

Here we start with a diagonal matrix whose entry of smallest valuation is in the 2 in the top left corner. However, the other entries are not all divisible by 2, so the matrix is not prenormal. We first perform a row operation to ensure that there is a non-divisible entry on the top row. We then subtract 2 times the first column from the second column, leaving a remainder of 1 in the top right corner, which has valuation smaller than 2. We can now clear the entries below and to the left of the 1 and then reorder to get a normal matrix.

Example 12.17. [eg-power-vecs-ii]

Put $v_k = (k, k^2, k^3, k^4) \in \mathbb{Z}^4$, and let N be the subgroup of \mathbb{Z}^4 generated by v_1, \dots, v_5 . We showed in Example 12.7 that the corresponding matrix can be column-reduced as follows:

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 9 & 16 & 25 \\ 1 & 8 & 27 & 64 & 125 \\ 1 & 16 & 81 & 256 & 625 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 \\ 1 & 6 & 6 & 0 & 0 \\ 1 & 14 & 36 & 24 & 0 \end{bmatrix}$$

We can now perform row operations to reduce this matrix further. This works easily in this particular example, because the entries underneath the 1 are divisible by 1, the entries under the 2 are divisible by 2, and the entry under the 6 is divisible by 6. Explicitly, we subtract the first row from each of the other rows; then we subtract 3 times the second row from the third row, and 7 times the second row from the last row; then we subtract 6 times the third row from the last row. This gives the matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 6 & 0 & 0 \\ 0 & 0 & 0 & 24 & 0 \end{bmatrix}.$$

The conclusion will be that $\mathbb{Z}^4/N \simeq \mathbb{Z}_2 \oplus \mathbb{Z}_6 \oplus \mathbb{Z}_{24}$. (We have omitted the \mathbb{Z}_1 because $\mathbb{Z}_1 = 0$.)

Definition 12.18. [defn-elementary-matrix]

Let E be an $n \times n$ matrix over a ring R . We say that E is *elementary* if either

- (1) It is obtained from the identity matrix by exchanging two rows (or equivalently, exchanging two columns); or
- (2) The diagonal entries are all equal to 1, there is a single nonzero entry off the diagonal, and all other entries are zero; or
- (3) One of the diagonal entries is an invertible element of R , the remaining diagonal entries are equal to 1, and all other entries are zero.

Note that elementary matrices are always invertible, and that their inverses are also elementary matrices.

Just as in the case of fields, if A' is obtained from A by a single elementary row operation then $A' = EA$ for some elementary matrix E . For example, let A be an $n \times 3$ -matrix. Then

- (1) Exchanging the first and third rows is the same as multiplying on the left by $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$.
- (2) Adding the third row to the first is the same as multiplying on the left by $\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.
- (3) Multiplying the second row by -1 is the same as multiplying the matrix on the left by $\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

Similarly, if A' is obtained from A by a single column operation then $A' = AE$ for some elementary matrix E . Thus, if A' is obtained from A by a sequence of row and column operations then $A' = PAQ$ for some matrices P and Q which are products of elementary matrices and thus are invertible. Using this, the following corollary follows immediately from Proposition 12.14

Corollary 12.19. [cor-gauss]

If A is an $n \times m$ matrix over a Euclidean domain, then there exist invertible square matrices P and Q (of size m and n respectively) such that PAQ is in normal form. \square

Lemma 12.20. [lem-standard-presentation]

Let F be a free module over R , and let u_1, \dots, u_m be a basis for F . Suppose that $r \leq m$ and $d_1, \dots, d_r \in R$. Let M be the submodule of F generated by d_1u_1, \dots, d_ru_r . Then $F/M \simeq R/d_1 \oplus \dots \oplus R/d_r \oplus R^{m-r}$.

Proof. Write $Q = R/d_1 \oplus \dots \oplus R/d_r \oplus R^{m-r}$ for brevity. Any element $x \in F$ can be written uniquely in the form $x_1u_1 + \dots + x_mu_m$ with $x_1, \dots, x_m \in R$. It therefore makes sense to define a map $\alpha: F \rightarrow Q$ by

$$\alpha(x_1u_1 + \dots + x_mu_m) = (x_1 + Rd_1, \dots, x_r + Rd_r, x_{r+1}, \dots, x_m).$$

This is easily seen to be a homomorphism. Suppose we have an element $y = (y_1, \dots, y_m) \in Q$, so $y_i \in R/d_i$ for $i \leq r$ and $y_i \in R$ for $i > r$. For $i \leq r$ we can choose $x_i \in R$ such that $y_i = x_i + Rd_i$. We can then

define $u = (x_1, \dots, x_r, y_{r+1}, \dots, y_m) \in R^m$ and we find that $\alpha(u) = y$. Thus α is surjective, and the First Isomorphism Theorem tells us that $Q \simeq F/\ker(\alpha)$.

We now need to understand $\ker(\alpha)$. If $x = \sum_i x_i u_i \in \ker(\alpha)$ then $(x_1 + Rd_1, \dots, x_r + Rd_r, x_{r+1}, \dots, x_m) = (0, \dots, 0)$ which means that $x_i = 0$ for $i > r$ and $x_i \in Rd_i$ for $i \leq r$. This means that for $i \leq r$ we have elements $y_i \in R$ such that $x_i = y_i d_i$ and so

$$x = x_1 u_1 + \dots + x_r u_r = y_1 (d_1 u_1) + \dots + y_r (d_r u_r),$$

so $x \in \text{span}\{d_1 u_1, \dots, d_r u_r\} = M$. This shows that $\ker(\alpha) \subseteq M$, and the reverse inclusion is easy so $\ker(\alpha) = M$. We thus have $Q \simeq F/M$ as claimed. \square

Proof of Theorem 12.9. Let A be an $n \times m$ matrix over a Euclidean domain R , let M be the submodule of R^m spanned by the columns of A , and put $N = R^m/M$. Choose P and Q as in Corollary 12.19, so the matrix $C := PAQ$ is in normal form, with diagonal entries d_1, \dots, d_r say. Put $B = AQ$. This is obtained from A by elementary column operations, so the columns of B span the same submodule as the columns of A ; in other words, they span N .

Next, note that $PB = C$ so $B = P^{-1}C$. Let u_1, \dots, u_m be the columns of P^{-1} ; as P^{-1} is invertible, these form a basis for R^m . I claim that the columns of $P^{-1}C$ are $d_1 u_1, \dots, d_r u_r, 0, \dots, 0$ (with $n - r$ zeros). This is clear in the following special case, where $n = 4$, $m = 3$ and $r = 2$:

$$P^{-1} = \begin{bmatrix} u_{11} & u_{21} & u_{31} \\ u_{12} & u_{22} & u_{32} \\ u_{13} & u_{23} & u_{33} \end{bmatrix}$$

$$C = \begin{bmatrix} d_1 & 0 & 0 & 0 \\ 0 & d_2 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$P^{-1}C = \begin{bmatrix} d_1 u_{11} & d_2 u_{21} & 0 & 0 \\ d_1 u_{12} & d_2 u_{22} & 0 & 0 \\ d_1 u_{13} & d_2 u_{23} & 0 & 0 \end{bmatrix}$$

The general case is the same except for more elaborate notation.

Thus M is spanned by $d_1 u_1, \dots, d_r u_r$, and it follows that $N = R^m/M$ is isomorphic to $R/d_1 \oplus \dots \oplus R/d_r \oplus R^{m-r}$. \square

Corollary 12.21. [cor-class]

Let M be a finitely generated module over a Euclidean domain R . Then

$$M \simeq R/d_1 \oplus \dots \oplus R/d_r \oplus R^s$$

for some $r, s \geq 0$ and some nonunits $d_1, \dots, d_r \in R$ with $d_1 | d_2 | \dots | d_r$.

Proof. As M is finitely generated, we can find a finite list v_1, \dots, v_m of elements that generate M . We can then define a homomorphism $\alpha: R^m \rightarrow M$ by

$$\alpha(a_1, \dots, a_m) = a_1 v_1 + \dots + a_m v_m.$$

This is surjective because the elements v_i span M . Thus, the First Isomorphism Theorem for modules tells us that $M \simeq R^m/\ker(\alpha)$. We also know that $\ker(\alpha)$ is a submodule of R^m and thus is a finite free module, with basis u_1, \dots, u_n say. Let A be the $n \times m$ matrix with columns u_1, \dots, u_n , so M is isomorphic to the quotient of R^m by the span of the columns of A . The claim now follows \square

Corollary 12.22. [cor-fg-Ab]

Let M be a finitely generated Abelian group. Then

$$M \simeq \mathbb{Z}_{d_1} \oplus \dots \oplus \mathbb{Z}_{d_r} \oplus \mathbb{Z}^s$$

for some $r, s \geq 0$ and some natural numbers d_1, \dots, d_r with $1 < d_1 | d_2 | \dots | d_r$. \square

Exercise 12.1. [ex-col-red-i]

Reduce the following matrix over $\mathbb{C}[x]$ to column echelon form.

$$A = \begin{bmatrix} x-1 & x & x+1 \\ x & 0 & x \\ x+1 & x & x-1 \end{bmatrix}.$$

Exercise 12.2. [ex-gauss-i]

Consider the following matrix over $\mathbb{C}[x]$:

$$A = \begin{bmatrix} x & 0 & -1 \\ -1 & x & -1 \\ 0 & -1 & x+1 \end{bmatrix}.$$

Let M be the quotient of $\mathbb{C}[x]^3$ by the span of the columns of A .

- Reduce A to normal form by row and column operations.
- Give a polynomial $f(x)$ such that $M \simeq \mathbb{C}[x]/f(x)$.
- Give a list of basic $\mathbb{C}[x]$ -modules whose direct sum is isomorphic to M .

Exercise 12.3. [ex-gauss-ii]

Consider the following matrix over $\mathbb{C}[x]$:

$$A = \begin{bmatrix} x & 0 & 1 \\ 1 & x & 0 \\ 0 & 1 & x \end{bmatrix}.$$

Let M be the quotient of $\mathbb{C}[x]^3$ by the span of the columns of A .

- Reduce A to normal form by row and column operations.
- Give a polynomial $f(x)$ such that $M \simeq \mathbb{C}[x]/f(x)$.

Exercise 12.4. [ex-gauss-iii]

Consider the following matrix over $\mathbb{C}[x]$:

$$A = \begin{bmatrix} x & 0 & 0 \\ 0 & x-1 & 0 \\ 0 & 0 & x^2-x \end{bmatrix}.$$

Let M be the quotient of $\mathbb{C}[x]^3$ by the span of the columns of A .

- Reduce A to normal form by row and column operations.
- Describe M as a direct sum of cyclic modules over $\mathbb{C}[x]$.

Exercise 12.5. [ex-gauss-iv]

Let M be the subgroup of \mathbb{Z}^5 generated by the columns of the following matrix.

$$A = \begin{bmatrix} 1 & 1 & 2 & 3 & 3 \\ 1 & -1 & 0 & 1 & -1 \\ 1 & 1 & 2 & -1 & -1 \\ 1 & 1 & -1 & 0 & 0 \\ 1 & -1 & 0 & -1 & 1 \end{bmatrix}$$

Determine the structure of \mathbb{Z}^5/M .

[Readers familiar with representation theory may be interested to know that A is the character table of the symmetric group S_4 . This means that $\mathbb{Z}^5/M = C/R$, where R is the character ring of S_4 and C is the ring of integer-valued class functions. You can do the question without knowing any of this, however.]

Exercise 12.6. [ex-col-red-primes]

Reduce the following matrix over \mathbb{Z} to column echelon form.

$$A = \begin{bmatrix} 71 & 97 & 113 & 149 \\ 1 & 1 & 1 & 1 \end{bmatrix}.$$

(The entries in the top row are the 20'th, 25'th, 30'th and 35'th prime numbers. You could actually derive the answer from this fact, but it is probably easier to just do the calculation.)

Exercise 12.7. [ex-det-deg]

- Let A be an $n \times n$ matrix over \mathbb{Z} , and suppose that B is obtained from A by row and column operations. Show that $|\det(A)| = |\det(B)|$.
- Let A be an $n \times n$ matrix over \mathbb{Z} , and suppose that $|\det(A)| = d \neq 0$. Let M be the quotient of \mathbb{Z}^n by the span of the columns of A . Show that M is a finite Abelian group of order d .
- Let A be an $n \times n$ matrix over $\mathbb{C}[x]$, with $\det(A) = f(x) \neq 0$. Let M be the quotient of $\mathbb{C}[x]^n$ by the span of the columns of A . Show that $\dim_{\mathbb{C}}(M)$ is the degree of the polynomial $f(x)$.

Exercise 12.8. [ex-hilbert-matrix]

Let A be the 3×3 matrix whose (i, j) 'th entry is $1/(i + j - 1)$, as shown below:

$$A = \begin{bmatrix} 1 & 1/2 & 1/3 \\ 1/2 & 1/3 & 1/4 \\ 1/3 & 1/4 & 1/5 \end{bmatrix}.$$

This is called a *Hilbert matrix*. Put $B = 120A$ (so the entries in B are integers) and let M be the quotient of \mathbb{Z}^3 by the span of the columns of B . Describe M as a direct sum of cyclic groups.

Exercise 12.9. [ex-cubic-span]

Let M be the subgroup of \mathbb{Z}^2 generated by the vectors (k, k^3) for $k = 2, 3, 4, 5$. Find a basis for M over \mathbb{Z} , and determine the order of \mathbb{Z}^2/M .

Exercise 12.10. [ex-thirty]

Let A be the following matrix over \mathbb{Z} :

$$A = \begin{bmatrix} 6 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & 15 \end{bmatrix}.$$

- Reduce A to normal form by row and column operations over \mathbb{Z} .
- What does the answer tell you about the group $\mathbb{Z}_6 \oplus \mathbb{Z}_{10} \oplus \mathbb{Z}_{15}$?
- Can you prove this fact in an easier way?

13. PRIMARY DECOMPOSITION

Corollary 12.21 gives us in some sense a complete list of all finitely generated R -modules. However, it turns out not to be the most convenient kind of list to work with. Moreover, it still leaves us with a question about uniqueness. For example, both $\mathbb{Z}_4 \oplus \mathbb{Z}_{12}$ and $\mathbb{Z}_2 \oplus \mathbb{Z}_{24}$ appear in the list of finitely generated Abelian groups, and so far we have no way of knowing whether they might be isomorphic. In this section we will modify the classification given in Corollary 12.21 to get a new classification which is more useful in practice and which allows us to prove a uniqueness theorem.

Let R be a Euclidean domain, and choose a complete set of irreducibles \mathcal{P} .

Let M be an R -module. Recall that in Definition 11.6 we defined $\text{tors}(M)$ to be the set of torsion elements in M , in other words the set of elements $m \in M$ such that there is a nonzero element $a \in R$ with $am = 0$.

Proposition 13.1. [prop-tors-submodule]
 $\text{tors}(M)$ is a submodule of M .

Proof. Suppose that $m, n \in \text{tors}(M)$. Then there are nonzero elements $a, b \in R$ such that $am = 0$ and $bn = 0$. It follows that $ab \neq 0$ and $ab(m+n) = b(am) + a(bn) = b \cdot 0 + a \cdot 0 = 0$, so $m+n \in \text{tors}(M)$. Similarly, for any $c \in R$ we have $a(cm) = c(am) = 0$, so $cm \in \text{tors}(M)$, so $\text{tors}(M)$ is a submodule as claimed. \square

Definition 13.2. [defn-torsion-module]

We say that M is a *torsion module* if $\text{tors}(M) = M$. We say that M is a *finite torsion module* if it is finitely generated as well as being a torsion module.

Definition 13.3. A *basic* R -module is an R -module of the form R/p^k for some $p \in \mathcal{P}$ and $k > 0$.

We will show that any finite torsion module is isomorphic to a direct sum of basic modules.

Example 13.4. [eg-torsion-group]

For any $m \in \mathbb{Z}_{12}$ we have $12m = 0$, so m is a torsion element. This shows that \mathbb{Z}_{12} is a torsion module over \mathbb{Z} . More generally, let M be any finite Abelian group, considered as a \mathbb{Z} -module. If d is the order of M then $dm = 0$ for all $m \in M$, which shows that M is a torsion module. We saw in Example 5.25 that M is also finitely generated, so it is a finite torsion module.

Example 13.5. [eg-poly-torsion]

Let W_2 be the set of functions of the form $f(t) = a + bt + ct^2$, considered as an $\mathbb{R}[D]$ -module in the usual way. For a function f of this form we have $f'(t) = b + 2ct$ so $f''(t) = 2c$ so $f'''(t) = 0$. This means that $D^3f = 0$, so f is a torsion element of W . It follows that W_2 is a torsion module. We saw in Example 5.26 that W_2 is a cyclic module over $\mathbb{R}[D]$ and thus is finitely generated, so it is a finite torsion module over $\mathbb{R}[D]$. Similarly, the space W_d of polynomials of degree at most d is a finite torsion module over $\mathbb{R}[D]$.

Example 13.6. [eg-QZ-torsion]

Consider the Abelian group $M = \mathbb{Q}/\mathbb{Z}$ as a \mathbb{Z} -module. Any element $m \in M$ has the form $a/b + \mathbb{Z}$ for some $a, b \in \mathbb{Z}$ with $b \neq 0$. It follows that $bm = a + \mathbb{Z}$, but $a \in \mathbb{Z}$ so $a + \mathbb{Z} = \mathbb{Z}$, which is the zero element of the group M . Thus $bm = 0$, proving that m is a torsion element. It follows that M is a torsion module. It is not finitely generated, however.

Example 13.7. [eg-eigen-torsion]

Let K be a field, and let λ and μ be elements of K . Consider K^2 as a module over $K[x]$ using the endomorphism $x.(u, v) = \phi(u, v) = (\lambda u, \mu v)$. Then

$$(x - \lambda).(u, v) = (\lambda u, \mu v) - (\lambda u, \lambda v) = (0, (\mu - \lambda)v),$$

and

$$(x - \mu).(u, v) = (\lambda u, \mu v) - (\mu u, \mu v) = ((\lambda - \mu)u, 0),$$

so

$$(x - \mu)(x - \lambda).(u, v) = (x - \mu).(0, (\mu - \lambda)v) = (0, 0).$$

In other words, the element $p(x) = (x - \mu)(x - \lambda) \in K[x]$ satisfies $p(x).(u, v) = (0, 0)$ for all $(u, v) \in K^2$, which proves that K^2 is a torsion module.

Example 13.8. [eg-fin-dim-torsion]

More generally, let V be any module over $K[x]$ that is finite-dimensional (of dimension d say) when considered as a vector space over K . I claim that V is a finite torsion module over $K[x]$. Indeed, suppose that $v \in V$. We then have vectors v, xv, x^2v, \dots, x^dv in V . There are $d+1$ vectors in this list, but V only has dimension d , so the vectors in our list must be linearly dependent. Thus, there are scalars $a_0, \dots, a_d \in K$ (not all zero) such that $a_0v + a_1xv + \dots + a_dx^dv = 0$. This means that the polynomial $f(x) = a_0 + a_1x + \dots + a_dx^d$ is nonzero and satisfies $f.v = 0$, so v is a torsion element. This shows that V is a torsion module. Moreover, we can choose a basis v_1, \dots, v_d for V as a vector space, and it follows that these elements generate V as a module over $K[x]$, so V is finitely generated.

Next recall that for any two rings R_0, R_1 we can form the product ring $R_0 \times R_1$. The elements of $R_0 \times R_1$ are pairs (a_0, a_1) with $a_0 \in R_0$ and $a_1 \in R_1$. Addition and multiplication are defined in the obvious way:

$$\begin{aligned}(a_0, a_1) + (b_0, b_1) &= (a_0 + b_0, a_1 + b_1) \\ (a_0, a_1)(b_0, b_1) &= (a_0b_0, a_1b_1).\end{aligned}$$

The additive identity is the element $(0, 0)$, and the multiplicative identity is the element $(1, 1)$.

The following result is called the *Chinese remainder theorem*.

Proposition 13.9. [prop-chinese]

If a and b are coprime then $R/ab \simeq R/a \times R/b$ as rings (or as R -modules).

Proof. Define $\alpha: R \rightarrow R/a \times R/b$ by $\alpha(t) = (t + aR, t + bR)$. Note that $\alpha(s + t) = \alpha(s) + \alpha(t)$ and $\alpha(st) = \alpha(s)\alpha(t)$ and $\alpha(1) = 1$, so α is a homomorphism of rings.

As a and b are coprime, we have $xa + yb = 1$ for some $x, y \in R$.

Suppose that $\alpha(t) = (0, 0)$. Then $t + aR$ is the zero coset $0 + aR$, so t is divisible by a , say $t = au$ for some u . Similarly $t = bv$ for some v . This means that

$$t = 1.t = (xa + yb)t = xat + ybt = xa(bv) + yb(au) = (xv + yu)ab,$$

so t is divisible by ab . Conversely, if t is divisible by ab then it is divisible by both a and b , so $\alpha(t) = (0, 0)$. Thus $\ker(\alpha) = Rab$.

Now suppose we have some element $(u + aR, v + bR) \in R/a \times R/b$. Consider the element $t = ybu + xav \in R$. Note that $t = (1 - xa)u + xav = u + xa(v - u) = u \pmod{a}$ and $t = ybu + (1 - yb)v = v + yb(u - v) = v \pmod{b}$, so $\alpha(t) = (t + Ra, t + Rb) = (u + Ra, v + Rb)$. This shows that α is surjective. Thus, Theorem 8.10 says that $R/ab \simeq R/a \oplus R/b$. \square

Corollary 13.10. [cor-basic-split]

Any finite torsion R -module is isomorphic to a direct sum of basic R -modules.

Proof. Let M be a finite torsion R -module. By Corollary 12.21 we have $M \simeq R/d_1 \oplus \dots \oplus R/d_r \oplus R^s$ say. If $s > 0$ then M contains a copy of R so it cannot be a torsion module, contrary to assumption. Thus we must have $s = 0$ and so $M = R/d_1 \oplus \dots \oplus R/d_r$. It will thus be enough to show that R/d is a direct sum of basic modules for all $d \neq 0$. For this we factor d as $up_1^{n_1} \dots p_r^{n_r}$ as in Theorem 10.11. As $Rd = Rdu^{-1}$ we may replace d by du^{-1} and thus assume that $d = p_1^{n_1} \dots p_r^{n_r}$. Put $q_i = p_i^{n_i}$ and $b_i = q_i q_{i+1} \dots q_r$, so $b_1 = d$. Note that $d = q_1 b_2$ and q_1 and b_2 are coprime. Thus Proposition 13.9 tells us that $R/d \simeq R/q_1 \oplus R/b_2$. Similarly, $b_2 = q_2 b_3$ and q_2 and b_3 are coprime so $R/b_2 \simeq R/q_2 \oplus R/b_3$ so $R/d = R/q_1 \oplus R/q_2 \oplus R/b_3$. Continuing in the obvious way we find that $R/d = R/q_1 \oplus \dots \oplus R/q_r$, and the modules R/q_i are basic, as required. \square

Example 13.11. [eg-finite-abelian]

In the case $R = \mathbb{Z}$, we deduce that any finite Abelian group is a direct sum of groups of the form \mathbb{Z}/p^k where p is prime and $k > 0$. Suppose that $M = B_1 \oplus \dots \oplus B_t$ where $B_i \simeq \mathbb{Z}/p_i^{n_i}$. (Note that the primes p_i need not all be different.) We then have

$$|M| = |B_1| \dots |B_t| = p_1^{n_1} \dots p_t^{n_t}.$$

Consider for example the case where $|M| = 81 = 3^4$. Clearly all the primes p_i must be equal to 3, and $3^4 = |M| = 3^{n_1} \dots 3^{n_t} = 3^{n_1 + \dots + n_t}$ so $n_1 + \dots + n_t = 4$. Given this, it is not hard to see that the possibilities are as follows:

$$\begin{aligned}M &\simeq \mathbb{Z}_{81} \\ M &\simeq \mathbb{Z}_{27} \oplus \mathbb{Z}_3 \\ M &\simeq \mathbb{Z}_9 \oplus \mathbb{Z}_9 \\ M &\simeq \mathbb{Z}_9 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_3 \\ M &\simeq \mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_3.\end{aligned}$$

Now consider instead the case where $|M| = 36 = 2^2 3^2$. Then all the primes p_i must be either 2 or 3. The orders $|B_i| = p_i^{n_i}$ must divide 36 so we must have $n_i = 1$ or 2. Using this it is not hard to see that the

possibilities are as follows:

$$\begin{aligned} M &\simeq \mathbb{Z}_4 \oplus \mathbb{Z}_9 \\ M &\simeq \mathbb{Z}_4 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_3 \\ M &\simeq \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_9 \\ M &\simeq \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_3. \end{aligned}$$

It remains to discuss the question of uniqueness. Could it happen, for example, that the groups $A := \mathbb{Z}_9 \oplus \mathbb{Z}_9$ and $B := \mathbb{Z}_9 \times \mathbb{Z}_3 \times \mathbb{Z}_3$ are isomorphic? The answer is no, because A contains 9 elements satisfying $3a = 0$, whereas B contains 27 elements satisfying $3b = 0$. An elaboration of this argument will show that any finite torsion module can be written in an essentially unique way as a direct sum of basic modules.

Definition 13.12. [defn-f-p-k]

Suppose that $p \in \mathcal{P}$ and $k \in \mathbb{N}$. For any finite torsion module M , we define

$$F_p^k(M) = \{x \in p^{k-1}M \mid px = 0\}.$$

This is easily seen to be a submodule of M . As M is finitely generated, we see from Corollary 11.11 that $F_p^k(M)$ is finitely generated. As $px = 0$ for all $x \in F_p^k(M)$, we can regard $F_p^k(M)$ as a module over R/p . As R/p is a field, every finitely generated module over it has a well-defined dimension, so we can define

$$f_p^k(M) = \dim_{R/p}(F_p^k(M)).$$

We also define

$$g_p^k(M) = f_p^k(M) - f_p^{k+1}(M).$$

Remark 13.13. [rem-f-p-k-sum]

It is easy to see that a pair $(x, y) \in M \oplus N$ lies in $F_p^k(M \oplus N)$ if and only if $x \in F_p^k(M)$ and $y \in F_p^k(N)$. It follows that $F_p^k(M \oplus N) = F_p^k(M) \oplus F_p^k(N)$ and thus that $f_p^k(M \oplus N) = f_p^k(M) + f_p^k(N)$ and $g_p^k(M \oplus N) = g_p^k(M) + g_p^k(N)$.

Remark 13.14. [rem-f-p-k-iso]

Suppose that $M \simeq M'$. I claim that $F_p^k(M) \simeq F_p^k(M')$, and thus that $f_p^k(M) = f_p^k(M')$ and $g_p^k(M) = g_p^k(M')$. Indeed, let $\phi: M \rightarrow M'$ be an isomorphism. Then if $x \in F_p^k(M)$ then $x = p^{k-1}y$ for some y and $px = 0$, so $\phi(x) = p^{k-1}\phi(y)$ and $p\phi(x) = 0$, so $\phi(x) \in F_p^k(M')$. Thus, ϕ gives a homomorphism from $F_p^k(M)$ to $F_p^k(M')$. Similarly, the homomorphism $\phi^{-1}: M' \rightarrow M$ restricts to give a homomorphism from $F_p^k(M')$ to $F_p^k(M)$. It is easy to see that these two maps are inverse to each other, so $F_p^k(M) \simeq F_p^k(M')$ as claimed.

Proposition 13.15. [prop-g-p-k-basic]

We have

$$g_p^k(R/q^j) = \begin{cases} 1 & \text{if } p = q \text{ and } k = j \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We first prove that

$$f_p^k(R/q^j) = \begin{cases} 0 & \text{if } p \neq q \\ 0 & \text{if } p = q \text{ and } k > j \\ 1 & \text{if } p = q \text{ and } k \leq j. \end{cases}$$

First suppose that $p \neq q$. Then p^k and q^j are coprime, so $ap^k + bq^j = 1$ for some $a, b \in R$. If $x \in F_p^k(R/q^j)$ then $x = p^{k-1}y$ for some y and $px = 0$ so $p^k y = 0$. On the other hand, it is clear from the definition of R/q^j that $q^j z = 0$ for all $z \in R/q^j$, so $q^j y = 0$. We thus have $y = 1 \cdot y = ap^k y + bq^j y = 0$, and thus $x = p^{k-1}y = 0$. Thus $F_p^k(R/q^j) = 0$ and so $f_p^k(R/q^j) = 0$, as required.

Now suppose that $q = p$ and $j < k$. If $x \in R/p^j$ then $p^j x = 0$ and $k - 1 \geq j$ so $p^{k-1}x = 0$. Thus $p^{k-1} \cdot (R/p^j) = 0$ and therefore $F_p^k(R/p^j) = 0$ so $f_p^k(R/p^j) = 0$.

Now suppose instead that $q = p$ and $k \leq j$. Put $e = p^{j-1} + p^j R \in R/p^j$, so clearly $pe = 0$. We also have $e = p^{k-1}e'$, where $e' = p^{j-k} + p^j R \in R/p^j$, so $e \in p^{k-1}(R/p^j)$, so $e \in F_p^k(R/p^j)$. If \bar{a} is another element of

$F_p^k(R/p^j)$ then $p\bar{a} = 0$ so pa is a multiple of p^j so a is a multiple of p^{j-1} so \bar{a} is a multiple of e . As $e \neq 0$ and e spans $F_p^k(R/p^j)$ we conclude that $F_p^k(R/p^j)$ has dimension one, so $f_p^k(R/p^j) = 1$, as required.

It is now easy to deduce our description of $g_p^k(R/q^j)$. If $q \neq p$ then $f_p^k(R/q^j) = 0$ for all k and it follows easily that $g_p^k(R/q^j) = 0$. Suppose instead that $p = q$. If $k > j$ then $k + 1 > j$ as well so $f_p^k(R/p^j) = f_p^{k+1}(R/p^j) = 0$ so $g_p^k(R/p^j) = 0$ as claimed. If $k < j$ then both k and $k + 1$ are less than or equal to j , so $f_p^k(R/p^j) = f_p^{k+1}(R/p^j) = 1$ so $g_p^k(R/p^j) = 0$ as claimed. If $k = j$ then $f_p^k(R/p^j) = 1$ and $f_p^{k+1}(R/p^j) = 0$ so $g_p^k(R/p^j) = 1$ as claimed. \square

Corollary 13.16. [cor-primary]

Let M be an finite torsion module. Then M can be expressed in a unique way as a direct sum of basic modules. The number of copies of R/p^k in the direct sum is $g_p^k(M)$.

Proof. We know from Corollary 13.10 that M can be written as $B_1 \oplus \dots \oplus B_t$, where each B_i is a basic module. Let n_p^k be the number of copies of R/p^k in this list. We know that $g_p^k(M) = g_p^k(B_1) + \dots + g_p^k(B_t)$. In this sum we get a 1 for every B_i that is a copy of R/p^k and a 0 for all other B_i 's. Thus implies that $g_p^k(M) = n_p^k$.

Now suppose we have another splitting, say $M = C_1 \oplus \dots \oplus C_s$ where each C_j is basic. Let m_p^k be the number of copies of R/p^k in this list. The same argument as before shows that $g_p^k(M) = m_p^k$, so $m_p^k = n_p^k$. Thus the lists B_1, \dots, B_t and C_1, \dots, C_s contain the same number of copies of R/p^k for all p and k , so the two lists must be the same up to reordering. Thus, M can be written in an essentially unique way as a direct sum of basic modules. \square

Exercises

Exercise 13.1. [eg-decompose-i]

Write the $\mathbb{C}[x]$ -module $M := \mathbb{C}[x]/(x^4 - x^2) \oplus \mathbb{C}[x]/(x^4 - 2x^2 + 1)$ as a direct sum of basic $\mathbb{C}[x]$ -modules.

Exercise 13.2. [ex-FpkM-calc]

Let M be the \mathbb{Z} -module $\mathbb{Z}_2 \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_9$. List all the elements of the subgroups $F_2^1(M)$, $F_2^2(M)$ and $F_3^1(M)$.

Exercise 13.3. [ex-classify-CCV]

- (a) List all the Abelian groups of order 225 up to isomorphism. You should write all the groups as direct sums of basic \mathbb{Z} -modules.
- (b) Which of the groups in your list is isomorphic to \mathbb{Z}_{225} ?
- (c) Let M be an Abelian group of order 225. Suppose that there is an element in M of order 25, and that there are 9 elements $x \in M$ satisfying $3x = 0$. Which of the groups in your list is isomorphic to M ?

Exercise 13.4. [ex-ten-thousand]

Let p be a prime number.

- (a) List all the Abelian groups of order p^4 , up to isomorphism.
- (b) Let M be an Abelian group of order p^4 , and put $N = \{x \in M \mid px = 0\}$. Suppose that $\dim_{\mathbb{Z}/p} N = 3$. Which of the groups in your list is isomorphic to M ?
- (c) How many isomorphism classes of Abelian groups of order 10000 are there? You should justify your answer, but you need not list all the groups.

Exercise 13.5. [ex-p-fifth]

- (a) Given a prime number p , list all the isomorphism classes of Abelian groups of order p^5 .

- (b) Let M be an Abelian group of order 2^5 . Suppose that M has precisely 8 elements satisfying $2m = 0$, but that all elements satisfy $4m = 0$. Which of the groups in your list is isomorphic to M ?

Exercise 13.6. [ex-diffop-basis]

Suppose that $f \in C^\infty(\mathbb{R}, \mathbb{R})$ satisfies $f''' = f'$. Find elements $e_0, e_1, e_{-1} \in \mathbb{R}[D]$ such that

$$\begin{aligned} f &= e_{-1}f + e_0f + e_1f \\ (D - 1)e_{-1}f &= 0 \\ De_0f &= 0 \\ (D + 1)e_1f &= 0. \end{aligned}$$

Exercise 13.7. [ex-diffop-solve]

In this problem we solve the differential equation $f''' = f'$.

- Write the equation $f''' = f'$ in the form $p(D)f = 0$ for some polynomial p .
- Put $M = \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid f''' = f'\}$. Factorise $p(D)$ and thus write M as a direct sum of three submodules.
- Show that if $f''' = f'$ then there are constants $a, b, c \in \mathbb{R}$ such that $f(t) = ae^t + be^{-t} + c$ for all t .

Exercise 13.8. [ex-diffop-torsion]

Consider $C^\infty(\mathbb{R}, \mathbb{C})$ as a module over $\mathbb{C}[D]$. Let W be the space of functions of the form $p_1(t)e^{\lambda_1 t} + \dots + p_r(t)e^{\lambda_r t}$ for some $\lambda_1, \dots, \lambda_r \in \mathbb{C}$ and polynomials $p_1, \dots, p_r \in \mathbb{C}[t]$. In this (quite substantial) problem we will show that $\text{tors}(C^\infty(\mathbb{R}, \mathbb{C})) = W$.

- For each $\lambda \in \mathbb{C}$, let W_λ be the space of functions of the form $f(t) = p(t)e^{\lambda t}$, where p is polynomial. Calculate $(D - \lambda)^k f$, and deduce that W_λ is a torsion module over $\mathbb{C}[D]$.
- Suppose that $f \in C^\infty(\mathbb{R}, \mathbb{C})$ and that $(D - \lambda)^k f = 0$ for some $k \geq 0$. Prove that the function $g(t) := f(t)e^{-\lambda t}$ satisfies $D^k g = 0$.
- Suppose that a function $g \in C^\infty(\mathbb{R}, \mathbb{C})$ satisfies $D^k g = 0$. Prove by induction on k that g is a polynomial of degree less than k .
- Deduce that every function f with $(D - \lambda)^k f = 0$ lies in W_λ .
- Suppose that $f \in C^\infty(\mathbb{R}, \mathbb{C})$ and that $p(D)f = 0$ for some nonzero element $p(D) \in \mathbb{C}[D]$. By factoring $p(D)$ and considering the module $\{g \in C^\infty(\mathbb{R}, \mathbb{C}) \mid p(D)g = 0\}$, show that $f \in W_{\lambda_1} + \dots + W_{\lambda_r}$ for some r .
- Deduce that $\text{tors}(C^\infty(\mathbb{R}, \mathbb{C})) = W$.

Exercise 13.9. [ex-matrix-monomials]

Consider the following matrix over $\mathbb{C}[x]$:

$$A = \begin{bmatrix} x^3 & x^2 & x \\ x & x^2 & x \\ x & x & x \end{bmatrix}.$$

Let M be the quotient of $\mathbb{C}[x]^3$ by the span of the columns of A .

- Reduce A to normal form by row and column operations.
- Give a list of three cyclic $\mathbb{C}[x]$ -modules whose direct sum is isomorphic to M .
- Give a list of basic $\mathbb{C}[x]$ -modules whose direct sum is isomorphic to M .

14. CANONICAL FORMS FOR SQUARE MATRICES

Given any $n \times n$ matrix A over a field K we have a module M_A over $K[x]$. We see from Example 13.8 that M_A is a finite torsion module, and Corollary 13.16 gives a classification of such modules. In this section, we will see what this tells us about square matrices.

For simplicity, we will restrict attention to the case $K = \mathbb{C}$, where the irreducibles are easy to understand. As we saw in Example 10.6, the set

$$\mathcal{P} = \{x - \lambda \mid \lambda \in \mathbb{C}\}$$

is a complete set of irreducibles in $\mathbb{C}[x]$. The basic $\mathbb{C}[x]$ modules are thus the modules

$$B(\lambda, k) := B_{x-\lambda}^k = \mathbb{C}[x]/(x - \lambda)^k.$$

Thus Corollary 13.16 says that any finite torsion module over $\mathbb{C}[x]$ can be written essentially uniquely as a direct sum of $B(\lambda, k)$'s, in an essentially unique way.

The next proposition explains the most basic case of this. Recall that M_λ is the module whose elements are the complex numbers, with the multiplication rule $f.z = f(\lambda)z$.

Proposition 14.1. [prop-B-1m-one]
 $B(\lambda, 1) \simeq M_\lambda$.

Proof. Define a map $\alpha: \mathbb{C}[x] \rightarrow M_\lambda$ by $\alpha(f) = f(\lambda)$. This is a $\mathbb{C}[x]$ -module map because

$$\alpha(g.f) = (gf)(\lambda) = g(\lambda)f(\lambda) = g.f(\lambda) = g.\alpha(f).$$

If a is a constant polynomial then $\alpha(a) = a$, and this shows that α is surjective. We also have $\alpha(f) = 0$ iff $f(\lambda) = 0$ iff $f(x)$ is divisible by $x - \lambda$, so $\ker(\alpha)$ is the principal ideal $\mathbb{C}[x].(x - \lambda)$. Thus, the first isomorphism theorem gives us an isomorphism $\bar{\alpha}: B(\lambda, 1) = \mathbb{C}[x]/(x - \lambda) \rightarrow M_\lambda$. \square

Now suppose that A is a diagonalizable $n \times n$ matrix over \mathbb{C} , with eigenvalues $\lambda_1, \dots, \lambda_n$ say. Then there exists a matrix P (whose columns are eigenvectors of A) such that $D := P^{-1}AP$ is a diagonal matrix, with entries $\lambda_1, \dots, \lambda_n$ on the diagonal. Recall from Example 5.16 that the direct sum of matrices is defined by

$$A \oplus B = \left(\begin{array}{c|c} A & 0 \\ \hline 0 & B \end{array} \right).$$

If we regard λ_i as a 1×1 matrix and use this notation, we find that $D = \lambda_1 \oplus \dots \oplus \lambda_n$.

We saw in Corollary 6.19 that $M_A \simeq M_{\lambda_1} \oplus \dots \oplus M_{\lambda_n}$, and we can now rewrite this as

$$M_A \simeq B(\lambda_1, 1) \oplus \dots \oplus B(\lambda_n, 1).$$

To extend this picture to nondiagonalizable matrices, we need the following definition.

Definition 14.2. [defn-jordan-block]

Given $\lambda \in \mathbb{C}$ and $k > 0$ we let $J(\lambda, k)$ be the $k \times k$ matrix such that

- (1) Every entry on the diagonal is λ
- (2) Every entry just below the diagonal is 1
- (3) Every other entry is 0.

This is called a *Jordan block* of size k and eigenvalue λ . For example, we have

$$J(\lambda, 4) = \begin{pmatrix} \lambda & 0 & 0 & 0 \\ 1 & \lambda & 0 & 0 \\ 0 & 1 & \lambda & 0 \\ 0 & 0 & 1 & \lambda \end{pmatrix}.$$

Proposition 14.3. [prop-B-J]

$B(\lambda, k)$ is isomorphic to $M_{J(\lambda, k)}$.

Proof. Put $y = x - \lambda \in \mathbb{C}[x]$, so that $B(\lambda, k) = \mathbb{C}[x]/y^k$. Put $A = J(\lambda, k) - \lambda I$, so for $v \in M_{J(\lambda, k)} = \mathbb{C}^k$ we have $y.v = J(\lambda, k)v - \lambda v = Av$. From the definition of $J(\lambda, k)$ we see that A has 1's just below the diagonal and 0's everywhere else. In the case $k = 4$ we have

$$y.v = Av = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \begin{bmatrix} 0 \\ v_1 \\ v_2 \\ v_3 \end{bmatrix}.$$

It is not hard to see that the general case follows the same pattern, so we have

$$y.(v_1, \dots, v_k) = (0, v_1, \dots, v_{k-1}).$$

It follows that $y^2.(v_1, \dots, v_k) = (0, 0, v_1, \dots, v_{k-2})$ and so on, so $y^{k-1}.(v_1, \dots, v_k) = (0, \dots, 0, v_1)$ and $y^k v = 0$.

Let $\{e_1, \dots, e_k\}$ be the usual basis for $M_{J(\lambda, k)} = \mathbb{C}^k$ over \mathbb{C} , so $y.e_i = e_{i+1}$ for $i < k$ and $y.e_k = 0$. We define $\alpha: \mathbb{C}[x] \rightarrow M_{J(\lambda, k)}$ by $\alpha(f) = f.e_1 = f(J(\lambda, k))e_1$. This is easily seen to be a map of $\mathbb{C}[x]$ -modules.

I next claim that α is surjective. Indeed, for any $v = (v_1, \dots, v_k) \in M_{J(\lambda, k)}$ we can put $f = v_1 + v_2 y + \dots + v_k y^{k-1} \in \mathbb{C}[x]$ and we find that

$$\alpha(f) = f.e_1 = v_1 e_1 + v_2 y.e_1 + \dots + v_k y^{k-1}.e_k = v_1 e_1 + \dots + v_k e_k = v,$$

which proves surjectivity.

We next claim that $\ker(\alpha) = \mathbb{C}[x].y^k$. Indeed, suppose we have some polynomial $f(x)$ with $\alpha(f) = f.e_1 = 0$. By putting $x = y + \lambda$ and expanding everything out, we can write $f(x)$ in the form $a_0 + a_1 y + \dots + a_d y^d$. (For example, if $f(x) = x^2 + x + 1$ then $f(x) = (y + \lambda)^2 + (y + \lambda) + 1 = (1 + \lambda + \lambda^2) + (1 + 2\lambda)y + y^2$.) We then have

$$f.e_1 = a_0 e_1 + \dots + a_{k-1} y^{k-1}.e_1 = a_0 e_1 + \dots + a_{k-1} e_k = (a_0, a_1, \dots, a_{k-1}).$$

As $f.e_1 = 0$ we must have $a_0 = \dots = a_{k-1} = 0$ so $f(x) = a_k y^k + \dots + a_d y^d$, so $f(x)$ is divisible by y^k as required.

The first isomorphism theorem now tells us that

$$M_{J(\lambda, k)} = \text{image}(\alpha) \simeq \mathbb{C}[x]/\ker(\alpha) = \mathbb{C}[x]/y^k = B(\lambda, k)$$

as claimed. □

Theorem 14.4. [thm-jordan]

Any square matrix A over \mathbb{C} is conjugate to a matrix A' (called the Jordan normal form or JNF of A) that is a direct sum of Jordan blocks.

Proof. We know from Corollary 13.16 that M_A is isomorphic to a direct sum of modules of the form $B(\lambda, k)$, say

$$M_A \simeq B(\lambda_1, k_1) \oplus \dots \oplus B(\lambda_t, k_t).$$

Put $A' = J(\lambda_1, k_1) \oplus \dots \oplus J(\lambda_t, k_t)$, so

$$M_{A'} \simeq M_{J(\lambda_1, k_1)} \oplus \dots \oplus M_{J(\lambda_t, k_t)} \simeq B(\lambda_1, k_1) \oplus \dots \oplus B(\lambda_t, k_t) \simeq M_A.$$

As $M_A \simeq M_{A'}$, Proposition 6.18 tells us that A is conjugate to A' , as claimed. □

The row-reduction algorithm described previously can be used write M_A as a direct sum of cyclic modules, or in other words modules of the form $\mathbb{C}[x]/f(x)$. If we factor $f(x)$ as $(x - \lambda_1)^{k_1} \dots (x - \lambda_r)^{k_r}$ (with all the λ 's different) we see from Proposition 13.9 that

$$\mathbb{C}[x]/f(x) \simeq \mathbb{C}[x]/(x - \lambda_1)^{k_1} \oplus \dots \oplus \mathbb{C}[x]/(x - \lambda_r)^{k_r} = B(\lambda_1, k_1) \oplus \dots \oplus B(\lambda_r, k_r).$$

We can use this to give an explicit expression for M_A as a direct sum of $B(\lambda, k)$'s and thus to determine the JNF of A .

A different, and usually easier, approach is to use Corollary 13.16, which tells us that the number of copies of $B(\lambda, k)$ in the decomposition of M_A is just $g_{x-\lambda}^k(M_A)$, as defined in Definition 13.12. To calculate these numbers $g_{x-\lambda}^k(M_A)$, we need to recall the definition of the characteristic polynomial and minimal polynomial of a matrix.

Definition 14.5. [defn-char-poly]

The *characteristic polynomial* $\text{char}(A)$ of a square matrix A is the polynomial $\det(tI - A)$.

Note that this is easy to calculate directly from A ; for example, if

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

then the characteristic polynomial is

$$\begin{aligned} \begin{vmatrix} t-1 & -2 & -3 \\ -4 & t-5 & -6 \\ -7 & -8 & t-9 \end{vmatrix} &= (t-1) \begin{vmatrix} t-5 & -6 \\ -8 & t-9 \end{vmatrix} - (-2) \begin{vmatrix} -4 & -6 \\ -7 & t-9 \end{vmatrix} + (-3) \begin{vmatrix} -4 & t-5 \\ -7 & -8 \end{vmatrix} \\ &= (t-1)((t-5)(t-9) - 48) + 2(-4(t-9) - 42) - 3(32 + 7(t-5)) \\ &= t^3 - 15t^2 - 18t. \end{aligned}$$

It is also not hard to see that for any $n \times n$ matrix A we have $\text{char}(A) = t^n +$ lower terms ; in other words, $\text{char}(A)$ is a monic polynomial of degree n .

We now define the minimal polynomial of a square matrix A . First, we put $I = \{f \in \mathbb{C}[x] \mid f(A) = 0\}$. Clearly if $f(A) = g(A) = 0$ and h is arbitrary then $(f+g)(A) = f(A)+g(A) = 0$ and $(hf)(A) = h(A)f(A) = 0$, so I is an ideal. By Theorem 9.7, we see that $I = \mathbb{C}[x]g$ for some polynomial g . I next claim that I is never the zero ideal. Indeed, the set of all $n \times n$ -matrices over \mathbb{C} is a vector space over \mathbb{C} of dimension n^2 , so any list of $n^2 + 1$ such matrices must be linearly dependent. In particular, the list I, A, \dots, A^{n^2} is linearly dependent, so there is some list a_0, \dots, a_{n^2} (not all zero) such that $a_0I + a_1A + \dots + a_{n^2}A^{n^2} = 0$. If we put $f(x) = a_0 + a_1x + \dots + a_{n^2}x^{n^2}$ we find that $f \neq 0$ and $f \in I$, so $I \neq 0$. As $I = \mathbb{C}[x]g$, we deduce that $g \neq 0$. After multiplying g by a nonzero constant, we may assume that g is a monic polynomial. This justifies the following definition:

Definition 14.6. [defn-min-poly]

The *minimal polynomial* $\text{min}(A)$ of a square matrix A is the unique monic polynomial that generates the ideal $I_A = \{f \in \mathbb{C}[x] \mid f(A) = 0\}$.

Theorem 14.7. [thm-char-min]

Let A be an $n \times n$ matrix, with $\text{char}(A) = \prod_{i=1}^r (t - \lambda_i)^{r_i}$, where all the λ_i 's are distinct and $r_i > 0$. Then

- We have $\text{min}(A) = \prod_{i=1}^r (t - \lambda_i)^{s_i}$, where $0 < s_i \leq r_i$. In other words, the roots of $\text{min}(A)$ are precisely the same as the roots of $\text{char}(A)$, and the multiplicities of the roots in $\text{min}(A)$ are at most as large as the multiplicities in $\text{char}(A)$.
- The JNF of A contains only blocks of the form $J(\lambda_i, k)$, where λ_i is a root of the characteristic polynomial, as before. The number of such blocks is $\dim(\ker(A - \lambda_i I)) = n - \text{rank}(A - \lambda_i I)$. The maximum value of k that occurs with λ_i is precisely s_i , and the sum of all the k 's that occur with λ_i is r_i .

Remark 14.8. [rem-cayley-hamilton]

In particular, part (a) says that $\text{min}(A)(t)$ divides $\text{char}(A)(t)$, so $\text{char}(A)(t) \in I_A$, so if we substitute the matrix A into its characteristic polynomial (in other words, evaluate $\text{char}(A)(A)$) we get the zero matrix. This is the *Cayley-Hamilton Theorem*.

The proof will follow after some examples and preliminary results.

Example 14.9. [eg-JNF-i]

Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 2 & 3 \\ 0 & 0 & 2 \end{bmatrix}.$$

The characteristic polynomial is

$$\text{char}(A) = \det \begin{bmatrix} t-2 & -3 & -4 \\ 0 & t-2 & -3 \\ 0 & 0 & t-2 \end{bmatrix} = (t-2)^3.$$

Here we have used the fact that if a matrix has zeros everywhere below the diagonal, then the determinant is just the product of the entries on the diagonal. It follows from the theorem that the minimal polynomial must be $(t-2)$ or $(t-2)^2$ or $(t-2)^3$. However, we find that

$$A - 2I = \begin{bmatrix} 0 & 3 & 4 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix} \quad (A - 2I)^2 = \begin{bmatrix} 0 & 0 & 9 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

If the minimal polynomial is $f(t)$, we must have $f(A) = 0$. Thus, the above shows that $\min(A) \neq t-2$ and $\min(A) \neq (t-2)^2$, so $\min(A)$ must be $(t-2)^3$.

We also see that the JNF of A can only contain blocks of the form $J(2, k)$, so it must have the form $J(2, k_1) \oplus \dots \oplus J(2, k_r)$, and we may as well order the factors so that $0 < k_1 \leq \dots \leq k_r$. By comparing characteristic polynomials we see that $(t-2)^3 = \text{char}(A) = (t-2)^{k_1 + \dots + k_r}$. By comparing minimal polynomials (and noting that $\max(k_1, \dots, k_r) = k_r$) we see that $(t-2)^3 = \min(A) = (t-2)^{k_r}$. Thus $3 = k_1 + \dots + k_r = k_r$. As all the k_i 's are supposed to be positive, this can only work if $r = 1$ and $k_1 = 3$. Thus the JNF of A is just the single Jordan block $J(2, 3)$, and thus $M_A \simeq B(2, 3) = \mathbb{C}[x]/(x-2)^3$.

For an alternative approach, we can observe from our formula for $A - 2I$ that $\text{rank}(A - 2I) = 2$ and so $\dim(\ker(A - 2I)) = 3 - 2 = 1$. Part (b) of Theorem 14.7 therefore tells us that there is only one block in the JNF, so A is conjugate to $J(2, k)$ for some k . As A is a 3×3 matrix, we must have $k = 3$.

Example 14.10. [eg-JNF-ii]

Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} 2 & i & i & 2 \\ i & 0 & 0 & i \\ i & 0 & 0 & i \\ 2 & i & i & 2 \end{bmatrix}.$$

The characteristic polynomial is the determinant of the first matrix shown below:

$$\begin{bmatrix} t-2 & -i & -i & -2 \\ -i & t & 0 & -i \\ -i & 0 & t & -i \\ -2 & -i & -i & t-2 \end{bmatrix} \rightarrow \begin{bmatrix} t & 0 & 0 & -t \\ 0 & t & -t & 0 \\ -i & 0 & t & -i \\ -2 & -i & -i & t-2 \end{bmatrix} \rightarrow \begin{bmatrix} t & 0 & 0 & 0 \\ 0 & t & 0 & 0 \\ -i & 0 & t & -2i \\ -2 & -i & -2i & t-4 \end{bmatrix}.$$

It is perfectly possible to evaluate the determinant directly, but it is more efficient to perform some row and column operations first, as shown above. In the first step we have subtracted the fourth row from the first row and the third row from the second row. In the second step we have added the first column to the fourth column and the second column to the third column. None of these operations change the determinant. The determinant of our final matrix is t^2 times the determinant of the 2×2 block in the bottom right corner, which is $t^2 - 4t + 4$. Thus $\det(tI - A) = t^2(t^2 - 4t + 4) = t^2(t-2)^2$. The factor $(t-2)^2$ comes from some Jordan blocks of the form $J(2, k)$, each of which contributes a factor $(t-2)^k$. The only possibilities are $J(2, 2)$ and $J(2, 1) \oplus J(2, 1)$. Similarly, the factor t^2 comes from $J(0, 2)$ or $J(0, 1) \oplus J(0, 1)$. This gives four possibilities for the JNF of A ; we list these below, together with their minimal polynomials.

$$\begin{array}{ll} J(0, 1) \oplus J(0, 1) \oplus J(2, 1) \oplus J(2, 1) & t(t-2) \\ J(0, 2) \oplus J(2, 1) \oplus J(2, 1) & t^2(t-2) \\ J(0, 1) \oplus J(0, 1) \oplus J(2, 2) & t(t-2)^2 \\ J(0, 2) \oplus J(2, 2) & t^2(t-2)^2. \end{array}$$

The minimal polynomial of A must be one of the polynomials in the list above. By direct calculation, we find that

$$A(A - 2I) = 2 \begin{bmatrix} 1 & i & i & 1 \\ i & -1 & -1 & i \\ i & -1 & -1 & i \\ 1 & i & i & 1 \end{bmatrix}$$

$$A^2(A - 2I) = 4 \begin{bmatrix} 1 & i & i & 1 \\ i & -1 & -1 & i \\ i & -1 & -1 & i \\ 1 & i & i & 1 \end{bmatrix}$$

$$A(A - 2I)^2 = A^2(A - 2I)^2 = 0.$$

It follows that the minimal polynomial of A is $t(t - 2)^2$. By comparing this with the list, we deduce that the JNF of A must be $J(0, 1) \oplus J(0, 1) \oplus J(2, 2)$.

Example 14.11. [eg-need-rank]

Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} 0 & 0 & 1 & 1 \\ -1 & 1 & 1 & 0 \\ -1 & 0 & 2 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Here we will just calculate the characteristic polynomial directly by expanding along the top row, although more efficient methods are certainly possible. We have

$$\begin{vmatrix} t & 0 & -1 & -1 \\ 1 & t-1 & -1 & 0 \\ 1 & 0 & t-2 & -1 \\ 0 & 0 & 0 & t-1 \end{vmatrix} = t \begin{vmatrix} t-1 & -1 & 0 \\ 0 & t-2 & -1 \\ 0 & 0 & t-1 \end{vmatrix} - 0 \begin{vmatrix} 1 & -1 & 0 \\ 1 & t-2 & -1 \\ 0 & 0 & t-1 \end{vmatrix} +$$

$$(-1) \begin{vmatrix} 1 & t-1 & 0 \\ 1 & 0 & -1 \\ 0 & 0 & t-1 \end{vmatrix} - (-1) \begin{vmatrix} 1 & t-1 & -1 \\ 1 & 0 & t-2 \\ 0 & 0 & 0 \end{vmatrix}$$

$$\begin{vmatrix} t-1 & -1 & 0 \\ 0 & t-2 & -1 \\ 0 & 0 & t-1 \end{vmatrix} = (t-1)^2(t-2)$$

$$\begin{vmatrix} 1 & t-1 & 0 \\ 1 & 0 & -1 \\ 0 & 0 & t-1 \end{vmatrix} = -(t-1)^2$$

$$\begin{vmatrix} 1 & t-1 & -1 \\ 1 & 0 & t-2 \\ 0 & 0 & 0 \end{vmatrix} = 0$$

so

$$\begin{aligned} \text{char}(A) &= t(t-1)^2(t-2) - 0 + (-1)(-(t-1)^2) - 0 \\ &= (t-1)^2(t(t-2) + 1) \\ &= (t-1)^2(t^2 - 2t + 1) = (t-1)^4. \end{aligned}$$

It follows that the minimal polynomial is $(t - 1)^k$ for some k with $1 \leq k \leq 4$. We have

$$(A - I) = \begin{bmatrix} -1 & 0 & 1 & 1 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$(A - I)^2 = 0.$$

It follows that the minimal polynomial must be $(t-1)^2$. We also see that $\text{rank}(A-I) = 2$, so $\dim(\ker(A-I)) = 4 - 2 = 2$, so there are 2 blocks in the JNF. The JNF is thus $J(1, k_1) \oplus J(1, k_2)$ for some k_1, k_2 with $k_1 \leq k_2$. By comparing characteristic polynomials we find that $k_1 + k_2 = 2$, and by comparing minimal polynomials we find that $k_2 = \max(k_1, k_2) = 2$. It follows that $k_1 = k_2 = 2$, so the JNF is $J(1, 2) \oplus J(1, 2)$.

Proposition 14.12. [prop-conj-char]

If A is conjugate to B , then $\text{char}(A) = \text{char}(B)$ and $\min(A) = \min(B)$ and $\dim(\ker(A - \lambda I)) = \dim(\ker(B - \lambda I))$ for all $\lambda \in \mathbb{C}$.

Proof. As A and B are conjugate, we have $A = PBP^{-1}$ for some invertible matrix P . It follows that $P(tI - B)P^{-1} = tPP^{-1} - PBP^{-1} = tI - A$ so

$$\det(tI - A) = \det(P(tI - B)P^{-1}) = \det(P) \det(tI - B) \det(P)^{-1} = \det(tI - B),$$

so $\text{char}(A) = \text{char}(B)$ as claimed.

Next, we have $f(A) = 0$ iff $f.m = 0$ for all $m \in M_A$, and $f(B) = 0$ iff $f.m = 0$ for all $m \in M_B$. As A is conjugate to B the modules M_A and M_B are isomorphic, so $f(A) = 0$ iff $f(B) = 0$. (More directly, one can check that $f(A) = Pf(B)P^{-1}$, and again it follows that $f(A) = 0$ iff $f(B) = 0$.) Thus, the ideals I_A and I_B are the same, so they have the same monic generator, in other words $\min(A) = \min(B)$.

Finally, note that $\ker(A - \lambda I) = \{m \in M_A \mid (x - \lambda)m = 0\}$ and $\ker(B - \lambda I) = \{m \in M_B \mid (x - \lambda)m = 0\}$. As $M_A \simeq M_B$ we see that these two vector spaces are isomorphic and thus have the same dimension, as claimed. \square

Proposition 14.13. [prop-oplus-char]

Let A and B be square matrices of sizes n and m . Then $\text{char}(A \oplus B) = \text{char}(A) \text{char}(B)$, and $\min(A \oplus B)$ is the least common multiple of $\min(A)$ and $\min(B)$. Moreover, for any $\lambda \in \mathbb{C}$ we have

$$\dim \ker(A \oplus B - \lambda I_{n+m}) = \dim \ker(A - \lambda I_n) + \dim \ker(B - \lambda I_m).$$

Proof. We first claim that $\det(A \oplus B) = \det(A) \det(B)$. We have

$$A \oplus B = \left(\begin{array}{c|c} A & 0 \\ \hline 0 & B \end{array} \right) = \left(\begin{array}{c|c} A & 0 \\ \hline 0 & I_m \end{array} \right) \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & B \end{array} \right) = (A \oplus I_m)(I_n \oplus B),$$

so $\det(A \oplus B) = \det(A \oplus I_m) \det(I_n \oplus B)$. By expanding along the top row we find that $\det(I_n \oplus B) = \det(I_{n-1} \oplus B)$, and it follows inductively that $\det(I_n \oplus B) = \det(B)$ for all n . Similarly, by expanding along the bottom row we see that $\det(A \oplus I_m) = \det(A)$ for all m , so the equation $\det(A \oplus B) = \det(A \oplus I_m) \det(I_n \oplus B)$ gives $\det(A \oplus B) = \det(A) \det(B)$ as claimed.

Next, observe that $tI_{n+m} - A \oplus B = (tI_n - A) \oplus (tI_m - B)$, so $\det(tI_{n+m} - A \oplus B) = \det(tI_n - A) \det(tI_m - B) = \text{char}(A) \text{char}(B)$ as claimed.

Next, note that $f(A \oplus B) = f(A) \oplus f(B)$. Thus

$$\begin{aligned} f \text{ is divisible by } \min(A \oplus B) &\Leftrightarrow f(A \oplus B) = 0 \\ &\Leftrightarrow f(A) = 0 \text{ and } f(B) = 0 \\ &\Leftrightarrow f \text{ is divisible by both } \min(A) \text{ and } \min(B). \end{aligned}$$

This means that $\min(A \oplus B)$ is the least common multiple of $\min(A)$ and $\min(B)$.

Finally, we can regard \mathbb{C}^{n+m} as $\mathbb{C}^n \oplus \mathbb{C}^m$ and we then have $(A \oplus B).(u, v) = (Au, Bv)$. It follows that $(A \oplus B - \lambda I).(u, v) = (Au - \lambda u, Bv - \lambda v)$, and thus that $\ker(A \oplus B - \lambda I)$ is the set of pairs (u, v) for which $(A - \lambda I)u = 0$ and $(B - \lambda I)v = 0$. In other words we have $\ker(A \oplus B - \lambda I) = \ker(A - \lambda I) \oplus \ker(B - \lambda I)$ and so $\dim \ker(A \oplus B - \lambda I) = \dim \ker(A - \lambda I) + \dim \ker(B - \lambda I)$. \square

Proposition 14.14. [prop-jordan-char]

We have $\text{char}(J(\lambda, k)) = \min(J(\lambda, k)) = (t - \lambda)^k$. Moreover, we have

$$\dim \ker(J(\lambda, k) - \mu I) = \begin{cases} 1 & \text{if } \lambda = \mu \\ 0 & \text{if } \lambda \neq \mu. \end{cases}$$

Proof. I first claim that $\det(J(\lambda, k)) = \lambda^k$. To see this, we need to recall the usual row-expansion method for evaluating determinants. Suppose we have a $k \times k$ matrix A with entries a_1, \dots, a_k on the top row, and that A_i is obtained from A by deleting the top row and the i 'th column; then

$$\det(A) = a_1 \det(A_1) - a_2 \det(A_2) + \dots \pm a_k \det(A_k).$$

If we take $A = J(\lambda, k)$ then $a_1 = \lambda$, $A_1 = J(\lambda, k-1)$ and $a_2 = \dots = a_k = 0$, so $\det(J(\lambda, k)) = \lambda \det(J(\lambda, k-1))$. Moreover, $J(\lambda, 1)$ is the 1×1 matrix (λ) so $\det(J(\lambda, 1)) = \lambda$. It follows inductively that $\det(J(\lambda, k)) = \lambda^k$ for all $k > 0$, as claimed. Here we illustrate the case $k = 5$:

$$J(\lambda, 5) = \left(\begin{array}{c|cccc} \lambda & 0 & 0 & 0 & 0 \\ \hline 1 & \lambda & 0 & 0 & 0 \\ 0 & 1 & \lambda & 0 & 0 \\ 0 & 0 & 1 & \lambda & 0 \\ 0 & 0 & 0 & 1 & \lambda \end{array} \right) = \left(\begin{array}{c|c} \lambda & 0 \\ \hline * & J(\lambda, 4) \end{array} \right).$$

Next, note that $J(\lambda, k) - tI = J(\lambda - t, k)$, so $\det(J(\lambda, k) - tI) = (\lambda - t)^k$. For a $k \times k$ matrix we have $\det(-A) = (-1)^k \det(A)$, so $\det(tI - J(\lambda, k)) = (-1)^k (\lambda - t)^k = (t - \lambda)^k$ as claimed.

We next need to understand which polynomials $f(x)$ have $f(J(\lambda, k)) = 0$. This happens iff $f.m = 0$ for all m in the module $M_{J(\lambda, k)}$, which is isomorphic to $B(\lambda, k) = \mathbb{C}[x]/(x - \lambda)^k$. It is clear that $f.B(\lambda, k) = \{0\}$ iff f is divisible by $(x - \lambda)^k$, so $\min(J(\lambda, k)) = (x - \lambda)^k$ as well.

Finally, note that $J(\lambda, k) - \mu I = J(\lambda - \mu, k)$. If $\mu \neq \lambda$ we deduce that $\det(J(\lambda, k) - \mu I) = (\lambda - \mu)^k \neq 0$, so $J(\lambda, k) - \mu I$ is invertible and $\ker(J(\lambda, k) - \mu I) = \{0\}$. On the other hand, if $\mu = \lambda$ then $J(\lambda, k) - \mu I = J(0, k)$, and it is not hard to see that $J(0, k).(x_1, \dots, x_k) = (0, x_1, \dots, x_{k-1})$. Thus $J(0, k).\underline{x} = 0$ iff $x_1 = \dots = x_{k-1} = 0$, so $\underline{x} = (0, \dots, 0, x_k)$ for some $x_k \in \mathbb{C}$. It follows that $\ker(J(\lambda, k) - \lambda I)$ is the span of the standard basis vector e_k , and so $\dim \ker(J(\lambda, k) - \lambda I) = 1$. \square

Proof of Theorem 14.7. Let A' be the JNF of A , so A' is conjugate to A and is a block sum of matrices of the form $J(\lambda, k)$ for various λ 's and k 's. We know from Proposition 14.12 that $\text{char}(A) = \text{char}(A')$ and $\min(A) = \min(A')$ and $\dim \ker(A - \lambda I) = \dim \ker(A' - \lambda I)$ for all λ .

First suppose that all the Jordan blocks in A' have the same eigenvalue λ . Then we can write

$$A' = J(\lambda, k_1) \oplus \dots \oplus J(\lambda, k_d)$$

for some sequence k_1, \dots, k_d of positive integers. We then have

$$\text{char}(A') = \prod_i \text{char}(J(\lambda, k_i)) = \prod_i (t - \lambda)^{k_i} = (t - \lambda)^k,$$

where $r = \sum_i k_i$. We also see that $\min(A')$ is the least common multiple of the polynomials $(t - \lambda)^{k_i}$, which is $(t - \lambda)^s$, where $s = \max(k_1, \dots, k_d)$. As each k_i is positive this means that $0 < s \leq r$. Finally, we see that

$$\begin{aligned} \dim \ker(A - \lambda I) &= \dim \ker(A' - \lambda I) \\ &= \sum_{i=1}^d \dim \ker(J(\lambda, k_i) - \lambda I) \\ &= \sum_{i=1}^d 1 = d. \end{aligned}$$

A similar argument shows that $\dim \ker(A - \mu I) = 0$ for $\mu \neq \lambda$.

More generally, there will be a number of different eigenvalues, say $\lambda_1, \dots, \lambda_d$. Let A'_i be the block sum of all the terms of the form $J(\lambda_i, k)$ for some k . The above argument shows that $\text{char}(A'_i) = (t - \lambda_i)^{r_i}$ and $\min(A'_i) = (t - \lambda_i)^{s_i}$ for some integers r_i, s_i with $0 < s_i \leq r_i$. We also have $A' = A'_1 \oplus \dots \oplus A'_d$, so

$$\text{char}(A) = \text{char}(A') = \prod_i (t - \lambda_i)^{r_i}.$$

Similarly, we find that $\min(A)$ is the least common multiple of the polynomials $(t - \lambda_1)^{s_1}, \dots, (t - \lambda_d)^{s_d}$. As these polynomials are all powers of inequivalent irreducibles, their lcm is just their product, so

$$\min(A) = \prod_i (t - \lambda_i)^{s_i}.$$

We also see that $\dim \ker(A - \lambda_i) = \sum_j \dim \ker(A_j - \lambda_i)$. The terms for $j \neq i$ are zero, and the term for $j = i$ is just the number of Jordan blocks in A_i .

The theorem now follows immediately. \square

We conclude by studying the structure of cyclic modules over $\mathbb{C}[x]$.

Let $f(x)$ be a monic polynomial of degree n over \mathbb{C} . We then have a cyclic module $\mathbb{C}[x]/f(x)$ over $\mathbb{C}[x]$. For any polynomial $g(x) \in \mathbb{C}[x]$ we have an element $\overline{g(x)} = g(x) + \mathbb{C}[x]f(x) \in \mathbb{C}[x]/f(x)$, and $\overline{g(x)} = \overline{h(x)}$ iff $g(x) - h(x)$ is divisible by $f(x)$.

Proposition 14.15. [prop-cyclic-basis]

If f is as above then the elements $\overline{1}, \overline{x}, \dots, \overline{x^{n-1}}$ form a basis for $\mathbb{C}[x]/f(x)$ over \mathbb{C} . In particular, we have $\dim_{\mathbb{C}}(\mathbb{C}[x]/f(x)) = n$.

Proof. Every element of $\mathbb{C}[x]/f(x)$ can be written as $\overline{g(x)}$ for some $g(x) \in \mathbb{C}[x]$. We can divide g by f to get $g(x) = f(x)q(x) + r(x)$ for some polynomial $r(x)$ of degree at most $n - 1$. It follows that $g(x) - r(x)$ is divisible by $f(x)$, so $\overline{g(x)} = \overline{r(x)}$. As $\deg(r) < n$ we have $r(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1}$ for some $a_0, \dots, a_{n-1} \in \mathbb{C}$. It follows that $r(x) = \sum_{i=0}^{n-1} a_i \overline{x^i}$, so the elements $\overline{1}, \overline{x}, \dots, \overline{x^{n-1}}$ span $\mathbb{C}[x]/f(x)$ over \mathbb{C} .

Now suppose we have a linear relation among these elements, say $b_0\overline{1} + \dots + b_{n-1}\overline{x^{n-1}} = \overline{0}$. This means that the polynomial $g(x) := b_0 + b_1x + \dots + b_{n-1}x^{n-1}$ satisfies $\overline{g(x)} = \overline{0}$, so g is divisible by f . As the degree of g is less than the degree of f , this can only happen if $g = 0$, which means that $b_0 = \dots = b_{n-1} = 0$. Thus, the elements $\overline{1}, \dots, \overline{x^{n-1}}$ are linearly independent over \mathbb{C} . \square

Theorem 14.16. [thm-JNF-cyclic]

Let A be an $n \times n$ matrix over \mathbb{C} , with JNF $J(\lambda_1, k_1) \oplus \dots \oplus J(\lambda_r, k_r)$. Then the following statements are all equivalent (so if any one of them is true, then all of them are true):

- (a) M_A is a cyclic module over $\mathbb{C}[x]$.
- (b) The numbers λ_i are all different.
- (c) $\min(A) = \text{char}(A)$.
- (d) $M_A \simeq \mathbb{C}[x]/\text{char}(A)(x)$.
- (e) There is a vector $v \in M_A$ such that $\{v, Av, \dots, A^{n-1}v\}$ is a basis for M_A over \mathbb{C} .

Proof. (a) \Rightarrow (b): If M_A is cyclic then $M_A \simeq \mathbb{C}[x]/f(x)$ for some polynomial $f(x)$. This must be nonzero, otherwise M_A would be the same as $\mathbb{C}[x]$ and thus would have infinite dimension over \mathbb{C} , which is impossible because $\dim_{\mathbb{C}}(M_A) = n$. We can factor $f(x)$ as $c(x - \mu_1)^{k_1} \dots (x - \mu_s)^{k_s}$ for some nonzero constant c and numbers μ_1, \dots, μ_s . By collecting terms we may assume that the μ_i 's are all different. Using the Chinese Remainder Theorem we find that

$$M_A \simeq \mathbb{C}[x]/f(x) = \bigoplus_i \mathbb{C}[x]/(x - \mu_i)^{k_i} = \bigoplus_i B(\mu_i, k_i),$$

so the JNF of A is $J(\mu_1, k_1) \oplus \dots \oplus J(\mu_s, k_s)$. Thus the μ 's are the same as the λ 's (up to possible reordering) and so the λ_i 's are all different.

(b) \Rightarrow (a): Conversely, suppose that the λ_i 's are all different. Then the polynomials $(x - \lambda_i)^{k_i}$ are all coprime to each other, so if we put $f(x) = \prod_i (x - \lambda_i)^{k_i}$ the Chinese Remainder Theorem tells us that

$$\mathbb{C}[x]/f(x) \simeq \bigoplus_i \mathbb{C}[x]/(x - \lambda_i)^{k_i} = \bigoplus_i B(\lambda_i, k_i) \simeq M_A,$$

so M_A is cyclic.

(b) \Leftrightarrow (c): Note that A is conjugate to $J(\lambda_1, k_1) \oplus \dots \oplus J(\lambda_r, k_r)$. We know from Propositions 14.12, 14.13 and 14.14 that $\text{char}(A)$ is the product of the polynomials $(x - \lambda_i)^{k_i}$, and that $\min(A)$ is their least common multiple. The product is the same as the least common multiple iff the factors $(x - \lambda_i)^{k_i}$ are all coprime to each other, which is true iff the numbers λ_i are all different.

(a) \Leftrightarrow (d): If (d) holds (ie $M_A \simeq \mathbb{C}[x]/\text{char}(A)(x)$) then M_A has the form $\mathbb{C}[x]/f(x)$ and is certainly cyclic.

Conversely, suppose that (a) holds, so $M_A \simeq \mathbb{C}[x]/f(x)$ for some f . Just as in the proof that (a) \Rightarrow (b) we see that f is a unit multiple of $\prod_i (x - \lambda_i)^{k_i}$, which we know is the same as $\text{char}(A)(x)$. It follows that $M_A \simeq \mathbb{C}[x]/\text{char}(A)(x)$, so (d) holds.

(e) \Rightarrow (a): Suppose that $v \in M_A$ and that $\{v, Av, \dots, A^{n-1}v\}$ is a basis for M_A over \mathbb{C} . Then for any $w \in M_A$ there exist $a_0, \dots, a_{n-1} \in \mathbb{C}$ such that $w = a_0v + a_1Av + \dots + a_{n-1}A^{n-1}v$. Thus, if we put $g(x) = \sum_i a_i x^i \in \mathbb{C}[x]$ we find that $w = g.v$. It follows that v generates M_A as a $\mathbb{C}[x]$ -module, so M_A is cyclic.

(a) \Rightarrow (e): Suppose that M_A is cyclic, so we can choose an isomorphism $\alpha: \mathbb{C}[x]/f(x) \rightarrow M_A$. It follows that $\deg(f) = \dim(\mathbb{C}[x]/f(x)) = \dim(M_A) = n$. It follows from Proposition 14.15 that $\{\bar{1}, \dots, \bar{x}^{n-1}\}$ is a basis for $\mathbb{C}[x]/f(x)$, and thus that $\{\alpha(\bar{1}), \dots, \alpha(\bar{x}^{n-1})\}$ is a basis for M_A . If we put $v = \alpha(\bar{1})$ we find that $\alpha(\bar{x}^k) = \alpha(x^k \bar{1}) = x^k \cdot \alpha(\bar{1}) = A^k v$. This means that our basis is just $\{v, Av, \dots, A^{n-1}v\}$, as required. \square

Example 14.17. [eg-JNF-not-cyclic]

Consider the following matrices:

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

We find that $\text{char}(A) = \text{char}(B) = (t-1)^4$. Moreover $(A-I)^3 \neq 0$, so $\min(A) = (t-1)^4 = \text{char}(A)$, so M_A is cyclic. On the other hand, $(B-I)^2 = 0$ (and $B-I \neq 0$) so $\min(B) = (t-1)^2 \neq \text{char}(B)$, so M_B is not cyclic.

Example 14.18. [eg-JNF-cyclic]

Consider the following matrix:

$$A = \begin{bmatrix} 0 & 0 & 0 & -3 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 3 & 0 & 0 & 0 \end{bmatrix}.$$

We find that

$$\text{char}(A) = 9 + 10x^2 + x^4 = (x^2 + 1)(x^2 + 9) = (x+i)(x-i)(x+3i)(x-3i).$$

As the four roots are distinct, we see that the minimal polynomial is the same as the characteristic polynomial, so M_A is cyclic. If we put $v = (1, 1, 1, 1) \in \mathbb{C}^4$ we find that

$$\begin{aligned} v &= (1, 1, 1, 1) \\ Av &= (-3, -1, 1, 3) \\ A^2v &= (-9, -1, -1, -9) \\ A^3v &= (27, 1, -1, -27). \end{aligned}$$

By standard methods we can check that these vectors are linearly independent and thus form a basis of \mathbb{C}^4 . This gives another proof that M_A is cyclic.

Exercise 14.1. [ex-antisymmetric]

Let A be the matrix

$$\begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}.$$

We assume that $a, b, c \in \mathbb{R}$, but we consider A as a matrix over \mathbb{C} so we get a module M_A over $\mathbb{C}[x]$. Prove that

$$M_A \simeq M_0 \oplus M_{ir} \oplus M_{-ir},$$

where $r = \sqrt{a^2 + b^2 + c^2}$.

Exercise 14.2. [ex-circulant]

(This question is quite elaborate.) A 3×3 circulant matrix is a matrix of the following form:

$$A = \begin{bmatrix} u & v & w \\ v & w & u \\ w & u & v \end{bmatrix}.$$

We will assume that u , v and w are real numbers. Put $a = u+v+w$ and $b = ((u-v)^2 + (v-w)^2 + (w-u)^2)/2$.

- Show that the characteristic polynomial of A is $(x-a)(x^2-b)$ [You may wish to start by performing some row and column operations rather than just wading in and calculating the determinant.]
- Show that if $uv + vw + wu \neq 0$ and u, v, w are not all the same then the minimal polynomial is equal to the characteristic polynomial.
- Show that if $uv + vw + wu = 0$ and $a \neq 0$ then the minimal polynomial is $x^2 - a^2$.
- Calculate and factorise the minimal polynomial when $u = -13$, $v = 11$ and $w = 10$.
- Calculate and factorise the minimal polynomial when $u = -2$, $v = 3$ and $w = 6$.

Exercise 14.3. [ex-JNF-i]

Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} -1 & 1 & 1 & -1 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & -1 \end{bmatrix}.$$

- What is the characteristic polynomial of A ?
- What is the minimal polynomial of A ?
- What is the rank of the matrix $A + I$?
- Which direct sum of basic $\mathbb{C}[x]$ -modules is isomorphic to M_A ?

Exercise 14.4. [ex-JNF-ii]

Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 5 \\ 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

- What is the characteristic polynomial of A ?
- What is the minimal polynomial of A ?
- Which direct sum of basic $\mathbb{C}[x]$ -modules is isomorphic to M_A ?
- What is the Jordan normal form of A ?

Exercise 14.5. [ex-JNF-iii]

Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} -1 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 1 & 1 \end{bmatrix}.$$

- What is the characteristic polynomial of A ?
- What are the ranks of $A + I$ and $A - I$?
- What is the Jordan normal form of A ?
- Show that M_A is cyclic.

Exercise 14.6. [ex-JNF-iv]

Put $\alpha = \sqrt{-8}$. Consider the following matrix over \mathbb{C} :

$$A = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \alpha \\ 0 & 1 & \alpha & -4 \\ 1 & \alpha & -4 & -\alpha \end{bmatrix}.$$

- (a) What is the characteristic polynomial of A ?
- (b) What are the ranks of $A + I$ and $A - I$?
- (c) What is the Jordan normal form of A ?
- (d) Show that M_A is cyclic.

SOLUTIONS

Exercise 2.1:

- R_0 is a ring. The main point is to observe that R_0 is closed under addition and multiplication, because if $f(-x) = f(x)$ and $g(-x) = g(x)$ then

$$(f + g)(-x) = f(-x) + g(-x) = f(x) + g(x) = (f + g)(x)$$

$$fg(-x) = f(-x)g(-x) = f(x)g(x) = fg(x).$$

- R_1 is not a ring, because it is not closed under multiplication: if f and g lie in R_1 then

$$fg(-x) = f(-x)g(-x) = (-f(x))(-g(x)) = +fg(x) \neq -fg(x),$$

so $fg \notin R_1$ (except in trivial cases where $fg = 0$).

- R_2 is not a commutative ring, because matrix multiplication is not commutative in general. For example, if we take $a = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $b = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ then $ab = b$ and $ba = 0$ so $ab \neq ba$. All the other axioms are satisfied, however.
- R_3 is a ring. The additive identity is the zero matrix, and the multiplicative identity is the matrix $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.
- R_4 is not a ring. Firstly, it is not commutative, because $b \times a = -a \times b$. It is not even associative, because

$$a \times (b \times c) = (a.c)b - (a.b)c$$

$$(a \times b) \times c = (a.c)b - (b.c)a.$$

There is also no multiplicative identity: if there were, then we would have $1 \times 1 = 1$, but $a \times a$ is always zero for any vector a . (R_3 is in fact an example of a *Lie algebra*; these are rather different from rings, but also very important.)

Exercise 2.2:

- In $\mathbb{Z}_{(5)}$ we could take $a = 3/4$ and $b = 6/7$; these both lie in $\mathbb{Z}_{(5)}$ because 4 and 7 are not divisible by 5. We have $a + b = 45/28$ and $ab = 18/28 = 9/14$. These both lie in $\mathbb{Z}_{(5)}$ because 28 and 14 are not divisible by 5.
- In $\mathbb{Z}[i]$ we could take $a = 2 + 3i$ and $b = 4 - 5i$. We then have $a + b = 6 - 2i$ and $ab = 23 + 2i$; both of these clearly also lie in $\mathbb{Z}[i]$.
- In $\mathbb{Q}[x, y]$ we could take $a = (x + y)/2$ and $b = (x - y)/2$. Then $a + b = x$ and $ab = (x^2 - y^2)/4$, so $a + b$ and ab are again elements of $\mathbb{Q}[x, y]$.
- In \mathbb{Z}_{12} we could take $a = \bar{3}$ and $b = \bar{4}$, so $a + b = \bar{7} = \overline{-5}$ and $ab = \overline{12} = \bar{0}$.

Exercise 2.3: Put $a = 1/2$ and $b = -1/3$. Then $a, b \in R$ but $a + b = 1/6 \notin R$ and $ab = -1/6 \notin R$, so R is not closed under addition or multiplication.

Exercise 2.4: Let a and b be nonzero elements of K ; we must prove that $ab \neq 0$. As K is a field, we know that a and b are invertible, so we can find elements $c, d \in K$ with $ac = 1$ and $bd = 1$. It follows that $abcd = 1$. If ab were zero we would also have $abcd = 0$, so 1 would be equal to 0, contradicting the definition of a field. We must thus have $ab \neq 0$ as required.

Exercise 2.5:

- We have $A \cup \emptyset = A$ and $A \cap \emptyset = \emptyset$ so $A + \emptyset = A \setminus \emptyset = A$. Similarly, we have $A \cup A = A = A \cap A$, so $A + A = A \setminus A = \emptyset$.
- For the equation $\chi_{A+B}(x) = \chi_A(x) + \chi_B(x)$, there are four cases to consider.
 - x lies in both A and B , so x does not lie in $A+B$, so $\chi_{A+B}(x) = \bar{0}$. Here we have $\chi_A(x) + \chi_B(x) = \bar{1} + \bar{1} = \bar{2} = \bar{0}$ (because we are working in \mathbb{Z}_2), so $\chi_{A+B}(x) = \chi_A(x) + \chi_B(x)$ as required.

- (ii) x lies in A but not in B , so $x \in A+B$, so $\chi_{A+B}(x) = \bar{1}$. Here we have $\chi_A(x) + \chi_B(x) = \bar{1} + \bar{0} = \bar{1} = \chi_{A+B}(x)$ as required.
- (iii) x lies in B but not in A ; this works the same way as in (ii).
- (iv) x lies in neither A nor B . Here it is clear that $\chi_{A+B}(x) = \bar{0} = \bar{0} + \bar{0} = \chi_A(x) + \chi_B(x)$. The argument is similar but easier for the equation $\chi_{AB}(x) = \chi_A(x)\chi_B(x)$.
- (c) It is clear that $\{x \in X \mid \chi_A(x) = \bar{1}\} = A$. If $\chi_A = \chi_B$ then $\{x \mid \chi_A(x) = \bar{1}\} = \{x \mid \chi_B(x) = \bar{1}\}$, so $A = B$.
- (d) It is clear that the above rules do indeed define subsets of X , so R is closed under addition and multiplication. It is easy to see that $AB = A \cap B = B \cap A = BA$ and $(AB)C = (A \cap B) \cap C = A \cap (B \cap C) = A(BC)$, so multiplication is commutative and associative. Moreover, for $A \subseteq X$ we have $X \cap A = A$, so X is a multiplicative identity element.

It is also clear that $A+B = B+A$, so addition is commutative. Part (a) says that \emptyset is an additive identity, and A is an additive inverse for itself. We next show that addition is associative. By part (b), we have

$$\chi_{A+(B+C)}(x) = \chi_A(x) + \chi_{B+C}(x) = \chi_A(x) + \chi_B(x) + \chi_C(x) = \chi_{A+B}(x) + \chi_C(x) = \chi_{(A+B)+C}(x).$$

It follows using (c) that $A + (B + C) = (A + B) + C$, as required.

All that is left is to check distributivity, which can be done by the same method. We have

$$\begin{aligned} \chi_{A(B+C)}(x) &= \chi_A(x)\chi_{B+C}(x) \\ &= \chi_A(x)(\chi_B(x) + \chi_C(x)) \\ &= \chi_A(x)\chi_B(x) + \chi_A(x)\chi_C(x) \\ &= \chi_{AB}(x) + \chi_{AC}(x) \\ &= \chi_{AB+AC}(x), \end{aligned}$$

so $A(B+C) = AB+AC$ as required.

Exercise 3.1: The elements are $(\bar{0}, \bar{0}), (\bar{0}, \bar{1}), (\bar{0}, \bar{2}), (\bar{0}, \bar{3}), (\bar{0}, \bar{4}), (\bar{1}, \bar{0}), (\bar{1}, \bar{1}), (\bar{1}, \bar{2}), (\bar{1}, \bar{3})$ and $(\bar{1}, \bar{4})$. I claim that the element $x := (\bar{1}, \bar{1})$ has order 10. Indeed, we have $nx = (\bar{n}, \bar{n})$. The first \bar{n} is in \mathbb{Z}_2 , so it is zero iff n is divisible by 2. The second \bar{n} is in \mathbb{Z}_5 , so it is zero iff n is divisible by 5. Thus $nx = (\bar{0}, \bar{0})$ iff n is divisible by both 2 and 5, or equivalently iff n is divisible by 10. This means that x has order 10 as claimed.

Exercise 3.2:

- (a) We have $D.t^3 = 3t^2$ so $D^2.t^3 = 3D.t^2 = 6t$ so $D^3.t^3 = 6D.t = 6$. This means that $(D^2/2).t^3 = 3t$ and $(D^3/6).t^3 = 1$ so $(1 + D + D^2/2 + D^3/6).t^3 = t^3 + 3t^2 + 3t + 1$. We notice that this is just $(t+1)^3$. The generalisation is that

$$\left(\sum_{k=0}^m D^k/k!\right).t^m = (t+1)^m.$$

More generally, if $f(t)$ is any polynomial of degree less than or equal to m , it can be shown that

$$\left(\sum_{k=0}^m D^k/k!\right).f(t) = f(t+1).$$

This is essentially Taylor's theorem.

- (b) Put

$$g(t) = ((D+1)f)(t) = f'(t) + f(t) = (-e^{-t} \sin(t) + e^{-t} \cos(t)) + e^{-t} \sin(t) = e^{-t} \cos(t).$$

Then $((D+1)^2 f)(t) = ((D+1)g)(t) = g'(t) + g(t) = -e^{-t} \sin(t)$, by a similar calculation. In other words $(D+1)^2 f = -f$.

- (c) We have $((D-1)g_k)(t) = g'_k(t) - g_k(t) = (kt^{k-1}e^t + t^k e^t) - t^k e^t = kt^{k-1}e^t$, or in other words $(D-1)g_k = kg_{k-1}$. It follows that

$$\begin{aligned}(D-1)^2 g_k &= k(D-1)g_{k-1} = k(k-1)g_{k-2} \\ (D-1)^3 g_k &= k(k-1)(D-1)g_{k-2} = k(k-1)(k-2)g_{k-3}\end{aligned}$$

and so on. We eventually find that $(D-1)^k g_k = k!g_0$, so $((D-1)^k g_k)(t) = k!e^t$.

- (d) We certainly have $(D^0 f)(t) = f(t) = (0+t)e^t$. Assuming that $(D^k f)(t) = (k+t)e^t$ for some particular value of k , we have

$$(D^{k+1} f)(t) = D((k+t)e^t) = (k+t)D(e^t) + e^t D(k+t) = (k+t)e^t + e^t = ((k+1)+t)e^t.$$

It follows by induction that $(D^k f)(t) = (k+t)e^t$ for all k . Thus, for an operator $p(D) = \sum_k a_k D^k$, we have

$$(p(D)f)(t) = \sum_k a_k (k+t)e^t = \left(\sum_k k a_k\right) e^t + \left(\sum_k a_k\right) t e^t.$$

We also have $p(1) = \sum_k a_k \cdot 1^k = \sum_k a_k$. Similarly, we have $p'(D) = \sum_k k a_k D^{k-1}$, so $p'(1) = \sum_k k a_k$. Putting these into our earlier formula gives

$$(p(D)f)(t) = (p'(1) + p(1)t)e^t,$$

as claimed.

Exercise 3.3:

- (a) We just need to check that V is closed under differentiation. Note that $v'(t) = te^{t^2/2} = tv(t)$, so

$$\frac{d}{dt} f(t)v(t) = f(t)v'(t) + f'(t)v(t) = (tf(t) + f'(t))v(t).$$

If $f(t)$ is a polynomial, then clearly $tf(t) + f'(t)$ is also a polynomial, so the function $(tf(t) + f'(t))v(t)$ lies in V as required.

- (b) If we differentiate repeatedly using the above rule we find that

$$\begin{aligned}(D^0 v)(t) &= v(t) \\ (D^1 v)(t) &= tv(t) \\ (D^2 v)(t) &= (t^2 + 1)v(t) \\ (D^3 v)(t) &= (t^3 + 3t)v(t).\end{aligned}$$

- (c) As V is an $\mathbb{R}[D]$ -module, we must have $D^k v \in V$, so $D^k v = p_k v$ for some polynomial p_k . (From part (b) we see that $p_0(t) = 1$, $p_1(t) = t$, $p_2(t) = t^2 + 1$ and $p_3(t) = t^3 + 3t$.) Using part (a) we see that $p_{k+1}(t) = tp_k(t) + p'_k(t)$. The claim is that $p_k(t) = t^k + \text{lower terms}$. If this is true for some value of k , then $tp_k(t) = t^{k+1} + \text{lower terms}$ and $p'_k(t) = kt^{k-1} + \text{lower terms}$, so $p_{k+1}(t) = t^{k+1} + \text{lower terms}$, so the claim holds for the next value of k . Moreover, the claim visibly holds for $k=0$, so it holds for all k by induction.

- (d) Let k be the degree of q , so $q(D) = a_0 + a_1 D + \dots + a_k D^k$ for some $a_0, \dots, a_k \in \mathbb{R}$ with $a_k \neq 0$. Then $q(D)v = (a_0 p_0 + \dots + a_k p_k)v$, and using part (c) we see that $a_0 p_0(t) + \dots + a_k p_k(t) = a_k t^k + \text{lower terms}$, so in particular it is not zero.

- (e) First suppose that f has degree 0, say $f(t) = c$ for all t , where $c \in \mathbb{R}$. We can regard c as an element in $\mathbb{R}[D]$, and $fv = cv$ as required; this proves the claim for $k=0$.

Now suppose we have proved the claim for all polynomials of degree less than k , and that f has degree k . Then $f(t) = at^k + \text{lower terms}$ for some $a \in \mathbb{R}$. It follows that the function $g(t) = f(t) - ap_k(t)$ is a polynomial of degree less than k , so we have $gv = q(D)v$ for some $q(D) \in \mathbb{R}[D]$. It follows that

$$fv = gv + ap_k v = q(D)v + aD^k v = (q(D) + aD^k)v,$$

so $fv \in \mathbb{R}[D]v$ as required. The claim now follows for all degrees by induction.

- (f) We know from part (e) that v generates V as an $\mathbb{R}[D]$ -module. It follows that $V \simeq \mathbb{R}[D]/I$, where $I = \{q(D) \in \mathbb{R}[D] \mid q(D)v = 0\}$. Part (d) tells us that $I = 0$, so $V \simeq \mathbb{R}[D]$.

Exercise 4.1: The first few powers are

$$A^2 = \begin{bmatrix} 1 & 2 & 1 & 0 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 1 & 3 & 3 & 1 \\ 0 & 1 & 3 & 3 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A^4 = \begin{bmatrix} 1 & 4 & 6 & 4 \\ 0 & 1 & 4 & 6 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

In general, in the matrix A^n all the entries in the k 'th band above and parallel to the diagonal are equal to the binomial coefficient $\binom{n}{k}$. The entries below the diagonal are zero.

Exercise 4.2:

(a) $A^3 - I = \begin{bmatrix} 0 & 0 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ so

$$(x^3 - 1)m = \begin{bmatrix} 0 & 0 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix}.$$

- (b) $A^2 = I$ so $A^3 = A$ so $A^3 - I = A - I$. Moreover $Am = (3, 2, 1)$ so $(A - I)m = Am - m = (3, 2, 1) - (1, 2, 3) = (2, 0, -2)$. Thus $(x^3 - 1)m = (2, 0, -2)$.
- (c) $Am = (0, 0)$ so $x^k m = A^k m = 0$ for all $k > 0$. Thus when we expand out $(14x^{12} + 5x^{11} - 36x^7 - 22x^4 + 13x - 5)m$, all the terms except the last one are zero, so we are left with $-5m = (-5, 5)$.

Exercise 4.3: We have

$$A^2 = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{bmatrix}$$

and

$$(a + d)A = \begin{bmatrix} a^2 + ad & ab + bd \\ ac + cd & ad + d^2 \end{bmatrix}$$

so

$$\begin{aligned} f(A) &= A^2 - (a + d)A + (ad - bc)I \\ &= \begin{bmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{bmatrix} - \begin{bmatrix} a^2 + ad & ab + bd \\ ac + cd & ad + d^2 \end{bmatrix} + \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Exercise 4.4: $A^2 = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} = 2A$ so $A^4 = (A^2)^2 = (2A)^2 = 4A^2 = 8A$. Thus $f(A) = A^4 - 3A = 8A - 3A = 5A = \begin{bmatrix} 5 & 5 \\ 5 & 5 \end{bmatrix}$.

Exercise 4.5:

- (a) Clearly $A^0 = I$ and $A^1 = A$. We observe that $A^2 = I$, and it follows immediately that A^i is I whenever i is even, and A whenever i is odd.
- (b) We have $f(1) = \sum_i a_i = b + c$, and $f(-1) = \sum_i (-1)^i a_i = b - c$. It follows that $b = (f(1) + f(-1))/2$ and $c = (f(1) - f(-1))/2$.

- (c) We have $f(A) = \sum_i a_i A^i$. The term corresponding to an even number $i = 2j$ is $a_{2j}I$, whereas the term corresponding to an odd number $i = 2j + 1$ is $a_{2j+1}A$. We thus have

$$\begin{aligned} f(A) &= \sum_j (a_{2j}I + a_{2j+1}A) \\ &= bI + cA \\ &= ((f(1) + f(-1))/2)I + ((f(1) - f(-1))/2)A \\ &= f(1)(I + A)/2 + f(-1)(I - A)/2 \\ &= \frac{f(1)}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + \frac{f(-1)}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}. \end{aligned}$$

Exercise 5.1: Let $\alpha: M \rightarrow N$ be a homomorphism of modules over a Euclidean domain R . Suppose that $aM = \{0\}$ and $bN = \{0\}$ and let c be the gcd of a and b . I claim that $cn = 0$ for all $n \in \text{image}(\alpha)$. Indeed, we can write $c = au + bv$ for some $a, b \in R$. If $n \in \text{image}(\alpha)$ then $n = \alpha(m)$ for some $m \in M$. We have $am = 0$ (because $aM = \{0\}$) so $an = a\alpha(m) = \alpha(am) = \alpha(0) = 0$. We also have $n \in N$ and $bN = \{0\}$ so $bn = 0$. Thus $cn = uan + vbn = 0 + 0 = 0$ as claimed.

In the case considered we have $R = \mathbb{C}[x]$ and $a = x^3 - x = (x - 1)(x + 1)x$ and $b = x^5$ so it is clear that $c = x$. Thus $xn = 0$ for all $n \in \text{image}(\alpha)$.

Exercise 5.2: Suppose that $M \oplus N$ is cyclic, so there is an element $(x, y) \in M \oplus N$ as an R -module. This means that for any element $(m, n) \in M \oplus N$, there exists $a \in R$ such that $a(x, y) = (m, n)$. In particular, for any element $m \in M$ we have $(m, 0) \in M \oplus N$, so there exists $a \in R$ such that $a(x, y) = (m, 0)$, which means that $m = ax$. This shows that M is generated by the single element x , so M is cyclic. Similarly, N is generated by y and so is cyclic.

Exercise 5.3:

- (a) The set N_1 is not a submodule, because $(1, 0) \in N_1$ but $2(1, 0) = (2, 0) \notin N_1$. However, the set N_0 is a submodule. To see this, suppose that (n, m) and (n', m') lie in N_0 , so $n - m$ and $n' - m'$ are even. Then $(n, m) + (n', m') = (n + n', m + m')$ and the integer $(n + n') - (m + m') = (n - m) + (n' - m')$ is even so $(n, m) + (n', m') \in N_0$. Similarly, for any $a \in \mathbb{Z}$ we have $a(n, m) = (an, am)$ and $an - am = a(n - m)$ is even, so $a(n, m) \in N_0$. It is clear that $(0, 0) \in N_0$ and it follows that N_0 is a submodule as claimed.
- (b) I claim that both N_0 and N_1 are submodules of M_A . It is clear that they are both vector subspaces of \mathbb{R}^2 , so it is enough to check that they are both stable under A . An element $w \in N_0$ has the form $w = \begin{bmatrix} x \\ x \end{bmatrix}$ so $Aw = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ x \end{bmatrix} = \begin{bmatrix} 2x \\ 2x \end{bmatrix}$, so $Aw \in N_0$. This shows that N_0 is stable under A and thus is a submodule. Similarly, an element $w \in N_1$ has the form $w = \begin{bmatrix} x \\ -x \end{bmatrix}$ so $Aw = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$. As the zero vector certainly lies in N_1 we have $Aw \in N_1$ and so N_1 is also stable under A .
- (c) I claim that neither N_0 nor N_1 is an $\mathbb{R}[D]$ -submodule of $C^\infty(\mathbb{R}, \mathbb{R})$. Indeed, put $f(t) = t - 1$, so $f \in C^\infty(\mathbb{R}, \mathbb{R})$. Then $f(0) = 0$, so $f \in N_0$. However, $f'(0) = 1 \neq 0$, so $f' \notin N_0$, so N_0 is not closed under differentiation, so it is not an $\mathbb{R}[D]$ -submodule of $C^\infty(\mathbb{R}, \mathbb{R})$. To prove that N_1 is not a submodule, we can use the same function f . We have $\int_0^2 f = [t^2/2 - t]_0^2 = 0$, so $f \in N_1$. However, $\int_0^2 f' = \int_0^2 1 = 2$, so $f' \notin N_1$, so N_1 is not a submodule.

Exercise 5.4: If M were free, it would be isomorphic to R^d for some d , so we would have $20 = |M| = |R^d| = |R|^d = 10^d$. As 20 is not a power of 10, this is impossible, so M cannot be free.

Exercise 5.5: The elements of N_6 are the multiples of $\bar{6}$, which are $\bar{0}, \bar{6}, \bar{12}$ and $\bar{18}$. We can stop at this point because $\bar{24} = \bar{0}$, $\bar{30} = \bar{6}$ and so on. Thus $N_6 = \{\bar{0}, \bar{6}, \bar{12}, \bar{18}\}$, and similarly we have $N_4 = \{\bar{0}, \bar{4}, \bar{8}, \bar{12}, \bar{16}, \bar{20}\}$. From this we see that $N_4 \cap N_6 = \{\bar{0}, \bar{12}\} = N_{12}$, so we can take $d = 12$.

As $\bar{8} \in N_4$ and $\bar{18} \in N_6$, the group $N_4 + N_6$ contains $\bar{8} + \bar{18} = \bar{26} = \bar{2}$. As $N_4 + N_6$ is a subgroup of \mathbb{Z}_{24} we deduce that all multiples of $\bar{2}$ lie in $N_4 + N_6$, so $N_2 \subseteq N_4 + N_6$. On the other hand, as 4, 6 and 24 are all even we see that all elements of $N_4 + N_6$ have the form \bar{a} for some even integer a and thus they lie in N_2 . This shows that $N_4 + N_6 = N_2$, so we can take $e = 2$.

Exercise 5.6:

- (a) The order is $900/10 = 90$.
- (b) The factor group \mathbb{Z}_{900}/N_{10} is isomorphic to \mathbb{Z}_{10} .
- (c) Here d is the greatest common divisor of 70 and 900, which is 10.
- (d) Here d is the greatest common divisor of $12 = 2^2 \times 3$, $30 = 2 \times 3 \times 5$ and $100 = 2^2 \times 5^2$, so $d = 2$.
- (e) Here d is the least common multiple of $30 = 2 \times 3 \times 5$ and $50 = 2 \times 5^2$, so $d = 2 \times 3 \times 5^2 = 150$.

Exercise 5.7: Put $u = (1, 0, 0)$. Then $xu = (0, 0, 1)$ and $x^2u = (1, 1, 1)$. These three vectors are clearly linear independent, so they form a basis of \mathbb{Q}^3 , so they span \mathbb{Q}^3 . Thus any vector $v \in \mathbb{Q}^3$ can be written in the form $au + bxu + cx^2u$ for some $a, b, c \in \mathbb{Q}$. In other words, $v = (a + bx + cx^2)u \in \mathbb{Q}[x]u$, so we see that u generates M_A as a $\mathbb{Q}[x]$ -module. This means that $M_A \simeq \mathbb{Q}[x]/f(x)$ for some polynomial $f(x)$, which we can assume is monic. From the general theory we know that the degree of f is the size of A , which is 3. We also know that $f(x)$ is the only monic polynomial of degree 3 such that $f(x).u = 0$.

The polynomial f is in fact the characteristic polynomial of A , which can be calculated directly:

$$\det(xI - A) = \begin{vmatrix} x & 0 & -1 \\ 0 & x-1 & -1 \\ -1 & -1 & x-1 \end{vmatrix} = x^3 - 2x^2 - x + 1.$$

For another approach, consider the vector $x^3u = x(1, 1, 1) = (1, 2, 3)$. We would like to write this in the form $au + bxu + cx^2u$, so we want

$$(1, 2, 3) = a(1, 0, 0) + b(0, 0, 1) + c(1, 1, 1) = (a + c, c, b + c).$$

The solution is $a = -1, b = 1, c = 2$, so $x^3u = -u + xu + 2x^2u$, so $(x^3 - 2x^2 - x + 1)u = 0$, so $f(x) = x^3 - 2x^2 - x + 1$.

Exercise 5.8: Define $\alpha: \mathbb{R}[D] \rightarrow W_d$ by $\alpha(p(D)) = p(D).t^d$. Note that

$$\begin{aligned} D.t^d &= dt^{d-1} \\ D^2.t^d &= d(d-1)t^{d-2} \\ D^3.t^d &= d(d-1)(d-2)t^{d-3} \end{aligned}$$

and so on. In general, we have $D^k.t^d = m_k t^{d-k}$, where $m_k = d(d-1)\dots(d-k+1) = \prod_{i=0}^{k-1} (d-i)$, as one can easily check by induction. Note also that when $k \leq d$ all the factors $d-i$ for $0 \leq i \leq k-1$ are nonzero, so $m_k \neq 0$. However, we have $m_k = 0$ for $k > d$. Any element $f(t) \in W_d$ has the form $f(t) = a_0 + a_1t + \dots + a_d t^d$ for some $a_0, \dots, a_d \in \mathbb{R}$. If we define

$$p(D) = \sum_{i=0}^d a_{d-i} D^i / m_i = a_d + a_{d-1} D / m_1 + \dots + a_0 D^d / m_d$$

we find that

$$\begin{aligned} p(D).t^d &= a_d t^d + a_{d-1} m_1^{-1} D.t^d + \dots + a_0 m_d^{-1} D^d.t^d \\ &= a_d t^d + a_{d-1} t^{d-1} + \dots + a_0 t^0 \\ &= f(t). \end{aligned}$$

Thus every element $f \in W_d$ has the form $f = p(D).t^d$ for some $p(D) \in \mathbb{R}[D]$, so W_d is generated by t^d as a module over $\mathbb{R}[D]$.

Now put $I = \{p \in \mathbb{R}[D] \mid p(D).t^d = 0\}$, so $W_d \simeq \mathbb{R}[D]/I$. Suppose we have an element $p(D) = \sum_i b_i D^i \in \mathbb{R}[D]$. Then $p(D).t^d = b_0 t^d + b_1 m_1 t^{d-1} + \dots + b_d m_d t^0$, and this is zero iff $b_0 = \dots = b_d = 0$. This means that $p(D)$ has the form $b_{d+1} D^{d+1} + b_{d+2} D^{d+2} + \dots$, so it is divisible by D^{d+1} . Thus I is the principal ideal $\mathbb{R}[D].D^{d+1}$, and $W_d \simeq \mathbb{R}[D]/D^{d+1}$.

Exercise 5.9:

- (a) If $v \in L$ then $v = tu$ for some $t \in \mathbb{R}$, so $xv = \phi(v) = t(u \times u) = 0$ (using the fact that $a \times a = 0$ for any vector a). This shows that $xL = \phi(L) = 0$, so certainly $\phi(L) \leq L$, so L is a submodule.
- (b) For any $v \in M$ we have $xv = \phi(v) = u \times v$, which is always perpendicular to u , so it lies in K . This says that $xM = \phi(M) \leq K$, so certainly $\phi(K) \leq K$, so K is a submodule. Next, for any $v \in K$ we have $u.v = 0$ and so

$$x^2 v = \phi^2(v) = u \times (u \times v) = (u.v)u - (u.u)v = 0u - r^2 v = -r^2 v,$$

so $(x^2 + r^2)v = 0$. This shows that $(x^2 + r^2)K = 0$ as claimed.

- (c) We now know that $xM \leq K$ so

$$(x^3 + r^2 x)M = (x^2 + r^2)xM \leq (x^2 + r^2)K = 0.$$

Exercise 6.1: Such homomorphisms correspond to matrices P of the appropriate shape (the same number of columns as A , and the same number of rows as B) such that $PA = BP$.

- (a) Here P is a 2×2 matrix, say $P = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. We have $PA = \begin{bmatrix} a & a+b \\ c & c+d \end{bmatrix}$ and $BP = \begin{bmatrix} c & d \\ a & b \end{bmatrix}$. Thus $PA = BP$ if and only if $a = c$, $a + b = d$, $c = a$ and $c + d = b$. By solving these equations we find that $c = a = 0$ and $d = b$. Thus, the homomorphisms from M_A to M_B are precisely the matrices of the form $P = \begin{bmatrix} 0 & b \\ 0 & b \end{bmatrix}$ over \mathbb{Q} .
- (b) Here again we have $P = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for some a, b, c, d . Thus $PA = \begin{bmatrix} \lambda a & \mu b \\ \lambda c & \mu d \end{bmatrix}$ and $BP = \begin{bmatrix} \mu a & \mu b \\ \lambda c & \lambda d \end{bmatrix}$, so $PA = BP$ if and only if $\lambda a = \mu a$ and $\mu d = \lambda d$, or in other words $(\lambda - \mu)a = (\lambda - \mu)d = 0$. As $\lambda - \mu \neq 0$ this means that $a = d = 0$. Thus, the homomorphisms from M_A to M_B are precisely the matrices of the form $P = \begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}$.
- (c) Here P is a 2×3 matrix over \mathbb{Q} . We have $PA = PI_2 = P$ and $BP = I_3 P = P$ so the condition $PA = BP$ is automatically satisfied. Thus the homomorphisms from M_A to M_B are all the 2×3 matrices over \mathbb{Q} .

Exercise 6.2:

- (a) The homomorphisms correspond to matrices $C = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ with $CA = BC$, or equivalently $\begin{bmatrix} a & 2a+2b \\ c & 2c+2d \end{bmatrix} = \begin{bmatrix} a+c & b+d \\ a+c & b+d \end{bmatrix}$, so

$$\begin{aligned} a &= a + c \\ 2a + 2b &= b + d \\ c &= a + c \\ 2c + 2d &= b + d. \end{aligned}$$

The first and third equations give $a = c = 0$, and the remaining equations then give $d = b$, so C must have the form $\begin{bmatrix} 0 & b \\ 0 & b \end{bmatrix}$.

- (b) We need to find the 3×3 matrices with $CA = BC$. Note that $BC = C$. Put $D = A - I = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$. The condition $CA = BC$ now becomes $C(I + D) = C$, or equivalently $CD = 0$. If the columns of C are u, v and w , then the columns of CD are easily seen to be v, w and 0 . Thus $CD = 0$ iff

$v = w = 0$, so all the nonzero entries of C must be in the first column. Thus the homomorphisms from M_A to M_B are the matrices of the form $\begin{bmatrix} a & 0 & 0 \\ b & 0 & 0 \\ c & 0 & 0 \end{bmatrix}$.

(c) Here we need the matrices $C = \begin{bmatrix} c_{11} & c_{21} & c_{31} \\ c_{12} & c_{22} & c_{32} \end{bmatrix}$ such that

$$\begin{bmatrix} c_{11} & c_{21} & c_{31} \\ c_{12} & c_{22} & c_{32} \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} = \begin{bmatrix} \mu_1 & 0 \\ 0 & \mu_2 \end{bmatrix} \begin{bmatrix} c_{11} & c_{21} & c_{31} \\ c_{12} & c_{22} & c_{32} \end{bmatrix},$$

or equivalently

$$\begin{bmatrix} (\lambda_1 - \mu_1)c_{11} & (\lambda_2 - \mu_1)c_{21} & (\lambda_3 - \mu_1)c_{31} \\ (\lambda_1 - \mu_2)c_{11} & (\lambda_2 - \mu_2)c_{21} & (\lambda_3 - \mu_2)c_{31} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

For most choices of numbers $\lambda_1, \lambda_2, \lambda_3, \mu_1, \mu_2$, the λ 's will all be different from the μ 's, so all the numbers $\lambda_i - \mu_j$ will be nonzero. In this case, the only possible matrix C is the zero matrix. In general, if we have $\lambda_i = \mu_j$ for some pairs (i, j) , then the corresponding entries c_{ij} can be nonzero.

Exercise 6.3:

(a) $\text{ann}(4, \mathbb{Z}_{12}) = \{\bar{0}, \bar{3}, \bar{6}, \bar{9}\} \simeq \mathbb{Z}_4$.

(b) We have

$$(x-1)(u, v) = (A-I)(u, v) = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} 0 \\ u \end{bmatrix},$$

so $(x-1)(u, v) = (0, 0)$ iff $u = 0$. Thus $\text{ann}(x-1, M_A) = \{(0, v) \mid v \in \mathbb{R}\}$.

(c) $a\alpha(\bar{1}) = \alpha(a\bar{1}) = \alpha(\bar{a}) = \alpha(\bar{0}) = 0$, because $\bar{a} = \bar{0}$ in R/a .

(d) Suppose that $m \in \text{ann}(a, M)$. We can certainly define a homomorphism $\beta: R \rightarrow M$ by $\beta(x) = xm$. As $am = 0$ this satisfies $\beta(ya) = yam = 0$, so $\beta(x) = 0$ for $x \in Ra$. By the first isomorphism theorem we get an induced map $\alpha = \bar{\beta}: R/aR \rightarrow M$ defined by $\alpha(\bar{x}) = \beta(x) = xm$, and in particular $\alpha(\bar{1}) = m$.

(e) Note that $\text{ann}(D^2 - 1, C^\infty(\mathbb{R}, \mathbb{R}))$ is the space of solutions of the differential equation $f'' = f$, or equivalently the space of functions of the form $f(t) = ue^t + ve^{-t}$ with $u, v \in \mathbb{R}$. For each such function we get a homomorphism $\alpha: \mathbb{R}[D]/(D^2 - 1) \rightarrow C^\infty(\mathbb{R}, \mathbb{R})$ given by

$$\alpha(\overline{a + bD}) = (a + bD)(ue^t + ve^{-t}) = (a + b)ue^t + (a - b)ve^{-t}.$$

Exercise 6.4: Recall that there is a map $\phi: \mathbb{Z}_p \rightarrow \mathbb{Z}_q$ with $\phi(\bar{m}) = \bar{r}m$ iff rp is divisible by q . This is satisfied when $r = p = 3$ and $q = 9$, so α exists. It is also satisfied when $p = 9$ and $r = 1$ and $q = 3$, so β exists.

The elements of \mathbb{Z}_3 are $\bar{0}, \bar{1}$ and $\bar{2}$. We have $\alpha(\bar{0}) = \bar{0}$, $\alpha(\bar{1}) = \bar{3}$, $\alpha(\bar{2}) = \bar{6}$, so the image of α is $\{\bar{0}, \bar{3}, \bar{6}\}$.

Next, the elements of \mathbb{Z}_9 are $\bar{0}, \dots, \bar{8}$. We have $\beta(\bar{3}) = \bar{3}$. The $\bar{3}$ on the left hand side is interpreted as an element of \mathbb{Z}_9 and thus is nonzero, but the $\bar{3}$ on the right hand side is interpreted as an element of \mathbb{Z}_3 and thus is zero. In other words, we have $\beta(\bar{3}) = \bar{0}$. Similarly, we have

$$\beta(\bar{0}) = \beta(\bar{3}) = \beta(\bar{6}) = \bar{0}$$

$$\beta(\bar{1}) = \beta(\bar{4}) = \beta(\bar{7}) = \bar{1}$$

$$\beta(\bar{2}) = \beta(\bar{5}) = \beta(\bar{8}) = \bar{2}.$$

Thus $\ker(\beta) = \{a \in \mathbb{Z}_9 \mid \beta(a) = \bar{0}\} = \{\bar{0}, \bar{3}, \bar{6}\}$. We have shown that $\ker(\beta) = \text{image}(\alpha)$, so the sequence $\mathbb{Z}_3 \xrightarrow{\alpha} \mathbb{Z}_9 \xrightarrow{\beta} \mathbb{Z}_3$ is exact.

Exercise 6.5:

- (a) It is easy to see that $A^2 = 3A$, so for all $v \in M_A$ we have $(x^2 - 3x)v = 0$. In particular, for $u \in \mathbb{C}[x]/(x^2 - 1)$ we have $(x^2 - 3x)\alpha(u) = 0$. However, we also have $(x^2 - 1)u = 0$ and so $(x^2 - 1)\alpha(u) = 0$. The polynomials $x^2 - 3x = x(x - 3)$ and $x^2 - 1 = (x - 1)(x + 1)$ are coprime, so there exist polynomials $p(x)$ and $q(x)$ with $p(x)(x^2 - 3x) + q(x)(x^2 - 1) = 1$. It follows that

$$\alpha(u) = p(x)(x^2 - 3x)\alpha(u) + q(x)(x^2 - 1)\alpha(u) = 0 + 0 = 0.$$

As this holds for all $u \in \mathbb{C}[x]/(x^2 - 1)$, we have $\alpha = 0$ as claimed.

- (b) As $\sin'' = -\sin$ and $\cos'' = -\cos$ we have $(D^2 + 1)V = 0$. As $\sinh'' = \sinh$ and $\cosh'' = \cosh$ we have $(D^2 - 1)W = 0$. The elements $D^2 + 1$ and $D^2 - 1$ are coprime in $\mathbb{R}[D]$, so $\beta = 0$ by the same argument as in part (a).
- (c) We have $\gamma(\bar{1}) = (w, x, y, z) \in \mathbb{Z}^4$ for some $w, x, y, z \in \mathbb{Z}$. As $4\bar{1} = \bar{0}$ we see that $(4w, 4x, 4y, 4z) = \gamma(\bar{0}) = (0, 0, 0, 0)$, so $4w = 4x = 4y = 4z = 0$. As w, x, y and z are just integers, this implies that $w = x = y = z = 0$, so $\gamma(\bar{1}) = 0$. This in turn implies that $\gamma(\bar{n}) = n \cdot \gamma(\bar{1}) = n \cdot 0 = 0$ for all n .

Exercise 6.6: A homomorphism from R^d to M corresponds to a list (m_1, \dots, m_d) of elements of M . There are m possible choices for each entry in the list, so there are m^d possible lists, and thus m^d different homomorphisms from R^d to M .

Exercise 7.1: Let π be as described. Every element of $(L + N)/N$ has the form $z + N$ for some $z \in L + N$. We can write z as $x + y$ for some $x \in L$ and $y \in N$, so $z + N = x + y + N = x + N$ (because $y + N = N$). Thus every element of $(L + N)/N$ has the form $\pi(x)$ for some $x \in L$, which means that π is surjective.

Next, $\ker(\pi)$ is the set of those $x \in L$ for which $x + N = N$, or in other words those $x \in L$ for which we also have $x \in N$, so $\ker(\pi) = L \cap N$.

The First Isomorphism Theorem now tells us that

$$L/(L \cap N) = L/\ker(\pi) \simeq \text{image}(\pi) = (L + N)/N.$$

Exercise 7.2: The homomorphism σ is an isomorphism iff it is both injective and surjective, or equivalently $\ker(\sigma) = \{0\}$ and $\text{image}(\sigma) = M$. We have $\text{image}(\sigma) = M$ iff every element of M can be written in the form $n_0 + n_1$ for some $n_0 \in N_0$ and $n_1 \in N_1$, or equivalently $M = N_0 + N_1$. Next, $\ker(\sigma)$ is the set of pairs $(n, -n)$ where $n \in N_0$ and $-n \in N_1$. However, we have $-n \in N_1$ iff $n \in N_1$, so $\ker(\sigma) = \{(n, -n) \mid n \in N_0 \cap N_1\}$. It follows that $\ker(\sigma) = \{0\}$ iff $N_0 \cap N_1 = \{0\}$. Thus σ is an isomorphism iff $M = N_0 + N_1$ and $N_0 \cap N_1 = \{0\}$, which means precisely that M is the internal direct sum of N_0 and N_1 .

Exercise 8.1:

- (a) Let α be a ring homomorphism from \mathbb{Z}_3 to \mathbb{Z} . By applying α to the equation $\bar{1} + \bar{1} + \bar{1} = \bar{0}$ we obtain $\alpha(\bar{1}) + \alpha(\bar{1}) + \alpha(\bar{1}) = \alpha(\bar{0})$ but $\alpha(\bar{1}) = 1$ and $\alpha(\bar{0}) = 0$ so $1 + 1 + 1 = 0$ in \mathbb{Z} . This is clearly false, so no such α can exist.
- (b) Let α be a ring homomorphism from \mathbb{Q} to \mathbb{Z} . By applying α to the equation $\frac{1}{2} \cdot (1 + 1) = 1$ we get $\alpha(\frac{1}{2}) \cdot (\alpha(1) + \alpha(1)) = \alpha(1)$. We also have $\alpha(1) = 1$ so $\alpha(\frac{1}{2}) \cdot 2 = 1$. However, there is no element $x \in \mathbb{Z}$ with $x \cdot 2 = 1$ so this is impossible, so no such α can exist.
- (c) Let α be a ring homomorphism from \mathbb{C} to \mathbb{R} . By applying α to the equation $i^2 + 1 = 0$ we obtain $\alpha(i)^2 + \alpha(1) = \alpha(0)$ or in other words $\alpha(i)^2 + 1 = 0$. There is no element $x \in \mathbb{R}$ with $x^2 + 1 = 0$, so this is impossible, so no such α can exist.
- (d) The only reasonable example is given by $\alpha(z) = \bar{z}$ (the complex conjugate of z). This is a ring homomorphism because $\overline{z + w} = \bar{z} + \bar{w}$ and $\overline{zw} = \bar{z}\bar{w}$ and $\bar{\bar{z}} = z$. (There are some other examples defined by a bizarre procedure involving heavy set theory. The above example is the only one that is a continuous function from \mathbb{C} to itself.)

Exercise 8.2:

- (a) Define $\alpha: \mathbb{Q}[x] \rightarrow \mathbb{R}$ by $\alpha(f) = f(\sqrt{2})$; this is clearly a ring homomorphism. As $(\sqrt{2})^2 - 2 = 0$ we have $x^2 - 2 \in \ker(\alpha)$, so $\mathbb{Q}[x].(x^2 - 2) \subseteq \ker(\alpha)$. Conversely, suppose that $f \in \ker(\alpha)$, so $f(\sqrt{2}) = 0$. We can divide $f(x)$ by $x^2 - 2$ to get $f(x) = (x^2 - 2)q(x) + a + bx$ for some $a, b \in \mathbb{Q}$. We then have

$$0 = f(\sqrt{2}) = (\sqrt{2}^2 - 2)q(\sqrt{2}) + a + b\sqrt{2} = a + b\sqrt{2}.$$

If $b \neq 0$ we can deduce that $\sqrt{2} = -a/b$ which is impossible as $\sqrt{2}$ is irrational. Thus, we must have $b = 0$, in which case the equation $0 = a + b\sqrt{2}$ tells us that $a = 0$ also. Thus $f(x) = (x^2 - 2)q(x)$, so $f(x) \in \mathbb{Q}[x].(x^2 - 2)$. Thus $\ker(\alpha) = \mathbb{Q}[x].(x^2 - 2)$, so $\mathbb{Q}[x]/(x^2 - 2) \simeq \text{image}(\alpha)$ (by the First Isomorphism Theorem for rings), and $\text{image}(\alpha)$ is a subring of \mathbb{R} as required.

- (b) (i) We have $\alpha(-5) = -5 + I$, and we want to show that this is the same as $i + I$, or in other words that $-5 - i \in I$. By direct calculation we have $(-5 - i)/(2 + 3i) = 1 + i$ which lies in $\mathbb{Z}[i]$, so $-5 - i = (1 + i)(2 + 3i) \in \mathbb{Z}[i].(2 + 3i) = I$ as required.
Now suppose we have an element $a + ib + I \in \mathbb{Z}[i]/I$ (so $a, b \in \mathbb{Z}$). We find that $\alpha(a - 5b) = \alpha(a) + \alpha(b)\alpha(-5) = a + bi + I$, and it follows that α is surjective.
- (ii) Suppose that $n = (2 + 3i)(u + iv)$. By taking norms we find that $n^2 = N(n) = N(2 + 3i)N(u + iv) = 13(u^2 + v^2)$, so n^2 is divisible by 13 in \mathbb{Z} .
- (iii) As $13 = (2 + 3i)(2 - 3i) \in I$ we have $\alpha(13) = 0$ so $13\mathbb{Z} \subseteq \ker(\alpha)$. Conversely, suppose that $\alpha(n) = 0$, so n is divisible by $2 + 3i$ in $\mathbb{Z}[i]$. By (ii) we see that n^2 is divisible by 13, but 13 is prime so n itself must be divisible by 13, so $n \in 13\mathbb{Z}$. Thus $\ker(\alpha) = 13\mathbb{Z}$ and $\mathbb{Z}_{13} = \mathbb{Z}/13\mathbb{Z} \simeq \text{image}(\alpha) = \mathbb{Z}[i]/I$.

Exercise 8.3:

- (a) Define $\alpha: \mathbb{R}[x] \rightarrow \mathbb{C}$ by $\alpha(f) = f(2i)$. This is clearly a ring homomorphism. Any complex number $a + ib$ can be written as $\alpha(a + bx/2)$, so α is surjective. Put

$$I = \ker(\alpha) = \{f \in \mathbb{R}[x] \mid f(2i) = 0\}.$$

The First Isomorphism Theorem for rings now tells us that $\mathbb{R}[x]/I \simeq \mathbb{C}$, so it will be enough to show that $I = \mathbb{R}[x].(x^2 + 4)$. It is clear that the polynomial $f(x) = x^2 + 4$ satisfies $f(2i) = 0$, so $x^2 + 4 \in I$, so $\mathbb{R}[x].(x^2 + 4) \subseteq I$.

Conversely, suppose that $g(x) \in I$, so $g(2i) = 0$. By the division algorithm we have $g(x) = q(x)(x^2 + 4) + ax + b$ for some polynomial $q(x) \in \mathbb{R}[x]$ and some $a, b \in \mathbb{R}$. If we substitute $x = 2i$ in this relation and use the fact that $g(2i) = 0$ we find that $2ai + b = 0$. As a and b are real, we can conclude that $a = b = 0$, so $g(x) = q(x)(x^2 + 4)$, so $g(x) \in \mathbb{R}[x].(x^2 + 4)$. Thus $I = \mathbb{R}[x].(x^2 + 4)$, as required.

- (b) In $\mathbb{R}[x]/(x^2 - 4)$ the elements $\overline{x - 2}$ and $\overline{x + 2}$ are nonzero, but their product is $\overline{x^2 - 4} = \bar{0}$. Thus $\mathbb{R}[x]/(x^2 - 4)$ is not an integral domain, and thus not a field.

Exercise 8.4:

- (a) I claim that

$$R = \{\bar{0}, \bar{1}, \bar{2}, \bar{i}, \overline{1 + i}, \overline{2 + i}, \overline{2i}, \overline{1 + 2i}, \overline{2 + 2i}\},$$

or equivalently $R = \{\overline{a + bi} \mid a, b \in \{0, 1, 2\}\}$. Indeed, the listed elements are certainly contained in R , and it is easy to see that they are all different. Conversely, any element $x \in R$ can be written as $x = \overline{a + ib}$ for some $a, b \in \mathbb{Z}$. We can then write $a = 3c + a'$ for some $c \in \mathbb{Z}$ and $a' \in \{0, 1, 2\}$ and $c \in \mathbb{Z}$. Similarly we have $b = 3d + b'$ for some $d \in \mathbb{Z}$ and $b' \in \{0, 1, 2\}$. It follows that $(a + bi) = (a' + b'i) + 3(c + di)$, so $x = \overline{a' + b'i}$, so x is in our list.

(b) Here we will just write $a + bi$ for $\overline{a + bi}$. We have

$$\begin{aligned} u^0 &= 1 \\ u^1 &= 1 + i \\ u^2 &= (1 + i)^2 = 2i \\ u^3 &= u^2 \cdot u = 2i(1 + i) = 2i - 2 = 2i + 1 \\ u^4 &= (u^2)^2 = -4 = 2 \\ u^5 &= u^4 \cdot u = 2 + 2i \\ u^6 &= u^4 \cdot u^2 = 4i = i \\ u^7 &= u^4 \cdot u^3 = 4i + 2 = i + 2 \\ u^8 &= (u^4)^2 = 4 = 1. \end{aligned}$$

- (c) On comparing (a) with (b) we see that every nonzero element of R is a power of u . We also have $u^8 = 1$, so u^{8-k} is an inverse for u^k , so every nonzero element of the ring R is invertible. This means that R is a field.
- (d) If we can prove that 3 is irreducible in $\mathbb{Z}[i]$, it will follow that $\mathbb{Z}[i]/3$ is a field (Proposition 10.6 in the notes). It is a general fact that prime numbers of the form $4k - 1$ are irreducible in $\mathbb{Z}[i]$, and this obviously covers the case of the prime 3. More explicitly, if 3 were reducible we would have $3 = rs$ for some nonunits r and s . We would then have $9 = N(3) = N(r)N(s)$, and $N(r), N(s) \neq 1$ as r and s are not units. This means we must have $N(r) = N(s) = 3$. However, 3 cannot be written as $a^2 + b^2$ for any integers a and b , so we cannot have $N(r) = 3$, so 3 must be irreducible after all.

Exercise 9.1: Let h be the gcd of f and g , which we can take to be monic. Note that $f(1) = g(1) = 0$, so both f and g are divisible by $x - 1$, so h is divisible by $x - 1$, or equivalently $h(1) = 0$. We claim that this is the only root of h . Indeed, suppose that $h(\zeta) = 0$. As h divides both f and g and $h(\zeta) = 0$, we see that $f(\zeta) = g(\zeta) = 0$, so $\zeta^n = \zeta^m = 1$. We are also given that n and m are coprime, so $nu + mv = 1$ for some integers u and v . We deduce that

$$\zeta = \zeta^1 = \zeta^{nu+mv} = (\zeta^n)^u (\zeta^m)^v = 1^u 1^v = 1,$$

so $\zeta = 1$ as claimed. As this is the only root of h , we see that $h(x) = (x - 1)^k$ for some $k > 0$. To show that $k = 1$, it will suffice to check that f and g are not divisible by $(x - 1)^2$, or equivalently that $f'(1) \neq 0 \neq g'(1)$. This is clear from the formulae: we have $f'(x) = nx^{n-1}$, so $f'(1) = n > 0$, and similarly $g'(1) = m > 0$.

Exercise 9.2: We can write $a = p^n r/s$ for some integer $n \geq 0$ and some integers r, s that are not divisible by p . Similarly, we can write $b = p^m t/u$ for some integer $m \geq 0$ and some integers t, u that are not divisible by p . If $n \leq m$ then the number $x = b/a = p^{m-n} ts/ru$ lies in $\mathbb{Z}_{(p)}$ and $b = ax$. This says that b is divisible by a , and it follows easily that a is a gcd of a and b . Similarly, if $n \geq m$ then b is a gcd of a and b .

Exercise 11.1:

- (a) Put $u = (1, 1, 0)$ and $v = (0, 1, 1)$ and $w = (0, 0, 5)$. I claim that these vectors form a basis for M_0 over \mathbb{Z} . Indeed, it is easy to see that u, v and w all lie in M_0 . Moreover, if $m = (x, y, z) \in M_0$ then $x - y + z = 5t$ for some t , so $z = 5t - x + y$, and one checks that

$$\begin{aligned} xu + (y - x)v + tw &= (x, x, 0) + (0, y - x, y - x) + (0, 0, 5t) \\ &= (x, y, 5t + y - x) = (x, y, z) = m. \end{aligned}$$

This shows that m lies in the submodule generated by u, v and w , and it is clear that u, v and w are linearly independent over \mathbb{Z} , so they form a basis as claimed.

- (b) Here we put $u = (2, 0, 0)$ and $v = (3, 3, 0)$ and $w = (1, 1, 1)$; these are easily seen to be elements of M_1 . Given an arbitrary element $m = (x, y, z) \in M_1$, we note that $x - y = 2s$ and $y - z = 3t$ for some integers s, t . It follows that

$$\begin{aligned} su + tv + zw &= (2s, 0, 0) + (3t, 3t, 0) + (z, z, z) \\ &= (x - y, 0, 0) + (y - z, y - z, 0) + (z, z, z) \\ &= (x, y, z) = m. \end{aligned}$$

This shows that m lies in the submodule generated by u, v and w , and it is clear that u, v and w are linearly independent over \mathbb{Z} , so they form a basis as claimed.

- (c) Here we put $u = (5, -2, 0)$ and $v = (0, 2, -3)$; these are easily seen to be elements of M_2 . Now consider an arbitrary element $m = (x, y, z) \in M_2$, so $6x + 15y + 10z = 0$. We can reduce this equation modulo 5: as $6x = x \pmod{5}$ and $15y = 10z = 0 \pmod{5}$, we deduce that $x = 0 \pmod{5}$. Similarly, we can reduce modulo 2 to show that $y = 0 \pmod{2}$, and reduce mod 3 to see that $z = 0 \pmod{3}$. We thus have $(x, y, z) = (5r, 2s, 3t)$ for some $r, s, t \in \mathbb{Z}$. The equation $6x + 15y + 10z = 0$ now gives $30r + 30s + 30t = 0$, so $r + s + t = 0$. It follows that

$$\begin{aligned} ru - tv &= (5r, -2r, 0) - (0, 2t, -3t) \\ &= (5r, -2(t + r), 3t) \\ &= (5r, 2s, 3t) = (x, y, z) = m. \end{aligned}$$

This shows that m lies in the submodule generated by u and v , and it is clear that u, v and w are linearly independent over \mathbb{Z} , so they form a basis as claimed.

Exercise 11.2: We can define $\alpha: R^n \rightarrow R/d_1 \oplus \dots \oplus R/d_n$ by

$$\alpha(a_1, \dots, a_n) = (a_1 + Rd_1, \dots, a_n + Rd_n).$$

Suppose we have an element $(u_1, \dots, u_n) \in R/d_1 \oplus \dots \oplus R/d_n$, so $u_i \in R/d_i$ for $i = 1, \dots, n$. For each i we can choose $a_i \in R$ such that $u_i = a_i + Rd_i$, and then $\alpha(a_1, \dots, a_n) = (u_1, \dots, u_n)$. Thus α is surjective, and the First Isomorphism Theorem now tells us that $R^n / \ker(\alpha) \simeq R/d_1 \oplus \dots \oplus R/d_n$. We now need to determine the kernel of α . If $\alpha(a_1, \dots, a_n) = 0$ then $a_i + Rd_i$ must be the zero element of R/d_i for all i . This means that $a_i \in Rd_i$, so $a_i = b_i d_i$ say. It follows that $\underline{a} = \sum_i a_i e_i = \sum_i b_i \cdot (d_i e_i) \in N$. Conversely, it is clear that $\alpha(d_i e_i) = 0$ for all i , so that $N \subseteq \ker(\alpha)$, so $\ker(\alpha) = N$. Thus $R^n / N \simeq R/d_1 \oplus \dots \oplus R/d_n$ as claimed.

Exercise 11.3: One solution is as follows:

$$\begin{array}{ll} u_1 = (1, 1, 1, 1) & d_1 = 1 \\ u_2 = (0, 0, 0, 2) & d_1 = 2 \\ u_3 = (0, 1, 1, 0) & d_1 = 4 \\ u_4 = (0, 0, 1, 1) & d_1 = 4. \end{array}$$

Put $v_i = d_i u_i$. It is clear that the set $\{u_1, \dots, u_4\}$ is linearly independent, as is the set $\{v_1, \dots, v_4\}$. It will thus be enough to show that the elements u_i generate F , and the elements v_i generate G .

It is clear that the elements u_i all lie in F . Suppose we have an element $f = (w, x, y, z) \in F$, so $w + x + y + z = 2t$ for some integer t . We then have

$$\begin{aligned} wu_1 + (t - w - x)u_2 + (x - w)u_3 + (y - x)u_4 &= (w, w, w, w) + (0, 0, 0, 2t - 2w - 2x) + \\ &\quad (0, x - w, x - w, 0) + (0, 0, y - x, y - x) \\ &= (w, x, y, 2t - w - x - y) = (w, x, y, z). \end{aligned}$$

Thus f lies in the \mathbb{Z} -submodule generated by $\{u_1, \dots, u_4\}$, as required.

Next, we have

$$\begin{aligned}v_1 &= (1, 1, 1, 1) \\v_2 &= (0, 0, 0, 4) \\v_3 &= (0, 4, 4, 0) \\v_4 &= (0, 0, 4, 4).\end{aligned}$$

These vectors clearly lie in G . Suppose we have an element $g = (w, x, y, z) \in G$, so $w - x = 4r$, $x - y = 4s$ and $y - z = 4t$ for some integers r, s, t . We then have $x = w - 4r$ and $y = x - 4s = w - 4r - 4s$ and $z = y - 4t = w - 4r - 4s - 4t$, so $g = (w, w - 4r, w - 4r - 4s, w - 4r - 4s - 4t)$. From this we see directly that $g = wv_1 - (t+r)v_2 + rv_3 + sv_4$. Thus f lies in the \mathbb{Z} -submodule generated by $\{u_1, \dots, u_4\}$, as required.

We now deduce that

$$\frac{F}{G} \simeq \frac{\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}}{\mathbb{Z} \oplus 2\mathbb{Z} \oplus 4\mathbb{Z} \oplus 4\mathbb{Z}} \simeq \mathbb{Z}_2 \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_4.$$

Exercise 12.1:

$$\begin{aligned}\begin{bmatrix} x-1 & x & x+1 \\ x & 0 & x \\ x+1 & x & x-1 \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} -1 & x & 1 \\ x & 0 & x \\ 1 & x & -1 \end{bmatrix} \\ &\xrightarrow{2} \begin{bmatrix} 1 & 0 & 0 \\ -x & x^2 & 2x \\ -1 & 2x & 0 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 1 & 0 & 0 \\ -x & 2x & 0 \\ -1 & 0 & 2x \end{bmatrix} \\ &\xrightarrow{4} \begin{bmatrix} 1 & 0 & 0 \\ 0 & x & 0 \\ -1 & 0 & x \end{bmatrix}.\end{aligned}$$

We will refer to the three columns as C_1, C_2 and C_3 . In step 1 we subtracted C_2 from C_1 and from C_3 columns. In step 2 we multiplied C_1 by -1 , then subtracted $x C_1$ from C_2 and subtracted C_1 from C_3 . In step 3 we exchanged C_2 and C_3 , and then subtracted $\frac{1}{2}x C_2$ from C_3 . This gives us a matrix in column echelon form. In step 4 we tidy up a little further by multiplying C_2 and C_3 by $\frac{1}{2}$ and then add C_2 to C_1 .

Exercise 12.2:

- (a) In step 1 we add x times the middle row to the top row, and then multiply the middle row by -1 . In step 2 we add x times the first column to the middle column, and subtract the first column from the last column. In step 3 we subtract x^2 times the bottom row from the top row. Finally, in step 4 we add $1 + x$ times the middle column to the last column, and then move the top row down to the bottom.

$$\begin{aligned}\begin{bmatrix} x & 0 & -1 \\ -1 & x & -1 \\ 0 & -1 & x+1 \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} 0 & x^2 & -1-x \\ 1 & -x & 1 \\ 0 & 1 & -1-x \end{bmatrix} \xrightarrow{2} \begin{bmatrix} 0 & x^2 & -1-x \\ 1 & 0 & 0 \\ 0 & 1 & -1-x \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 0 & 0 & x^3+x^2-x-1 \\ 1 & 0 & 0 \\ 0 & 1 & -1-x \end{bmatrix} \xrightarrow{4} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & x^3+x^2-x-1 & 0 \end{bmatrix}.\end{aligned}$$

This gives us a matrix in normal form.

- (b) It follows that

$$M \simeq \mathbb{C}[x]/(x^3 + x^2 - x - 1).$$

- (c) We have $x^3 + x^2 - x - 1 = (x^2 - 1)(x + 1) = (x - 1)(x + 1)^2$. As $x - 1$ and $x + 1$ are coprime, The Chinese Remainder Theorem implies that

$$M \simeq \mathbb{C}[x]/(x^3 + x^2 - x - 1) \simeq \mathbb{C}[x]/(x - 1) \oplus \mathbb{C}[x]/(x + 1)^2 = B(1, 1) \oplus B(-1, 2).$$

Exercise 12.3:

(a)

$$\begin{aligned} \begin{bmatrix} x & 0 & 1 \\ 1 & x & 0 \\ 0 & 1 & x \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} 0 & 0 & 1 \\ 1 & x & 0 \\ -x^2 & 1 & x \end{bmatrix} \\ &\xrightarrow{2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & x \\ 0 & -x^2 & 1 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -x^2 & 1 + x^3 \end{bmatrix} \\ &\xrightarrow{4} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 + x^3 \end{bmatrix}. \end{aligned}$$

We refer to the rows as R_1, R_2 and R_3 and to the columns as C_1, C_2 and C_3 .

In step 1 we subtract xC_3 from C_1 . In step 2 we move C_3 to the front and then subtract xR_1 from R_3 . In step 3 we subtract x^2C_2 from C_3 . In step 4 we add x^2R_2 to R_3 .

- (b) It follows from (a) that $M \simeq R/1 \oplus R/1 \oplus R/(1 + x^3)$. As $R/1 = \{0\}$ it follows that $M \simeq R/(1 + x^3)$, so we may take $f(x) = 1 + x^3$.

Exercise 12.4:

(a)

$$\begin{aligned} \begin{bmatrix} x & 0 & 0 \\ 0 & x - 1 & 0 \\ 0 & 0 & x^2 - x \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} x & x - 1 & 0 \\ 0 & x - 1 & 0 \\ 0 & 0 & x^2 - x \end{bmatrix} \xrightarrow{2} \begin{bmatrix} 1 & x - 1 & 0 \\ 1 - x & x - 1 & 0 \\ 0 & 0 & x^2 - x \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 1 & 0 & 0 \\ 1 - x & x^2 - x & 0 \\ 0 & 0 & x^2 - x \end{bmatrix} \xrightarrow{4} \begin{bmatrix} 1 & 0 & 0 \\ 0 & x^2 - x & 0 \\ 0 & 0 & x^2 - x \end{bmatrix}. \end{aligned}$$

In step 1 we add R_2 to R_1 , and in step 2 we subtract C_2 from C_1 . In step 3 we subtract $(x - 1)C_1$ from C_2 , and in step 4 we subtract $(1 - x)R_1$ from R_2 .

- (b) It follows that $M \simeq \mathbb{C}[x]/(x^2 - x) \oplus \mathbb{C}[x]/(x^2 - x)$. (As usual, we have omitted the factor of $\mathbb{C}[x]/1$, because $\mathbb{C}[x]/1 = \{0\}$.)

Exercise 12.5: We need to reduce A to normal form by row and column operations. We start by subtracting multiples of the first column from the other columns to clear out the entries in the top row. We then subtract the first row from each of the other rows to clear out the first column. Finally, we multiply all the rows except the first one by -1 . This leaves the following matrix:

$$\left(\begin{array}{c|cccc} 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 2 & 2 & 2 & 4 \\ 0 & 0 & 0 & 4 & 4 \\ 0 & 0 & 3 & 3 & 3 \\ 0 & 2 & 0 & 4 & 2 \end{array} \right)$$

From here onwards, we can ignore the first row and column and just work with the remaining 4×4 block. This can be reduced as follows:

$$\begin{bmatrix} 2 & 2 & 2 & 4 \\ 0 & 0 & 4 & 4 \\ 0 & 3 & 3 & 3 \\ 2 & 0 & 4 & 2 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 1 & 2 & -1 & -1 \\ 4 & 0 & 0 & 4 \\ 3 & 0 & 3 & 3 \\ 2 & 2 & 0 & 4 \end{bmatrix} \xrightarrow{2} \left(\begin{array}{c|ccc} 1 & 0 & 0 & 0 \\ \hline 0 & -8 & 4 & 8 \\ 0 & -6 & 6 & 6 \\ 0 & -2 & 2 & 6 \end{array} \right)$$

In step 1 we subtracted the third row from the first row to get $4 - 3 = 1$ in the top right corner, then moved the last column to the front so as to get a 1 in the top left corner. In step 2 we subtracted multiples of the first column from the remaining columns to clear out the top row, and then cleared out the first column.

From here onwards we can ignore the first row and column and just work with the remaining 3×3 block. This can be reduced as follows:

$$\begin{bmatrix} -8 & 4 & 8 \\ -6 & 6 & 6 \\ -2 & 2 & 6 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 2 & -2 & 6 \\ -8 & 4 & 8 \\ -6 & 6 & 6 \end{bmatrix} \xrightarrow{2} \begin{bmatrix} 2 & 0 & 0 \\ 0 & -4 & -16 \\ 0 & 0 & -12 \end{bmatrix} \xrightarrow{3} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 12 \end{bmatrix}.$$

In step 1, we moved the bottom row to the top and multiplied it by -1 . In step 2 we added multiples of the first column to the other columns to clear out the top row, then cleared out the first column. In step 3 we subtracted 4 times the middle column from the last column, and then multiplied both these columns by -1 .

The conclusion is that $\mathbb{Z}^5/M \simeq \mathbb{Z}_2 \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_{12}$.

Exercise 12.6:

$$\begin{bmatrix} 71 & 97 & 113 & 149 \\ 1 & 1 & 1 & 1 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 71 & 26 & 42 & 7 \\ 1 & 0 & 0 & -1 \end{bmatrix} \\ \xrightarrow{2} \begin{bmatrix} 1 & 5 & 0 & 7 \\ 11 & 3 & 6 & -1 \end{bmatrix} \\ \xrightarrow{3} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 11 & -52 & 6 & -78 \end{bmatrix} \\ \xrightarrow{4} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 11 & 2 & 6 & 0 \end{bmatrix} \\ \xrightarrow{5} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 11 & 2 & 0 & 0 \end{bmatrix}.$$

We refer to the columns as C_1, \dots, C_4 . In step 1 we subtract C_1 from C_2 and C_3 , and $2C_1$ from C_4 . In step 2 we subtract $10C_4$ from C_1 , $3C_4$ from C_2 and $6C_4$ from C_3 . In step 3 we subtract $5C_1$ from C_2 , and $7C_1$ from C_4 . In step 4 we add $9C_3$ to C_2 and $13C_3$ to C_4 . Finally, in step 5 we subtract $3C_2$ from C_3 .

Exercise 12.7:

- (a) If we add one row of a matrix to another row, then the determinant is unchanged. If we swap two rows, then the determinant just changes sign. The only invertible elements in \mathbb{Z} are 1 and -1 . If we multiply a row by one of these invertible elements, then the determinant is either unchanged or just changes sign. Similar remarks apply to column operations. Thus, if B is obtained from A by a sequence of row and column operations then $\det(B) = \pm \det(A)$ and so $|\det(B)| = |\det(A)|$.
- (b) Let A be an $n \times n$ matrix over \mathbb{Z} , and suppose that $|\det(A)| = d \neq 0$. Let B be a matrix in normal form obtained from A by row and column operations, so B has the form

$$B = \left(\begin{array}{c|ccc} D & & & 0_{(n-r) \times r} \\ \hline 0_{r \times (n-r)} & & & 0_{(n-r) \times (n-r)} \end{array} \right),$$

where D is a diagonal matrix with nonzero diagonal entries d_1, \dots, d_r say. By multiplying some rows by -1 if necessary, we may assume that $d_i > 0$ for all i . By part (a) we have $|\det(B)| = |\det(A)| = d > 0$. If we had $r < n$ then the last $n - r$ columns in B would be zero and we would have $\det(B) = 0$, giving a contradiction. We must thus have $r = n$ and so $B = D$. As D is a diagonal

matrix we have $d = |\det(B)| = |\det(D)| = d_1 d_2 \dots d_r$. We also know that $M \simeq \mathbb{Z}_{d_1} \oplus \dots \oplus \mathbb{Z}_{d_r}$ and thus $|M| = |\mathbb{Z}_{d_1}| \dots |\mathbb{Z}_{d_r}| = d_1 \dots d_r$. Thus $|M| = d$ as claimed.

- (c) Let B be a matrix in normal form obtained from A by row and column operations. Just as in (a) we see that $\det(B)$ is an invertible element of $\mathbb{C}[x]$ times $\det(A)$. The invertible elements of $\mathbb{C}[x]$ are the nonzero constants, and it follows that the degree of $\det(B)$ is the same as the degree of $\det(A)$.

In particular, we see that $\det(B) \neq 0$ so B cannot contain blocks of zeros, so it must just be a diagonal matrix with nonzero entries $g_1(x), \dots, g_n(x)$ say. Let m_i be the degree of g_i . Then $\det(B) = g_1 \dots g_n$ so $\deg(f) = \deg(\det(B)) = m_1 + \dots + m_n$.

On the other hand, we also have $M \simeq \mathbb{C}[x]/g_1 \oplus \dots \oplus \mathbb{C}[x]/g_n$, so $\dim_{\mathbb{C}}(M) = \sum_{i=1}^n \dim_{\mathbb{C}}(\mathbb{C}[x]/g_i)$. Using the division algorithm we see that every element of $\mathbb{C}[x]/g_i$ has the form $a_0 + \dots + a_{m_i-1}x^{m_i-1} + \mathbb{C}[x]g_i(x)$ for some unique list of coefficients a_0, \dots, a_{m_i-1} . This means that $\{1, x, \dots, x^{m_i-1}\}$ is a basis for $\mathbb{C}[x]/g_i$, so $\dim_{\mathbb{C}}(\mathbb{C}[x]/g_i) = m_i$. It follows that $\dim_{\mathbb{C}}(M) = \sum_{i=1}^n \dim_{\mathbb{C}}(\mathbb{C}[x]/g_i) = \sum_{i=1}^n m_i = \deg(f)$ as claimed.

Exercise 12.8: We have

$$B = \begin{bmatrix} 120 & 60 & 40 \\ 60 & 40 & 30 \\ 40 & 30 & 24 \end{bmatrix}.$$

This can be put in normal form as follows:

$$\begin{aligned} \begin{bmatrix} 120 & 60 & 40 \\ 60 & 40 & 30 \\ 40 & 30 & 24 \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} 40 & 20 & 40 \\ 0 & 10 & 30 \\ -8 & 6 & 24 \end{bmatrix} \xrightarrow{2} \begin{bmatrix} 60 & 20 & -40 \\ 10 & 10 & -10 \\ -2 & 6 & 0 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} 60 & 200 & -40 \\ 10 & 40 & -10 \\ -2 & 0 & 0 \end{bmatrix} \xrightarrow{4} \begin{bmatrix} 0 & 200 & 40 \\ 0 & 40 & 10 \\ 2 & 0 & 0 \end{bmatrix} \\ &\xrightarrow{5} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 40 & 10 \\ 0 & 200 & 40 \end{bmatrix} \xrightarrow{6} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 10 & 40 \\ 0 & 40 & 200 \end{bmatrix} \\ &\xrightarrow{7} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 40 & 40 \end{bmatrix} \xrightarrow{8} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & 40 \end{bmatrix}. \end{aligned}$$

In step 1 we subtract $2C_3$ from C_1 and C_3 from C_2 . In step 2 we add C_2 to C_1 . In step 3 we add $3C_1$ to C_2 . In step 4 we multiply C_3 and R_3 by -1 and then subtract $30R_3$ from R_1 and $5R_3$ from R_2 . In step 5 we swap R_1 and R_3 , and in step 6 we swap C_2 and C_3 . In step 7 we subtract $4C_2$ from C_3 , and in step 8 we subtract $4R_2$ from R_3 .

The conclusion is that $M \simeq \mathbb{Z}_2 \oplus \mathbb{Z}_{10} \oplus \mathbb{Z}_{40}$.

Exercise 12.9: The generators of M are the columns of the matrix $\begin{bmatrix} 2 & 3 & 4 & 5 \\ 8 & 27 & 64 & 125 \end{bmatrix}$. We can perform column operations to simplify this matrix as follows:

$$\begin{aligned} \begin{bmatrix} 2 & 3 & 4 & 5 \\ 8 & 27 & 64 & 125 \end{bmatrix} &\rightarrow \begin{bmatrix} 2 & 1 & 0 & 1 \\ 8 & 19 & 48 & 109 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & 1 \\ 19 & 8 & 48 & 109 \end{bmatrix} \\ &\rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 19 & -30 & 48 & 90 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 19 & 30 & 18 & 0 \end{bmatrix} \\ &\rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 19 & 12 & 18 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 6 & 0 & 0 \end{bmatrix}. \end{aligned}$$

It follows that M is generated by the vectors $(1, 1)$ and $(0, 6)$, and these elements are clearly linearly independent, so they form a basis for M over \mathbb{Z} . Note also that $(1, 1)$ and $(0, 1)$ form a basis for \mathbb{Z}^2 ; using this, it is easy to see that $\mathbb{Z}^2/M \simeq \mathbb{Z}_6$, and so $|\mathbb{Z}^2/M| = 6$.

Exercise 12.10:

- (a) Note that A is not in normal form to start with, because 10 and 15 are not divisible by 6. The first step is to bring some numbers that are not divisible by 6 up onto the top row. It is convenient to add both the second and third rows to the first row. This has the effect that the greatest common divisor of the numbers on the new first row is 1. We can then perform column operations until we have a 1 in the top right corner:

$$\begin{bmatrix} 6 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & 15 \end{bmatrix} \rightarrow \begin{bmatrix} 6 & 10 & 15 \\ 0 & 10 & 0 \\ 0 & 0 & 15 \end{bmatrix} \rightarrow \begin{bmatrix} 6 & -2 & 3 \\ 0 & 10 & 0 \\ 0 & 0 & 15 \end{bmatrix} \rightarrow \begin{bmatrix} 6 & -2 & 1 \\ 0 & 10 & 10 \\ 0 & 0 & 15 \end{bmatrix}.$$

We then subtract multiples of the first row from the other rows, to clear out the entries underneath the 1. After that, we perform column operations to clear out the entries to the left of the 1. We then exchange the first and last columns to put the 1 in the top left corner.

$$\begin{bmatrix} 6 & -2 & 1 \\ 0 & 10 & 10 \\ 0 & 0 & 15 \end{bmatrix} \rightarrow \begin{bmatrix} 6 & -2 & 1 \\ -60 & 30 & 0 \\ -90 & 30 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 0 & 1 \\ -60 & 30 & 0 \\ -90 & 30 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 30 & -60 \\ 0 & 30 & -90 \end{bmatrix}.$$

We now add twice the middle column to the last column, subtract the middle row from the last row, and multiply the last row by -1 :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 30 & -60 \\ 0 & 30 & -90 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 30 & 0 \\ 0 & 30 & -30 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 30 & 0 \\ 0 & 0 & -30 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 30 & 0 \\ 0 & 0 & 30 \end{bmatrix}.$$

The last matrix (which we will call B) is in normal form.

- (b) As B is obtained from A by row and column operations, the quotient of \mathbb{Z}^3 by the span of the columns of A is isomorphic to the quotient of \mathbb{Z}^3 by the span of the columns of B . In other words,

$$\frac{\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}}{6\mathbb{Z} \oplus 10\mathbb{Z} \oplus 15\mathbb{Z}} \simeq \frac{\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}}{1\mathbb{Z} \oplus 30\mathbb{Z} \oplus 30\mathbb{Z}},$$

or equivalently $\mathbb{Z}_6 \oplus \mathbb{Z}_{10} \oplus \mathbb{Z}_{15} \simeq \mathbb{Z}_{30} \oplus \mathbb{Z}_{30}$. (We have omitted the factor of \mathbb{Z}_1 , because $\mathbb{Z}_1 = \{0\}$.)

- (c) The Chinese Remainder Theorem implies that

$$\begin{aligned} \mathbb{Z}_6 &\simeq \mathbb{Z}_2 \oplus \mathbb{Z}_3 \\ \mathbb{Z}_{10} &\simeq \mathbb{Z}_2 \oplus \mathbb{Z}_5 \\ \mathbb{Z}_{15} &\simeq \mathbb{Z}_3 \oplus \mathbb{Z}_5 \\ \mathbb{Z}_{30} &\simeq \mathbb{Z}_2 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_5, \end{aligned}$$

so

$$\begin{aligned} \mathbb{Z}_6 \oplus \mathbb{Z}_{10} \oplus \mathbb{Z}_{15} &\simeq (\mathbb{Z}_2 \oplus \mathbb{Z}_3) \oplus (\mathbb{Z}_2 \oplus \mathbb{Z}_5) \oplus (\mathbb{Z}_3 \oplus \mathbb{Z}_5) \\ &\simeq (\mathbb{Z}_2 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_5) \oplus (\mathbb{Z}_2 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_5) \\ &\simeq \mathbb{Z}_{30} \oplus \mathbb{Z}_{30}. \end{aligned}$$

Exercise 13.1: We have

$$\begin{aligned} x^4 - x^2 &= (x-1)(x+1)x^2 \\ x^4 - 2x^2 + 1 &= (x-1)^2(x+1)^2 \end{aligned}$$

so

$$\begin{aligned}
\mathbb{C}[x]/(x^4 - x^2) &\simeq \mathbb{C}[x]/(x-1) \oplus \mathbb{C}[x]/(x+1) \oplus \mathbb{C}[x]/x^2 \\
\mathbb{C}[x]/(x^4 - 2x^2 + 1) &\simeq \mathbb{C}[x]/(x-1)^2 \oplus \mathbb{C}[x]/(x+1)^2 \\
M &\simeq \mathbb{C}[x]/(x-1) \oplus \mathbb{C}[x]/(x-1)^2 \oplus \\
&\quad \mathbb{C}[x]/(x+1) \oplus \mathbb{C}[x]/(x+1)^2 \oplus \\
&\quad \mathbb{C}[x]x^2 \\
&= B(1, 1) \oplus B(1, 2) \oplus B(-1, 1) \oplus B(-1, 2) \oplus B(0, 2).
\end{aligned}$$

Exercise 13.2: The elements of M have the form $(\bar{a}, \bar{b}, \bar{c})$ with $\bar{a} \in \mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ and $\bar{b} \in \mathbb{Z}_4 = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}\}$ and $\bar{c} \in \mathbb{Z}_9 = \{\bar{0}, \bar{1}, \dots, \bar{8}\}$. We have

$$\begin{aligned}
F_2^1(M) &= \{(\bar{a}, \bar{b}, \bar{c}) \mid 2\bar{a} = \bar{0}, 2\bar{b} = \bar{0}, 2\bar{c} = \bar{0}\} \\
&= \{(\bar{a}, \bar{b}, \bar{c}) \mid 2|2a, 4|2b \text{ and } 9|2c\}.
\end{aligned}$$

Clearly 2 always divides $2a$, and 4 divides $2b$ iff b is even, and 9 divides $2c$ iff 9 divides c . Thus \bar{a} can be either element of \mathbb{Z}_2 , \bar{b} can be $\bar{0}$ or $\bar{2}$, and \bar{c} must be $\bar{0}$. Thus

$$F_2^1(M) = \{(\bar{0}, \bar{0}, \bar{0}), (\bar{0}, \bar{2}, \bar{0}), (\bar{1}, \bar{0}, \bar{0}), (\bar{1}, \bar{2}, \bar{0})\}.$$

The elements of $F_2^2(M)$ are the elements of $F_2^1(M)$ that have the form $2m$ for some $m \in M$. Thus

$$F_2^2(M) = \{(\bar{0}, \bar{0}, \bar{0}), (\bar{0}, \bar{2}, \bar{0})\}.$$

By similar arguments we have

$$F_3^1(M) = \{(\bar{0}, \bar{0}, \bar{0}), (\bar{0}, \bar{0}, \bar{3}), (\bar{0}, \bar{0}, \bar{6})\}.$$

Exercise 13.3:

- (a) The prime factorisation of 225 is $3^2 5^2$. Thus any group of order 225 is the direct sum of a group of order $3^2 = 9$ and a group of order $5^2 = 25$. The 3-primary part could be \mathbb{Z}_9 or $\mathbb{Z}_3 \oplus \mathbb{Z}_3$, and the 5-primary part could be \mathbb{Z}_{25} or $\mathbb{Z}_5 \oplus \mathbb{Z}_5$. The possibilities are:

$$M_1 = \mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_5 \oplus \mathbb{Z}_5$$

$$M_2 = \mathbb{Z}_9 \oplus \mathbb{Z}_5 \oplus \mathbb{Z}_5$$

$$M_3 = \mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_{25}$$

$$M_4 = \mathbb{Z}_9 \oplus \mathbb{Z}_{25}.$$

- (b) As 9 and 25 are coprime, The Chinese Remainder Theorem says that $\mathbb{Z}_{225} = \mathbb{Z}/(9 \times 25) \simeq \mathbb{Z}_9 \oplus \mathbb{Z}_{25} = M_4$.
- (c) Only the groups M_3 and M_4 have any elements of order 25. The group M_4 has only 3 elements satisfying $3x = 0$, so we must have $M \simeq M_3$.

Exercise 13.4:

- (a) Any Abelian group M of order p^4 can be written as a direct sum of groups of the form \mathbb{Z}_{p^k} with $1 \leq k \leq 4$. If $M \simeq \mathbb{Z}_{p^{k_1}} \oplus \dots \oplus \mathbb{Z}_{p^{k_r}}$ then

$$p^4 = |M| = p^{k_1} \times \dots \times p^{k_r} = p^{k_1 + \dots + k_r},$$

so $k_1 + \dots + k_r = 4$. As each k_i is at least 1 this means that $r \leq 4$. Using this and some trial and error we see that the possibilities are as follows:

$$\begin{aligned} M_1 &= \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \\ M_2 &= \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_{p^2} \\ M_3 &= \mathbb{Z}_p \oplus \mathbb{Z}_{p^3} \\ M_4 &= \mathbb{Z}_{p^2} \oplus \mathbb{Z}_{p^2} \\ M_5 &= \mathbb{Z}_{p^4}. \end{aligned}$$

- (b) In the usual notation, the question tells us that $f_p^1(M) = 3$. It is a standard fact that $f_p^1(\mathbb{Z}_{p^k}) = 1$ for all $k \geq 1$, and that $f_p^1(A \oplus B) = f_p^1(A) + f_p^1(B)$. Using this, we find that

$$\begin{aligned} f_p^1(M_1) &= 4 \\ f_p^1(M_2) &= 3 \\ f_p^1(M_3) &= 2 \\ f_p^1(M_4) &= 2 \\ f_p^1(M_5) &= 1. \end{aligned}$$

The only possibility is thus that $M \simeq M_2$.

- (c) We have $10000 = 2^4 \times 5^4$. As 2 and 5 are coprime, any Abelian group of order $2^4 \times 5^4$ is the direct sum of a group of order 2^4 and a group of order 5^4 , in a unique way. By part (a), there are 5 possibilities for the 2-primary part and 5 possibilities for the 5-primary part, giving $5 \times 5 = 25$ possible groups of order 10000.

Exercise 13.5:

- (a) The possibilities are as follows:

$$\begin{aligned} M_1 &= \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \\ M_2 &= \mathbb{Z}_{p^2} \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \\ M_3 &= \mathbb{Z}_{p^2} \oplus \mathbb{Z}_{p^2} \oplus \mathbb{Z}_p \\ M_4 &= \mathbb{Z}_{p^3} \oplus \mathbb{Z}_p \oplus \mathbb{Z}_p \\ M_5 &= \mathbb{Z}_{p^3} \oplus \mathbb{Z}_{p^2} \\ M_6 &= \mathbb{Z}_{p^4} \oplus \mathbb{Z}_p \\ M_7 &= \mathbb{Z}_{p^5}. \end{aligned}$$

- (b) Take $p = 2$. The groups M_4, \dots, M_7 all contain elements of order $p^3 = 8$, but all elements in M have $4m = 0$, so we must have $M \simeq M_i$ for some $i \leq 3$. By assumption we have $|F_2^1(M)| = 8 = 2^3$ so $f_2^1(M) = \dim_{\mathbb{Z}_2} F_2^1(M) = 3$. However, we know that $f_p^k(\mathbb{Z}_{p^j}) = 1$ for all $j > 0$ so $f_2^1(M_1) = 5$, $f_2^1(M_2) = 4$ and $f_2^1(M_3) = 3$. We must therefore have $M \simeq M_3$.

Exercise 13.6: First note that $(D - 1)D(D + 1)f = (D^3 - D)f = f''' - f' = 0$. Suppose we take $e_{-1} = a_{-1}D(D + 1)$ for some $a_{-1} \in \mathbb{R}$. Then $(D - 1)e_{-1}f = a_{-1}(D - 1)D(D + 1)f = 0$, as required. Similarly, if we take $e_0 = a_0(D - 1)(D + 1)$ and $e_1 = a_1(D - 1)D$ then we will have $De_0f = 0$ and $(D + 1)e_1f = 0$.

If we can arrange that $e_{-1} + e_0 + e_1 = 1$ then the remaining condition $f = e_{-1}f + e_0f + e_1f$ will clearly also be satisfied. We have

$$\begin{aligned} e_{-1} + e_0 + e_1 &= a_{-1}(D^2 + D) + a_0(D^2 - 1) + a_1(D^2 - D) \\ &= -a_0 + (a_{-1} - a_1)D + (a_{-1} + a_0 + a_1)D^2. \end{aligned}$$

For this to equal 1, we must have $a_0 = -1$ and $a_1 = a_{-1}$ and $a_{-1} + a_0 + a_1 = 0$, so $a_1 = a_{-1} = 1/2$. Thus

$$\begin{aligned}e_{-1} &= (D^2 + D)/2 \\e_0 &= 1 - D^2 \\e_1 &= (D^2 - D)/2.\end{aligned}$$

Exercise 13.7:

- (a) $p(D) = D^3 - D$.
 (b) Note that $p(D) = (D - 1)(D + 1)D$ and all the factors are coprime to each other and $p(D)M = \{0\}$. It follows that if we put

$$\begin{aligned}M_0 &= \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid (D - 1)f = 0\} = \{f \mid f' = f\} \\M_1 &= \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid (D + 1)f = 0\} = \{f \mid f' = -f\} \\M_2 &= \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid Df = 0\} = \{f \mid f' = 0\}\end{aligned}$$

then $M = M_0 \oplus M_1 \oplus M_2$.

- (c) By standard methods we have

$$M_0 = \{ae^t \mid a \in \mathbb{R}\} \quad M_1 = \{be^{-t} \mid b \in \mathbb{R}\} \quad M_2 = \mathbb{R}.$$

As $M = M_0 \oplus M_1 \oplus M_2$, any function f satisfying $f''' = f'$ can be written as $f = f_0 + f_1 + f_2$ with $f_i \in M_i$, or equivalently as $ae^t + be^{-t} + c$ for some $a, b, c \in \mathbb{R}$.

Exercise 13.8:

- (a) We have $f'(t) = p'(t)e^{\lambda t} + \lambda p(t)e^{\lambda t} = p'(t)e^{\lambda t} + \lambda f(t)$. Thus $((D - \lambda)f)(t) = p'(t)e^{\lambda t}$. It follows inductively that $((D - \lambda)^k f)(t) = p^{(k)}(t)e^{\lambda t}$ for all $k \geq 0$. If p is a polynomial of degree d then $p^{(k)}$ is a polynomial of degree $d - k$ for $k = 1, \dots, d$, and $p^{(k)} = 0$ for $k > d$. It follows that $(D - \lambda)^k f = 0$ for $k > d$. This means that f is a torsion element of W_λ . This holds for every element of W_λ , so W_λ is a torsion module.
 (b) We have

$$g'(t) = f'(t)e^{-\lambda t} - \lambda f(t)e^{-\lambda t} = ((D - \lambda)f)(t)e^{-\lambda t}.$$

By extending this inductively, we find that $g^{(k)}(t) = ((D - \lambda)^k f)(t)e^{-\lambda t} = 0$, or in other words $D^k g = 0$.

- (c) In the case $k = 1$ we have $Dg = 0$ so g is constant and so certainly polynomial of degree less than 1. Suppose we have shown that all functions with $D^{k-1}g = 0$ are polynomial of degree less than $k - 1$. If $D^k f = 0$ then $D^{k-1}f' = 0$ so f' is polynomial, say $f'(t) = a_0 + \dots + a_{k-2}t^{k-2}$. Put

$$F(t) = f(t) - f(0) = \int_{s=0}^t f'(s) ds = a_0 t + \dots + a_{k-2} t^{k-1} / (k-1),$$

which is clearly a polynomial of degree less than k . As $f(0)$ is just a constant we deduce that $f(t) = F(t) + f(0)$ is also a polynomial of degree less than k .

- (d) If $(D - \lambda)^k f = 0$ then the function $g(t) = f(t)e^{-\lambda t}$ satisfies $D^k g = 0$, so g is a polynomial, so $f(t) = g(t)e^{\lambda t}$ is an element of W_λ .
 (e) Suppose that $p(D)f = 0$. We can factor $p(D)$ as $u(D - \lambda_1)^{k_1} \dots (D - \lambda_r)^{k_r}$ for some nonzero constant u , where the λ_i 's are all different and the k_i 's are all strictly positive. It follows that the terms $(x - \lambda_i)^{k_i}$ are all coprime to each other. If we put $V = \{f \mid p(D)f = 0\}$ and $V_i = \{f \mid (D - \lambda_i)^{k_i} f = 0\}$, it follows that $V = V_1 \oplus \dots \oplus V_r$. However, we know from (d) that $V_i \subseteq W_{\lambda_i}$. Thus

$$f \in V_1 + \dots + V_r \subseteq W_{\lambda_1} + \dots + W_{\lambda_r}.$$

It follows immediately from the definitions that $f \in W$.

- (f) Suppose that $f \in \text{tors}(C^\infty(\mathbb{R}, \mathbb{C}))$. Then $p(D)f = 0$ for some nonzero element $p(D) \in \mathbb{C}[D]$, so (e) tells us that $f \in W$. Conversely, if $f \in W$ then we can write $f(t) = p_1(t)e^{\lambda_1 t} + \dots + p_r(t)e^{\lambda_r t}$ for some $\lambda_1, \dots, \lambda_r \in \mathbb{C}$ and polynomials $p_1, \dots, p_r \in \mathbb{C}[t]$. If we put $f_i(t) = p_i(t)e^{\lambda_i t}$ then $f_i \in W_{\lambda_i}$, so f_i is a torsion element by (a). Thus $f = f_1 + \dots + f_r$ is a sum of torsion elements and thus is a torsion element, as claimed. More concretely, if the degree of p_i is $k_i - 1$ then the element $q(D) = (D - \lambda_1)^{k_1} \dots (D - \lambda_r)^{k_r}$ is nonzero and satisfies $q(D)f = 0$.

Exercise 13.9:

- (a) In step 1 we subtract x^2 times the last column from the first column, and subtract x times the last column from the second column. In step 2 we swap the first and last columns. In step 3 we subtract the middle row from the bottom row, and then subtract the top row from the middle row. Finally, we swap the middle row and the bottom row.

$$\begin{aligned} \begin{bmatrix} x^3 & x^2 & x \\ x & x^2 & x \\ x & x & x \end{bmatrix} &\xrightarrow{1} \begin{bmatrix} 0 & 0 & x \\ x - x^3 & 0 & x \\ x - x^3 & x - x^2 & x \end{bmatrix} \xrightarrow{2} \begin{bmatrix} x & 0 & 0 \\ x & 0 & x - x^3 \\ x & x - x^2 & x - x^3 \end{bmatrix} \\ &\xrightarrow{3} \begin{bmatrix} x & 0 & 0 \\ 0 & 0 & x - x^3 \\ 0 & x - x^2 & 0 \end{bmatrix} \xrightarrow{4} \begin{bmatrix} x & 0 & 0 \\ 0 & x - x^2 & 0 \\ 0 & 0 & x - x^3 \end{bmatrix}. \end{aligned}$$

Note that $x - x^2 = x(1 - x)$ and $x - x^3 = x(1 - x)(1 + x)$, so $x|(x - x^2)|(x - x^3)$, so the last matrix is in normal form.

- (b) It follows that

$$M \simeq \mathbb{C}[x]/x \oplus \mathbb{C}[x]/(x - x^2) \oplus \mathbb{C}[x]/(x - x^3).$$

- (c) The Chinese Remainder Theorem implies that

$$\begin{aligned} \mathbb{C}[x]/(x - x^2) &\simeq \mathbb{C}[x]/x \oplus \mathbb{C}[x]/(x - 1) \\ &\simeq B(0, 1) \oplus B(1, 1) \\ \mathbb{C}[x]/(x - x^3) &\simeq \mathbb{C}[x]/x \oplus \mathbb{C}[x]/(x - 1) \oplus \mathbb{C}[x]/(x + 1) \\ &\simeq B(0, 1) \oplus B(1, 1) \oplus B(-1, 1). \end{aligned}$$

Thus

$$\begin{aligned} M &\simeq \mathbb{C}[x]/x \oplus \mathbb{C}[x]/(x - x^2) \oplus \mathbb{C}[x]/(x - x^3) \\ &\simeq B(0, 1) \oplus (B(0, 1) \oplus B(1, 1)) \oplus (B(0, 1) \oplus B(1, 1) \oplus B(-1, 1)) \\ &\simeq B(0, 1)^3 \oplus B(1, 1)^2 \oplus B(-1, 1). \end{aligned}$$

Exercise 14.1: The characteristic polynomial is

$$\begin{aligned} \begin{vmatrix} t & -a & -b \\ a & t & -c \\ b & c & t \end{vmatrix} &= t \begin{vmatrix} t & -c \\ c & t \end{vmatrix} - (-a) \begin{vmatrix} a & -c \\ b & t \end{vmatrix} + (-b) \begin{vmatrix} a & t \\ b & c \end{vmatrix} \\ &= (t^3 + c^2t) + (a^2t + abc) + (-abc + b^2t) \\ &= t^3 + r^2t = t(t + ir)(t - ir). \end{aligned}$$

If $r \neq 0$ then the three roots of $\text{char}(A)$ are distinct. It follows easily that the Jordan normal form is $J(0, 1) \oplus J(ir, 1) \oplus J(-ir, 1)$ and thus that

$$M_A \simeq B(0, 1) \oplus B(ir, 1) \oplus B(-ir, 1) = M_0 \oplus M_{ir} \oplus M_{-ir}.$$

In the exceptional case where $r = 0$ we must have $a = b = c = 0$ so A is just the zero matrix and it is clear that $M_A \simeq M_0 \oplus M_0 \oplus M_0$.

Exercise 14.2:

- (a) The characteristic polynomial is by definition the determinant of the first matrix below. The second matrix is obtained by adding the second and third rows to the top row, and the third matrix is obtained by subtracting the first column from each of the other columns. It is a standard fact that these operations do not change the determinant.

$$\begin{bmatrix} x-u & -v & -w \\ -v & x-w & -u \\ -w & -u & x-v \end{bmatrix} \rightarrow \begin{bmatrix} x-a & x-a & x-a \\ -v & x-w & -u \\ -w & -u & x-v \end{bmatrix} \rightarrow \begin{bmatrix} x-a & 0 & 0 \\ -v & x+(v-w) & v-u \\ -w & w-u & x-(v-w) \end{bmatrix}.$$

The determinant of the last matrix can be expanded out directly to give

$$\begin{aligned} (x-a)(x^2 - (v-w)^2 - (w-u)(v-u)) &= (x-a)(x^2 - v^2 - w^2 + 2vw - vw + uv + uw - u^2) \\ &= (x-a)(x^2 - u^2 - v^2 - w^2 + uv + vw + wu). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} b &= ((u-v)^2 + (v-w)^2 + (w-u)^2)/2 \\ &= (u^2 + v^2 - 2uv + v^2 + w^2 - 2vw + w^2 + u^2 - 2uw)/2 \\ &= (2u^2 + 2v^2 + 2w^2 - 2uv - 2vw - 2wu)/2 \\ &= u^2 + v^2 + w^2 - uv - vw - wu. \end{aligned}$$

Using this, we can rewrite the characteristic polynomial as $(x-a)(x^2 - b)$.

- (b) Suppose that the numbers u , v and w are not all the same, and that $uv + vw + wu \neq 0$. The characteristic polynomial of A is $(x-a)(x-\sqrt{b})(x+\sqrt{b})$; if we can show that the three roots are distinct, it will follow that this is the same as the minimal polynomial.

As u , v and w are not all the same, at least one of the numbers $u-v$, $v-w$ and $w-u$ is nonzero. Thus $(u-v)^2$, $(v-w)^2$ and $(w-u)^2$ are all nonnegative, and one of them is strictly positive, so the number $b = ((u-v)^2 + (v-w)^2 + (w-u)^2)/2$ is strictly positive. This means that $\sqrt{b} > 0$, so $\sqrt{b} \neq -\sqrt{b}$.

We also have

$$a^2 - b = (u+v+w)^2 - (u^2 + v^2 + w^2 - uv - vw - wu) = 3(uv + vw + wu) \neq 0,$$

so $a \neq \pm\sqrt{b}$. Thus, the three roots are all different, as required.

- (c) If $uv + vw + wu = 0$ we see from the above equation that $a^2 = b$, so the characteristic polynomial is $(x-a)(x^2 - a^2) = (x-a)^2(x+a)$. By assumption we have $a \neq 0$ so $x-a$ and $x+a$ are coprime. We know from the general theory that the minimal polynomial divides the characteristic polynomial and has precisely the same roots, so it is either $(x-a)(x+a) = x^2 - a^2$ or $(x-a)(x+a)^2$. By direct calculation one finds that

$$A^2 - a^2I = (uv + vw + wu) \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix},$$

which is the zero matrix because $uv + vw + wu = 0$. Thus the minimal polynomial is $x^2 - a^2$, as claimed.

- (d) If $u = -13$, $v = 11$ and $w = 2$ then $a = -13 + 11 + 2 = 0$ and $b = ((-13 - 11)^2 + (11 - 2)^2 + (2 - (-13))^2)/2 = 441 = 21^2$. The characteristic polynomial is $x(x^2 - 441) = x(x-21)(x+21)$. The three roots are clearly distinct, so this is also the minimal polynomial.
- (e) If $u = -2$, $v = 3$ and $w = 6$ then $a = 7$ and $uv + vw + wu = -6 + 18 - 12 = 0$ so part (c) applies and the minimal polynomial is $x^2 - 49$.

Exercise 14.3:

- (a) The characteristic polynomial is the determinant of the matrix

$$tI - A = \begin{bmatrix} 1+t & -1 & -1 & 1 \\ 0 & 1+t & 0 & -1 \\ 0 & 0 & 1+t & 1 \\ 0 & 0 & 0 & 1+t \end{bmatrix}.$$

As this is an upper-triangular matrix, the determinant is just the product of the diagonal entries, which is $(1+t)^4$.

- (b) The minimal polynomial is a divisor of the characteristic polynomial, so it must be $(1+t)^k$ for some integer $k \leq 4$. The matrix $I + A$ is clearly nonzero, but

$$(I + A)^2 = \begin{bmatrix} 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

It follows that the minimal polynomial of A is $(t+1)^2$.

- (c) The first two rows of $I + A$ are clearly linearly independent, but the third row is minus the second, and the last row is zero. Thus, the rank is two.
- (d) If M_A is a direct sum of modules of the form $B(\lambda, k)$, then the numbers λ must be roots of the minimal polynomial of multiplicity at least k . The only root of the minimal polynomial is -1 , which has multiplicity 2. It follows that the only possible summands in M_A are $B(-1, 1)$ (which has dimension 1 over \mathbb{C}) and $B(-1, 2)$ (which has dimension 2 over \mathbb{C}). As $x+1$ has rank 0 on $B(-1, 1)$ and rank 1 on $B(-1, 2)$, we see that the number of copies of $B(-1, 2)$ is the rank of $x+1$ on M_A , or in other words the rank of $A + I$, which is 2. The two copies of $B(-1, 2)$ have total dimension 4, which is the dimension of M_A , so there cannot be any copies of $B(-1, 1)$ and we have $M_A \simeq B(-1, 2) \oplus B(-1, 2)$.

Exercise 14.4:

- (a) The characteristic polynomial is the determinant of the matrix

$$tI - A = \begin{bmatrix} t-1 & -2 & -3 & -4 \\ 0 & t-1 & 0 & -5 \\ 0 & 0 & t-1 & -6 \\ 0 & 0 & 0 & t-1 \end{bmatrix}.$$

As this is an upper-triangular matrix, the determinant is just the product of the diagonal entries, which is $(t-1)^4$.

- (b) The minimal polynomial is a divisor of the characteristic polynomial, so it must be $(t-1)^k$ for some integer $k \leq 4$. We have

$$(A - I)^2 = \begin{bmatrix} 0 & 2 & 3 & 4 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 2 & 3 & 4 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 28 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \neq 0$$

$$(A - I)^3 = \begin{bmatrix} 0 & 0 & 0 & 28 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 2 & 3 & 4 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

and it follows that the minimal polynomial is $(t-1)^3$.

- (c) As 1 is the only root of the minimal polynomial, the module M_A is isomorphic to a direct sum of modules of the form $B(1, k)$. As 1 is a triple root of the minimal polynomial, we must have at least one summand of the form $B(1, 3)$. This has dimension 3 and M_A has dimension 4 so there is only one dimension left over, so the other summand must be $B(1, 1)$. We thus have $M_A \simeq B(1, 3) \oplus B(1, 1)$.

(d) We read off from this that A is conjugate to $J(1, 3) \oplus J(1, 1)$, which is the following matrix:

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Exercise 14.5:

(a) The characteristic polynomial is the determinant of the matrix

$$tI - A = \begin{bmatrix} t+1 & 0 & -1 & 0 \\ 0 & t-1 & -2 & 0 \\ 0 & 0 & t+1 & 0 \\ 0 & -1 & -1 & t-1 \end{bmatrix}.$$

Expanding along the top row and ignoring the 0's we get

$$\det(tI - A) = (t+1) \begin{vmatrix} t-1 & -2 & 0 \\ 0 & t+1 & 0 \\ -1 & -1 & t-1 \end{vmatrix} + (-1) \begin{vmatrix} 0 & t-1 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & t-1 \end{vmatrix}$$

The second term here is easily seen to be 0. For the first term, we have

$$\begin{aligned} \begin{vmatrix} t-1 & -2 & 0 \\ 0 & t+1 & 0 \\ -1 & -1 & t-1 \end{vmatrix} &= (t-1) \begin{vmatrix} t+1 & 0 \\ -1 & t-1 \end{vmatrix} - (-2) \begin{vmatrix} 0 & 0 \\ -1 & t-1 \end{vmatrix} \\ &= (t-1)^2(t+1) - (-2)(0) = (t-1)^2(t+1). \end{aligned}$$

It follows that $\text{char}(A)(t) = (t-1)^2(t+1)^2$.

A slightly quicker approach is possible if you know that determinants can be expanded along any row or column, not just along the top row. We then have

$$\begin{aligned} \begin{vmatrix} t+1 & 0 & -1 & 0 \\ 0 & t-1 & -2 & 0 \\ 0 & 0 & t+1 & 0 \\ 0 & -1 & -1 & t-1 \end{vmatrix} &\stackrel{1}{=} (t+1) \begin{vmatrix} t-1 & -2 & 0 \\ 0 & t+1 & 0 \\ -1 & -1 & t-1 \end{vmatrix} \\ &\stackrel{2}{=} (t+1)(t-1) \begin{vmatrix} t+1 & 0 \\ -1 & t-1 \end{vmatrix} \\ &= (t+1)^2(t-1)^2, \end{aligned}$$

where in step 1 we have expanded using the first column, and in step 2 we have expanded using the third column.

(b) We have

$$A + I = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 2 & 2 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 2 \end{bmatrix} \quad A - I = \begin{bmatrix} -2 & 0 & 1 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 1 & 1 & 0 \end{bmatrix}.$$

In $A + I$ the nonzero rows are clearly linearly independent, so the rank is three. In $A - I$ the middle two rows are multiples of each other but the first, second and fourth rows are linearly independent, so the rank is again three.

(c) It follows from (a) that the JNF of A contains only blocks of type $J(1, k)$ or $J(-1, k)$. Theorem 14.7(b) tells us that the number of blocks of type $J(1, k)$ is $4 - \text{rank}(A - I)$, which is equal to 1 by (b). We thus have only one block of type $J(1, k)$, which has to contribute a factor $(t-1)^2$ in the characteristic polynomial, so it must be $J(1, 2)$. Similarly, we have only one block of type $J(-1, k)$ and it must be $J(-1, 2)$. Thus the JNF of A is $J(1, 2) \oplus J(-1, 2)$.

(d) It follows from (c) that

$$M_A \simeq B(1, 2) \oplus B(-1, 2) = \mathbb{C}[x]/(x-1)^2 \oplus \mathbb{C}[x]/(x+1)^2 \simeq \mathbb{C}[x]/((x-1)^2(x+1)^2).$$

(The last equality uses the Chinese Remainder Theorem, which is valid because $(x-1)^2$ and $(x+1)^2$ are coprime.) It follows that M_A is cyclic, as claimed.

Exercise 14.6:

(a) $(t-1)^2(t+1)^2$

(b) 3 and 3.

(c) It follows from (a) that the JNF of A contains only blocks of type $J(1, k)$ or $J(-1, k)$. Theorem 14.7(b) tells us that the number of blocks of type $J(1, k)$ is $4 - \text{rank}(A - I)$, which is equal to 1 by (b). We thus have only one block of type $J(1, k)$, which has to contribute a factor $(t-1)^2$ in the characteristic polynomial, so it must be $J(1, 2)$. Similarly, we have only one block of type $J(-1, k)$ and it must be $J(-1, 2)$. Thus the JNF of A is $J(1, 2) \oplus J(-1, 2)$.

(d) It follows from (c) that

$$M_A \simeq B(1, 2) \oplus B(-1, 2) = \mathbb{C}[x]/(x-1)^2 \oplus \mathbb{C}[x]/(x+1)^2 \simeq \mathbb{C}[x]/((x-1)^2(x+1)^2).$$

(The last equality uses the Chinese Remainder Theorem, which is valid because $(x-1)^2$ and $(x+1)^2$ are coprime.) It follows that M_A is cyclic, as claimed.